



CLINICAL GUIDELINES

Preface to the Imaging Guidelines

Version 1.0

Effective February 1, 2021



eviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Imaging requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2020 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Preface to the Imaging Guidelines

Preface-1: Guideline Development	3
Preface-2: Benefits, Coverage Policies, and Eligibility Issues	4
Preface-3: Clinical Information	5
Preface-4: Coding Issues	14
Preface-5: Whole Body Imaging	21
Preface-6: References	24
Preface-7: Copyright Information	25
Preface-8: Trademarks	26

Preface-1: Guideline Development

- The eviCore healthcare (eviCore) evidence-based, proprietary clinical guidelines evaluate a range of advanced imaging and procedures, including NM, US, CT, MRI, PET, and Radiation Oncology, Sleep Studies and Cardiac and Spine interventions.
- eviCore reserves the right to change and update the guidelines. The guidelines undergo a formal review annually. eviCore's guidelines are based upon major national and international association and society guidelines and criteria, peer-reviewed literature, major treatises as well as, input from health plans, practicing academic and community-based physicians.
- These Guidelines are not intended to supersede or replace sound medical judgment, but instead, should facilitate the identification of the most appropriate imaging procedure given the patient's clinical condition. These guidelines are written to cover medical conditions as experienced by the majority of patients. However, these guidelines may not be applicable in certain clinical circumstances, and physician judgment can override the guidelines.
- Clinical decisions, including treatment decisions, are the responsibility of the patient and his/her provider. Clinicians are expected to use independent medical judgment, which takes into account the clinical circumstances to determine patient management decisions.
- eviCore supports the Choosing Wisely initiative (www.choosingwisely.org) by the American Board of Internal Medicine (ABIM) Foundation and many national physician organizations, to reduce the overuse of diagnostic tests that are low value, no value, or whose risks are greater than the benefits.

Preface-2: Benefits, Coverage Policies, and Eligibility Issues

- Benefits, coverage policies, and eligibility issues pertaining to each Health Plan may take precedence over eviCore's guidelines. Providers are urged to obtain written instructions and requirements directly from each payor.

Medicare Coverage Policies

- For Medicare and Medicare Advantage enrollees, the coverage policies of CMS (Centers for Medicare and Medicaid Services) take precedence over eviCore's guidelines.
- CMS requires coverage for studies requested as part of a CMS approved clinical trial through the CMS CED program. A list of the currently approved studies is available at:
 - ◆ <http://www.cms.gov/Medicare/Coverage/Coverage-with-Evidence-Development/Index>

Investigational and Experimental Studies

- Certain imaging studies described in these guidelines are considered investigational by various payers, and their coverage policies may take precedence over eviCore's guidelines. Certain advanced imaging studies, or other procedures, may be considered investigational and experimental if there is a paucity of supporting evidence; if the evidence has not matured to exhibit improved health parameters or; the advanced imaging study/procedure lacks a collective opinion of support.

Clinical and Research Trials

- Similar to investigational and experimental studies, clinical trial imaging requests will be considered to determine whether they meet Health Plan coverage and eviCore's evidence-based guidelines.

Legislative Mandate

- State and federal legislations may need to be considered in the review of advanced imaging requests. For example:
 - ◆ Various State and Federal Breast Density Laws
 - ◆ Texas HB 1290 Coronary Calcium CT Law

Reference

1. Prospective Payment Systems - General Information. CMS. <https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/ProspectivePayment/ProspectivePayment>.
2. Medicare Coverage With Evidence Development: A Policy-Making Tool in Evolution. Journal of Oncology Practice. 2007;3(6):296-301. doi:10.1200/jop.0763501.

Preface-3: Clinical Information

Clinical Documentation and Age Considerations

- eviCore guidelines use an evidence-based approach to determine the most appropriate imaging procedure for each patient, at the most appropriate time in the diagnostic and treatment cycle. eviCore guidelines are framed by:
 - ◆ Clinical presentation of the patient, rather than the studies requested
 - ◆ Adequate clinical information that must be submitted to eviCore in order to establish medical necessity for advanced imaging includes, but is not limited to the following:
 - ◆ Recent clinical evaluation (within 60 days including a recent detailed history, physical examination, and/or laboratory and prior imaging studies.
 - The Spine and Musculoskeletal guidelines require x-ray studies from when the current episode of symptoms has started or changed; x-ray imaging does not have to be within the past 60 days.
 - Advanced imaging should not be ordered prior to clinical evaluation of a patient by the physician treating the individual. This may include referral to a consultant specialist who will make further treatment decisions.
 - Other meaningful contact (telephone or video call, electronic mail or messaging) by an established patient can substitute for an in-person clinical evaluation.
 - An recent clinical evaluation may be unnecessary if the patient is undergoing a guideline-supported, scheduled follow-up imaging evaluation. Exceptions due to routine surveillance indications are addressed in the applicable condition-specific guideline sections.
 - ◆ eviCore’s evidence based approach to determine the most appropriate imaging procedure for each patient requires submission of medical records pertinent to the requested imaging.
- Many conditions affecting the pediatric population are different diagnoses than those occurring in the adult population. For those diseases which occur in both pediatric and adult populations, minor differences may exist in management due to patient age, comorbidities, and differences in disease natural history between children and adults.
 - ◆ Patients who are <18 years old should be imaged according to the Pediatric Imaging Guidelines. Any conditions not specifically discussed in the Pediatric Imaging Guidelines should be imaged according to the General Imaging Guidelines. Patients who are ≥18 years old should be imaged according to the General Imaging Guidelines, except where directed otherwise by a specific guideline section.
- The terms “male” and “female” used in these guidelines refer to anatomic-specific diseases and disease predispositions associated with individuals’ sex assigned at birth rather than their gender identity. It should be noted that gender identity and anatomic-specific diseases as well as disease predispositions are not always linked. As such, these guidelines should be applied to the individual’s corresponding known or suspected anatomic-specific disease or disease predisposition. At eviCore, we

believe that it is important to understand how all individuals, including those who are gender-diverse, choose to identify themselves. To ensure that gender-diverse individuals are treated with respect and that decisions impacting their healthcare are made correctly and with sensitivity, eviCore recognizes all individuals with the following gender marker options: Male, Female, Transgender male, Transgender female, “X”, and “Not specified”.

General Imaging Information

- “Standard” or “conventional” imaging is most often performed in the initial and subsequent evaluations of malignancy. Standard or conventional imaging includes plain film, CT, MRI, or US.
 - ◆ Often, further advanced imaging is needed when initial imaging, such as ultrasound, CT, or MRI does not answer the clinical question. Uncertain, indeterminate, inconclusive, or equivocal may describe these situations.
- Requests for many Healthcare Common Procedure Coding System (HCPCS) codes, including nonspecific codes such as S8042 [Magnetic resonance imaging (MRI), low-field], should be redirected to a more appropriate and specific CPT® code. Exceptions are noted in the applicable guidelines.
- Inappropriate use of contrast is a very important component of evidence-based advanced imaging use.
 - ◆ The appropriate levels of contrast for an examination (i.e. without contrast, with contrast, without and with contrast) is determined by the evidence- based guidance reflected in the condition-specific guideline sections.
 - ◆ If, during the performance of a non-contrast imaging study, there is the unexpected need to use contrast in order to evaluate a possible abnormality, then that is appropriate.¹

Ultrasound

- Diagnostic ultrasound uses high frequency sound waves to evaluate soft tissue structures and vascular structures utilizing greyscale and Doppler techniques.
- Ultrasound allows for dynamic real-time imaging at the bedside
 - ◆ Ultrasound is limited in areas where there is dense bone or other calcification.
 - ◆ Ultrasound also has a relatively limited imaging window so may be of limited value to evaluate very large abnormalities
 - ◆ In general, ultrasound is highly operator-dependent, and proper training and experience are required to perform consistent, high quality evaluations.
- Indications for ultrasound may include, but are not limited to:
 - ◆ Obstetric and gynecologic imaging
 - ◆ Soft tissue and visceral imaging of the chest, abdomen, pelvis, and extremities
 - ◆ Brain and spine imaging when not obscured by dense bony structures
 - ◆ Vascular imaging when not obscured by dense bony structures
 - ◆ Procedural guidance when not obscured by dense bony structures
 - ◆ Initial evaluation of ill-defined soft tissue masses or fullness and differentiating adenopathy from mass or cyst. Prior to advanced imaging, ultrasound can be

very beneficial in selecting the proper modality, body area, image sequences, and contrast level that will provide the most definitive information for the patient.

- More specific guidance for ultrasound usage, including exceptions to this general guidance, can be found throughout the condition specific guidelines.

Computed Tomography (CT):

- The AMA CPT manual does not describe nor assign any minimum or maximum number of sequences for any CT study. CT imaging protocols are often influenced by the individual clinical situation of the patient and additional sequences are not uncommon. There are numerous CT protocols that may be performed to evaluate specific clinical questions, and this technology is constantly undergoing development.
- CT utilizes ionizing radiation to create cross-sectional and volumetric images of the body.
 - ◆ Advantages over ultrasound include a much larger field of view, and faster completion time in general. Disadvantages compared to ultrasound include lack of portability and exposure to ionizing radiation.
 - ◆ Advantages over MRI include faster imaging, and a more spacious scanner area limiting claustrophobia. Disadvantages compared to MRI include decreased soft tissue definition, especially with non-contrast imaging, and exposure to ionizing radiation.
- CT can be performed without, with, or without and with intravenous (IV) contrast depending on the clinical indication and body area.
 - ◆ In general, non-contrast imaging is appropriate for evaluating structures with significant tissue density differences such as lung parenchyma and bony structures, or when there is a contraindication to contrast.
 - ◆ In general, CT with contrast is the most common level of contrast, and can be used when there is need for improved vascular or soft tissue resolution, including better characterization of known or suspected malignancy, as well as infectious and inflammatory conditions.
 - ◆ CT without and with contrast has a limited role as the risks of doubling the ionizing radiation exposure rarely outweigh the benefits of multiphasic imaging, though there are some exceptions which include but are not limited to:
 - Characterization of a mass
 - Characterization of arterial and venous anatomy
 - ◆ More specific guidance for CT contrast usage, including exceptions to this general guidance can be found throughout the condition-specific guidelines.
- Shellfish allergy:
 - ◆ It is commonly assumed that an allergy to shellfish infers iodine allergy, and that this implies an allergy to CT iodinated contrast media. However, this is NOT true. Shellfish allergy is due to tropomyosins. Iodine plays no role in these allergic reactions. Allergies to shellfish do not increase the risk of reaction to IV CT contrast any more than that of other allergens.

- Enteric contrast (oral or rectal) is sometimes used in abdominal imaging. There is no specific CPT code which refers to enteric contrast.
- The appropriate contrast level and anatomic region in CT imaging is specific to the clinical indication, as listed in the specific guideline sections.
- CT should not be used to replace MRI in an attempt to avoid sedation unless it is listed as a recommended study the appropriate guideline.
- There are significant potential adverse effects associated with the use of iodinated contrast media. These include hypersensitivity reactions, thyroid dysfunction, and contrast-induced nephropathy (CIN). Patients with impaired renal function are at increased risk for CIN.²
- Both contrast CT and MRI may be considered to have the same risk profile with renal failure (GFR <30 mL/min).
- The use of CT contrast should proceed with caution in pregnant and breastfeeding patients. There is a theoretical risk of contrast toxicity to the fetal and infant thyroid. The procedure can be performed if the specific need for that contrast-enhanced procedure outweighs risk to the fetus. Breastfeeding patients may reduce this risk by choosing to pump and discard breast milk for 12-24 hours after the contrast injection.
- CT without contrast may be appropriate if clinical criteria for CT with contrast are met AND the patient has:
 - ◆ Elevated blood urea nitrogen (BUN) and/or creatinine
 - ◆ Renal insufficiency
 - ◆ Allergies to iodinated CT contrast
 - ◆ Thyroid disease which could be treated with I-131
 - ◆ Diabetes
 - ◆ Very elderly
 - ◆ Urgent or emergent settings due to availability
 - ◆ Trauma
- CT is superior to other imaging modalities in certain conditions, including but not limited to the following:
 - ◆ Screening following trauma
 - ◆ Imaging pulmonary disease
 - ◆ Imaging abdominal and pelvic viscera
 - ◆ Imaging of complex fractures
 - ◆ Evaluation of inconclusive findings on Ultrasound or MRI, or if there is a contraindication to MRI
- More specific guidance for CT usage, including exceptions to this general guidance can be found throughout the condition specific guidelines.

Magnetic Resonance Imaging (MRI):

- The AMA CPT manual does not describe nor assign any minimum or maximum number of sequences for any MRI study. MRI protocols are often influenced by the individual clinical situation of the patient and additional sequences are not uncommon. There are numerous MRI sequences that may be performed to evaluate specific clinical questions, and this technology is constantly undergoing development.
- Magnetic Resonance Imaging (MRI) utilizes the interaction between the intrinsic radiofrequency of certain molecules in the body (hydrogen in most cases) and a strong external magnetic field.
 - ◆ MRI is often superior for advanced imaging of soft tissues, and can also define physiological processes in some instances [e.g. edema, loss of circulation (AVN), and increased vascularity (tumors)].
 - ◆ MRI does not use ionizing radiation, and even noncontrast images have much higher soft tissue definition than CT or Ultrasound
 - ◆ MRI typically takes much longer than either CT or Ultrasound, and for some patients may require sedation. It is also much more sensitive to patient motion that can degrade image quality than either CT or Ultrasound.
- MRI Breast and MRI Chest are not interchangeable, as they focus detailed sequences on different adjacent body parts.
- MRI may be utilized either as the primary advanced imaging modality, or when further definition is needed based on CT or ultrasound imaging.
- Most orthopedic and dental implants are not magnetic. These include hip and knee replacements; plates, screws, and rods used to treat fractures; and cavity fillings. Yet, all of these metal implants can distort the MRI image if near the part of the body being scanned.
 - ◆ Other implants, however, may have contraindications to MRI. These include:
 - Pacemakers
 - ICD or heart valves
 - Metal implants in the brain
 - Metal implants in the eyes or ears
 - Infusion catheters and bullets or shrapnel.
 - ◆ CT can therefore be an alternative study to MRI in these scenarios.
- The contrast level and anatomic region in MRI imaging is specific to the clinical indication, as listed in the specific guideline sections.
- MRI is commonly performed without, without and with contrast.
 - ◆ Non-contrast imaging offers excellent tissue definition
 - ◆ Imaging without and with contrast is commonly used when needed to better characterize tissue perfusion and vascularization.
 - Most contrast is gadolinium based and causes T2 brightening of the vascular and extracellular spaces.
 - Some specialized gadolinium and non-gadolinium contrast agents are available, and most commonly used for characterizing liver lesions.

- ◆ MRI with contrast only is rarely appropriate, and is usually used to better characterize findings on a recent inconclusive non-contrast MRI, commonly called a completion study.
- ◆ MRI contrast is contraindicated in pregnant individuals
- ◆ More specific guidance for MRI contrast usage, including exceptions to this general guidance can be found throughout the condition specific guidelines.
- MRI may be preferred in patients with renal failure, and in patients allergic to intravenous CT contrast.
 - ◆ Both contrast CT and MRI may be considered to have the same risk profile with renal failure (GFR <30 mL/min).
 - ◆ Gadolinium can cause Nephrogenic Systemic Fibrosis (NSF). The greater the exposure to gadolinium in individuals with a low GFR (especially if on dialysis), the greater the chance of individuals developing NSF.
 - ◆ Multiple studies have demonstrated potential for gadolinium deposition following the use of gadolinium-based contrast agents (GBCAs) for MRI studies.^{3,4,5,6,7} The U.S. Food and Drug Administration (FDA) has noted that there is currently no evidence to suggest that gadolinium retention in the brain is harmful and restricting gadolinium-based contrast agents (GBCAs) use is not warranted at this time. It has been recommended that GBCA use should be limited to circumstances in which additional information provided by the contrast agent is necessary and the necessity of repetitive MRIs with GBCAs should be assessed.⁸
- A CT may be approved in place of an MRI when clinical criteria are met for MRI AND there is a contraindication to having an MRI (pacemaker, ICD, insulin pump, neurostimulator, etc.)
 - ◆ When replacing MRI with CT, contrast level matching should occur as follows:
 - MRI without contrast → CT without contrast
 - MRI without and with contrast → CT with contrast or CT without and with contrast
- The following situations may impact the appropriateness for MRI and or MR contrast
 - ◆ Caution should be taken in the use of gadolinium in patients with renal failure
 - ◆ The use of gadolinium contrast agents is contraindicated during pregnancy unless the specific need for that procedure outweighs risk to the fetus.
 - ◆ MRI can be performed for non-ferromagnetic body metals (i.e. titanium), although some imaging facilities will consider it contraindicated if recent surgery, regardless of the metal type
- MRI should not be used as a replacement for CT for the sole reason of avoidance of ionizing radiation when CT is not supported in the condition based guidelines, since it does not solve the problem of overutilization.
- MRI is superior to other imaging modalities in certain conditions, including but not limited to the following:
 - ◆ Imaging the brain and spinal cord
 - ◆ Characterizing visceral and musculoskeletal soft tissue masses
 - ◆ Evaluating musculoskeletal soft tissues including ligaments and tendons

- ◆ Evaluating inconclusive findings on ultrasound or CT
- ◆ Patients who are pregnant or have high radiation sensitivity
- ◆ Suspicion, diagnosis of or surveillance of infections
- More specific guidance for MRI usage, including exceptions to this general guidance can be found throughout the condition specific guidelines.

Positron Emission Tomography (PET):

- PET is a nuclear medicine study that uses a positron emitting radiotracer to create cross-sectional and volumetric images based on tissue metabolism.
- Conventional imaging (frequently CT, sometimes MRI or bone scan, see Preface 3.2 for more detail) of the affected area(s) drives much of initial and restaging and surveillance imaging for malignancy and other chronic conditions. PET is not indicated for surveillance imaging unless specifically stated in the condition-specific guideline sections.
- PET/MRI is generally not supported, see **Preface-5.3:PET-MRI**
- PET is rarely performed as a single modality, but is typically performed as a combined PET/CT.
 - ◆ The unbundling of PET/CT into separate PET and diagnostic CT CPT codes is not supported, because PET/CT is done as a single study.
- PET/CT lacks the tissue definition of CT or MRI, but is fairly specific for metabolic activity based on the radiotracer used
 - ◆ Fluorodeoxyglucose (fluorine-18-2-fluoro-2-deoxy-D-glucose [FDG]) is the most common PET radiotracer and images glucose metabolism
 - ◆ Some specialized radiotracers including Gallium-68 DOTATATE, C-11 Choline, and F-18 Fluciclovine are supported in evaluation for some oncologic conditions, while the use of other radiotracers including but not limited to F-18 Sodium Fluoride and Gallium-68 PSMA are not supported.
- Indications for PET/CT may include
 - ◆ Oncologic Imaging for evaluation of tumor metabolic activity
 - ◆ Cardiac Imaging for evaluation of myocardial metabolic activity
 - ◆ Brain Imaging for evaluation of metabolic activity for procedural planning
- More specific guidance for PET usage, including exceptions to this general guidance can be found throughout the condition specific guidelines.

Overutilization of Advanced Imaging:

- A number of recent reports describe overutilization in all areas of advanced imaging, which may include:
 - ◆ High level testing without consideration of less invasive, lower cost options which may adequately address the clinical question at hand
 - ◆ Excessive radiation and costs with unnecessary testing
 - ◆ Defensive medical practice
 - ◆ CT without and with contrast (so called “double contrast studies) requests, which have few current indications.

- ◆ MRI requested in place of CT to avoid radiation without considering the primary indication for imaging
 - ◆ Adult CT settings and protocols used for smaller people and children
 - ◆ Unnecessary imaging procedures when the same or similar studies have already been conducted.
- A review of the imaging histories of all patients presenting for studies has been recognized as one of the more important processes that can be significantly improved. By recognizing that a duplicate or questionably indicated examination has been ordered for patients, it may be possible to avoid exposing them to unnecessary risks.^{9, 10} To avoid these unnecessary risks, the precautions below should be considered.
- ◆ The results of initial diagnostic tests or radiologic studies to narrow the differential diagnosis should be obtained prior to performing further tests or radiologic studies.
 - ◆ The clinical history should include a potential indication such as a known or suspected abnormality involving the body part for which the imaging study is being requested. These potential indications are addressed in greater detail within the applicable guidelines.
 - ◆ The results of the requested imaging procedures should be expected to have an impact on patient management or treatment decisions.
 - ◆ Repeat imaging studies are not generally necessary unless there is evidence of disease progression, recurrence of disease, and/or the repeat imaging will affect a patient's clinical management.

References

1. Bettmann MA. Frequently Asked Questions: Iodinated Contrast Agents. *Radiographics* 2004; 24:S3–S10
2. Andreucci M, Solomon R, Tasanarong A. Side Effects of Radiographic Contrast Media: Pathogenesis, Risk Factors, and Prevention. *BioMed Research International*. 2014;2014:1-20. <https://doi.org/10.1155/2014/741018>.
3. McDonald RJ, McDonald JS, Kallmes DF, et al. Intracranial Gadolinium Deposition after Contrast-enhanced MR Imaging. *Radiology*. 2015;275(3):772-782.
4. Kanda T, et al. High signal intensity in the dentate nucleus and globus pallidus on unenhanced T1-weighted MR images: relationship with increasing cumulative dose of a gadolinium-based contrast material. *Radiology* 2014;270:834–841. pmid:24475844
5. Olchowy et al. The presence of the gadolinium-based contrast agent depositions in the brain and symptoms of gadolinium neurotoxicity - A systematic review. *PLOS ONE*. February 10, 2017. <https://doi.org/10.1371/journal.pone.0171704>
6. Ramalho J, et al. High Signal Intensity in Globus Pallidus and Dentate Nucleus on Unenhanced T1-weighted MR Images: Evaluation of Two Linear Gadolinium-based Contrast Agents. *Radiology* 2015;276:836–844. pmid:26079490
7. Radbruch A, et al. Intraindividual Analysis of Signal Intensity Changes in the Dentate Nucleus After Consecutive Serial Applications of Linear and Macrocyclic Gadolinium-Based Contrast Agents. *Investigative Radiology* 2016;51:683–690. pmid:27495187
8. Safety Alerts for Human Medical Products > Gadolinium-based Contrast Agents (GBCAs): Drug Safety Communication - Retained in Body; New Class Warnings [12-19-2017]. Accessed 2/27/2018. <https://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm589580.htm>
9. ES Amis, et al. American College of Radiology White Paper on Radiation Dose in Medicine. *J Am Coll Radiol* 2007;4:272-284.

10. Powell AC, Long JW, Kren EM, Gupta AK, Levin DC. Evaluation of a Program for Improving Advanced Imaging Interpretation. *J Patient Saf.* 2019 Mar;15(1):69-75
11. FDA. White Paper: Initiative to Reduce Unnecessary Radiation Exposure from Medical Imaging. Page Last Updated: 02/05/2018. <https://www.fda.gov/Radiation-EmittingProducts/RadiationSafety/RadiationDoseReduction/ucm199994.htm>
12. Update on FDA approach to safety issue of gadolinium retention after administration of gadolinium-based contrast agents. <https://www.fda.gov/media/116492/download>.
13. Blumfield E, Swenson DW, Iyer RS, Stanescu AL. Gadolinium-based contrast agents — review of recent literature on magnetic resonance imaging signal intensity changes and tissue deposits, with emphasis on pediatric patients. *Pediatric Radiology.* 2019;49(4):448-457. doi:10.1007/s00247-018-4304-8.
14. ACR – SPR –SRU PRACTICE PARAMETER FOR THE PERFORMING AND INTERPRETING DIAGNOSTIC ULTRASOUND EXAMINATIONS Revised 2017 (Resolution 32) <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Perf-Interpret.pdf>
15. ACR–SPR PRACTICE PARAMETER FOR PERFORMING FDG-PET/CT IN ONCOLOGY Revised 2016 (Resolution 25) <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/FDG-PET-CT.pdf>
16. ACR PRACTICE PARAMETER FOR PERFORMING AND INTERPRETING MAGNETIC RESONANCE IMAGING (MRI) Revised 2017 (Resolution 10) <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Perf-Interpret.pdf>
17. ACR PRACTICE PARAMETER FOR PERFORMING AND INTERPRETING DIAGNOSTIC COMPUTED TOMOGRAPHY (CT) Revised 2017 (Resolution 22) <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CT-Perf-Interpret.pdf>
18. Lohrke, J., Frenzel, T., Endrikat, J., Alves, F. C., Grist, T. M., Law, M., Lee, J. M., Leiner, T., Li, K. C., Nikolaou, K., Prince, M. R., Schild, H. H., Weinreb, J. C., Yoshikawa, K., & Pietsch, H. (2016). 25 Years of Contrast-Enhanced MRI: Developments, Current Challenges and Future Perspectives. *Advances in therapy*, 33(1), 1–28. <https://doi.org/10.1007/s12325-015-0275-4>

Preface-4: Coding Issues

Preface-4.1: 3D Rendering	15
Preface-4.2: CT-, MR-, or Ultrasound-Guided Procedures	16
Preface-4.3: Unlisted Procedures/Therapy Treatment Planning	19
Preface-4.4: Unilateral versus Bilateral Breast MRI	19
Preface-4.5: CPT® 76380 Limited or Follow-up CT	19
Preface-4.6: SPECT/CT Imaging	20

Preface-4.1: 3D Rendering

CPT® 76376 and CPT® 76377:

- Both codes require concurrent supervision of the image post-processing 3D manipulation of the volumetric data set and image rendering.
 - ◆ Concurrent supervision is defined as active physician participation in and monitoring of the reconstruction process including design of the anatomic region that is to be reconstructed; determination of the tissue types and actual structures to be displayed (eg, bone, organs, and vessels); determination of the images or cine loops that are to be archived; and monitoring and adjustment of the 3D work product. The American College of Radiology (ACR) recommends that it is best to document the physician's supervision or participation in the 3D reconstruction of images.
- These two codes differ in the need for and use of an independent workstation for post-processing.
 - ◆ CPT® 76376 reports procedures not requiring image post-processing on an independent workstation.
 - ◆ CPT® 76377 reports procedures that require image post-processing on an independent workstation.
- These 3D rendering codes should not be used for 2D reformatting.
- Two-dimensional reconstruction (e.g. reformatting an axial scan into the coronal plane) is now included in all cross-sectional imaging base codes and is not separately reimbursable.
- Some payers do not reimburse separately for CPT® 76376 or CPT® 76377. In addition, these CPT® codes are not included in every eviCore patient's radiology management program.
 - ◆ The codes used to report 3D rendering for ultrasound and echocardiography are also used to report the 3D post processing work on CT, MRI, and other tomographic modalities.
- Providers may be required to obtain prior authorization on these 3D codes even if prior authorization is not required for the echocardiography and/or ultrasound procedure codes. It may appear that eviCore pre-authorizes echocardiography and/or ultrasound when, in fact, it may only be the 3D code that needs the prior authorization.
 - ◆ Prior authorization requirements are established on a CPT® code level and vary by the individual health plan payor.
 - ◆ Providers are urged to obtain written instructions and requirements directly from each payor.
- CPT® codes for 3D rendering should not be billed in conjunction with computer-aided detection (CAD), MRA, CTA, nuclear medicine SPECT studies, PET, PET/CT, Mammogram, MRI Breast, US Breast, CT Colonography (virtual colonoscopy), Cardiac MRI, Cardiac CT, or Coronary CTA studies.

- CPT® 76377 (3D rendering requiring image post-processing on an independent workstation) or CPT® 76376 (3D rendering not requiring image post-processing on an independent workstation) can be considered in the following clinical scenarios:
 - ◆ Bony conditions:
 - Evaluation of congenital skull abnormalities in newborns, infants, and toddlers (usually for preoperative planning)
 - Complex joint fractures or pelvis fractures
 - Spine fractures (usually for preoperative planning)
 - Complex facial fractures
 - ◆ Preoperative planning for other complex surgical cases
 - ◆ Cerebral angiography
 - ◆ Pelvis conditions:
 - Uterine intra-cavitary lesion when initial US is equivocal (See **PV-2.1: Abnormal Uterine Bleeding (AUB)** and **PV-12.1: Leiomyomata** in the Pelvis Imaging Guidelines)
 - Hydrosalpinx or peritoneal cysts when initial US is indeterminate (See **PV-5.3: Complex Adnexal Masses** in the Pelvis Imaging Guidelines)
 - Lost IUD (inability to feel or see IUD string) with initial US (See **PV-10.1: Intrauterine Device** in the Pelvis Imaging Guidelines)
 - Uterine anomalies with initial US (See **PV-14.1: Uterine Anomalies** in the Pelvis Imaging Guidelines)
 - Infertility (See **PV-9.1: Infertility Evaluation, Female** in the Pelvis Imaging Guidelines)
 - ◆ Abdomen conditions:
 - CT Urogram (See **AB-39: Hematuria and Hydronephrosis** in the Abdomen Imaging Guidelines)
 - MRCP (See **AB-27: MR Cholangiopancreatography (MRCP)** in the Abdomen Imaging Guidelines)

Preface-4.2: CT-, MR-, or Ultrasound-Guided Procedures

- CT, MR, and Ultrasound guidance procedure codes contain all the imaging necessary to guide a needle or catheter. It is inappropriate to routinely bill a diagnostic procedure code in conjunction with a guidance procedure code.
- Imaging studies performed as part of a CT-, MR-, or Ultrasound-guided procedure should be reported using the CPT® codes in the following table.

TABLE: Imaging Guidance Procedure Codes

CPT®	Description
19085	Biopsy, breast, with placement of breast localization device(s), when performed, and imaging of the biopsy specimen, when performed, percutaneous; first lesion, including MR guidance
19086	Biopsy, breast, with placement of breast localization device(s), when performed, and imaging of the biopsy specimen, when performed, percutaneous; first lesion, including MR guidance; each additional lesion, including MR guidance
75989	Imaging guidance for percutaneous drainage with placement of catheter (all modalities)
77011	CT guidance for stereotactic localization
77012	CT guidance for needle placement
77013	CT guidance for, and monitoring of parenchymal tissue ablation
77021	MR guidance for needle placement
77022	MR guidance for, and monitoring of parenchymal tissue ablation
76942	Ultrasonic guidance for needle placement

CPT® 19085 and CPT® 19086:

- The proper way to bill an MRI guided breast biopsy is CPT® 19085 (Biopsy, breast, with placement of breast localization device(s), when performed, and imaging of the biopsy specimen, when performed, percutaneous; first lesion, including MR guidance). Additional lesions should be billed using CPT® 19086.

CPT® 75989:

- This code is used to report imaging guidance for a percutaneous drainage procedure in which a catheter is left in place.
- This code can be used to report whether the drainage catheter is placed under fluoroscopy, ultrasound, CT, or MR guidance modality.

CPT® 77011:

- A stereotactic CT localization scan is frequently obtained prior to sinus surgery. The dataset is then loaded into the navigational workstation in the operating room for use during the surgical procedure. The information provides exact positioning of surgical instruments with regard to the patient's 3D CT images.
- In most cases, the preoperative CT is a technical-only service that does not require interpretation by a radiologist.
 - ◆ The imaging facility should report CPT® 77011 when performing a scan not requiring interpretation by a radiologist.
 - ◆ If a diagnostic scan is performed and interpreted by a radiologist, the appropriate diagnostic CT code (e.g., CPT® 70486) should be used.

- ◆ It is not appropriate to report both CPT® 70486 and CPT® 77011 for the same CT stereotactic localization imaging session.
- ◆ 3D Rendering (CPT® 76376 or CPT® 76377) should not be reported in conjunction with CPT® 77011 (or CPT® 70486 if used). The procedure inherently generates a 3D dataset.

CPT® 77012 (CT) and CPT® 77021 (MR):

- These codes are used to report imaging guidance for needle placement during biopsy, aspiration, and other percutaneous procedures.
- They represent the radiological supervision and interpretation of the procedure and are often billed in conjunction with surgical procedure codes.
 - ◆ For example, CPT® 77012 is reported when CT guidance is used to place the needle for a conventional arthrogram.
 - ◆ Only codes representing percutaneous surgical procedures should be billed with CPT® 77012 and CPT® 77021. It is inappropriate to use with surgical codes for open, excisional, or incisional procedures.

CPT® 77013 (CT) and CPT® 77022 (MR):

- These codes include the initial guidance to direct a needle electrode to the tumor(s), monitoring for needle electrode repositioning within the lesion, and as necessary for multiple ablations to coagulate the lesion and confirmation of satisfactory coagulative necrosis of the lesion(s) and comparison to pre-ablation images.
 - ◆ **NOTE:** CPT® 77013 should only be used for non-bone ablation procedures.
 - ◆ CPT® 20982 includes CT guidance for bone tumor ablations.
 - ◆ Only codes representing percutaneous surgical procedures should be billed with CPT® 77013 and CPT® 77022. It is inappropriate to use with surgical codes for open, excisional, or incisional procedures.
- CPT® 77012 and CPT® 77021 (as well as guidance codes CPT® 76942 [US], and CPT® 77002 - CPT® 77003 [fluoroscopy]) describe radiologic guidance by different modalities.
 - ◆ Only one unit of any of these codes should be reported per patient encounter (date of service). The unit of service is considered to be the patient encounter, not the number of lesions, aspirations, biopsies, injections, or localizations.

Preface-4.3: Unlisted Procedures/Therapy Treatment Planning

CPT®	Description
76497	Unlisted CT procedure (e.g., diagnostic or interventional)
76498	Unlisted MR procedure (e.g., diagnostic or interventional)
78999	Unlisted procedure, diagnostic nuclear medicine

- In the absence of written payor instructions, these unlisted codes should be reported whenever a diagnostic or interventional CT or MR study is performed in which an appropriate anatomic site-specific code is not available.
 - ◆ A Category III code that describes the procedure performed must be reported rather than an unlisted code if one is available.
- CPT® 76497 or CPT® 76498 (Unlisted CT or MRI procedure) can be considered in the following clinical scenarios:
 - ◆ Studies done for navigation and planning for neurosurgical procedures (i.e. Stealth or Brain Lab Imaging)^{1,2}
 - ◆ Custom joint Arthroplasty planning if covered by payor (not as Alternative Recommendation) (See **MS-12.1: Osteoarthritis** in the Musculoskeletal Imaging Guidelines)
 - ◆ Any procedure/surgical planning if thinner cuts or different positional acquisition (than those on the completed diagnostic study) are needed. These could include navigational bronchoscopy.

Therapy Treatment Planning

- Radiation Therapy Treatment Planning: See **ONC-1.5: Unlisted Procedure Codes in Oncology** in the Oncology Imaging Guidelines

Preface-4.4: Unilateral versus Bilateral Breast MRI

- Diagnostic MRI of both breasts should be coded as CPT® 77049 regardless of whether both breasts are imaged simultaneously or whether unilateral breast MRI is performed in two separate imaging sessions.

Preface-4.5: CPT® 76380 Limited or Follow-up CT

- CPT® 76380 describes a limited or follow-up CT scan. The code is used to report any CT scan, for any given area of the body, in which the work of a full diagnostic code is not performed.
- Common examples include (but are not limited to):
 - ◆ Limited sinus CT imaging protocol
 - ◆ Limited or follow-up slices through a known pulmonary nodule
 - ◆ Limited slices to assess a non-healing fracture (such as the clavicle)
- It is inappropriate to report CPT® 76380, in conjunction with other diagnostic CT codes, to cover 'extra slices' in certain imaging protocols.
 - ◆ There is no specific number of sequences or slices defined in any CT CPT® code definition.

- ◆ The AMA, in *CPT*[®] 2019, does not describe nor assign any minimum or maximum number of sequences or slices for any CT study.
 - A few additional slices or sequences are not uncommon.
 - CT imaging protocols are often influenced by the individual clinical situation of the patient. Sometimes the protocols require more time and sometimes less.

Preface-4.6: SPECT/CT Imaging

- SPECT/CT involves SPECT (Single Photon Emission Computed Tomography) nuclear medicine imaging and CT for optimizing location, accuracy, and attenuation correction and combines functional and anatomic information.
 - ◆ Common studies using this modality include ¹²³I- or ¹³¹I-Metaiodobenzylguanidine (MIBG) and octreotide scintigraphy for neuroendocrine tumors.
- Hybrid Nuclear/CT scan can be CPT[®] 78830 - single area and single day, CPT[®] 78831 - 2 or more days, or CPT[®] 78832 - 2 areas with one day and 2 day study.
- A procedure code for SPECT/CT parathyroid nuclear imaging, (CPT[®] 78072), became effective January 1, 2013.

References

1. Society of Nuclear Medicine and Molecular Imaging Coding Corner <http://www.snmmi.org/ClinicalPractice/CodingCornerPT.aspx?ItemNumber=1786>.
2. Intraoperative MR. Brainlab. <https://www.brainlab.com/surgery-products/overview-neurosurgery-products/intraoperative-mr/>.
3. Experience the Advanced 3D Sinus Surgery Planning with Scopis Building Blocks planning software. Scopis Planning. <http://planning.scopis.com/>.

Preface-5: Whole Body Imaging

Preface-5.1: Whole Body CT Imaging	22
Preface-5.2: Whole Body MR Imaging	22
Preface-5.3: PET-MRI	23

Preface-5.1: Whole Body CT Imaging

- Whole body CT or LifeScan (CT Brain, Chest, Abdomen, and Pelvis) for screening of asymptomatic patients is not a covered benefit of any of the current health plans who have delegated utilization review to eviCore. The performance of whole body screening CT examinations in healthy patients does not meet any of the current validity criteria for screening studies and there is no clear documentation of benefit versus radiation risk.
- Whole body low dose CT is supported for oncologic staging in Multiple Myeloma (See **ONC-25: Multiple Myeloma and Plasmacytomas** in the Oncology Imaging Guidelines)

Preface-5.2: Whole Body MR Imaging

- Whole body MRI (WBMRI) is, with the exception of select cancer predisposition syndromes discussed below, generally not supported by eviCore at this time due to lack of standardization in imaging technique and lack of evidence that WBMRI improves patient outcome for any individual disease state.
 - ◆ While WBMRI has the benefit of whole body imaging and lack of radiation exposure, substantial variation still exists in the number of images, type of sequences (STIR vs. diffusion weighting, for example), and contrast agent(s) used.
- Coding considerations:
 - ◆ There are no established CPT® or HCPCS codes for reporting WBMRI.
 - ◆ WBMRI is at present only reportable using CPT® 76498. All other methods of reporting whole body MRI are inappropriate, including:
 - Separate diagnostic MRI codes for multiple individual body parts
 - MRI Bone Marrow Supply (CPT® 77084)
- Disease-specific considerations:
 - ◆ Cancer screening:
 - Interval WBMRI is recommended for cancer screening in patients with select cancer predisposition syndromes. Otherwise, WBMRI has not been shown to improve outcomes for cancer screening. See **PEDONC-2.2: Li-Fraumeni Syndrome (LFS)**, **PEDONC-2.13: Hereditary Paraganglioma-Pheochromocytoma (HPP) Syndromes**, **PEDONC-2.15: Constitutional Mismatch Repair Deficiency (CMMRD or Turcot Syndrome)** in the Pediatric Oncology Imaging Guidelines for additional information
 - ◆ Cancer staging and restaging
 - While the feasibility of WBMRI has been established, data remain conflicting on whether WBMRI is of equivalent diagnostic accuracy compared with standard imaging modalities such as CT, scintigraphy, and PET imaging. Evidence has not been published establishing WBMRI as a standard evaluation for any type of cancer.

- ◆ Autoimmune disease
 - WBMRI has been shown to increase the number of detected lesions in chronic multifocal osteomyelitis and other inflammatory arthritides, but no improvement in outcomes from the use of WBMRI has yet been shown. See **PEDMS-10.2: Chronic Recurrent Multifocal Osteomyelitis** in the Pediatric Musculoskeletal Imaging Guidelines for additional information.

Preface-5.3: PET-MRI

- PET-MRI is, generally, not supported by eviCore at this time due to lack of standardization in imaging technique and lack of evidence that PET-MRI improves patient outcome for any individual disease state.

References

1. Villani A, Tabori U, Schiffman J, et al, Biochemical and imaging surveillance in germline TP53 mutation carriers with Li-Fraumeni syndrome: a prospective observational study, *Lancet Oncol* 2011;12:559-567.
2. Siegel MJ, Acharyya S, Hoffer FA et al, Whole-Body MR Imaging for Staging of Malignant Tumors in Pediatric Patients: Results of the American College of Radiology Imaging Network 6660 Trial, *Radiology* 2013;266:599-609.
3. Antoch G, Vogt FM, Freudenberg LS, et al, Whole-Body Dual-Modality PET/CT and Whole-Body MRI for Tumor Staging in Oncology, *JAMA* 2003;290:3199-3206.
4. Lauenstein TC and Semelka RC, Emerging Techniques: Whole-Body Screening and Staging With MRI, *J Magn Reson Imaging* 2006;24:489-498.
5. Khanna G, Sato TP, and Ferguson P, Imaging of Chronic Recurrent Multifocal Osteomyelitis, *Radiographics* 2009;29:1159-1177.
6. Ferguson PJ and Sandu M, Current Understanding of the Pathogenesis and Management of Chronic Recurrent Multifocal Osteomyelitis, *Curr Rheumatol Rep* 2012;14:130-141.
7. National Comprehensive Cancer Network (NCCN) Guidelines Version 3.2019. – January 18, 2019, Genetic/Familial High Risk Assessment: Breast and Ovarian, available at: https://www.nccn.org/professionals/physician_gls/pdf/genetics_screening.pdf. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines™) for Genetic/Familial High Risk Assessment: Breast and Ovarian V3.2019. – January 18, 2019 ©. 2019 National Comprehensive Cancer Network, Inc. All rights reserved. The NCCN Guidelines™ and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most recent and complete version of the NCCN Guidelines™, go online to NCCN.org.

Preface-6: References

- Complete reference citations for the journal articles are embedded within the body of the guidelines and/or may be found on the Reference pages at the end of some guideline sections.
- The website addresses for certain references are included in the body of the guidelines but are not hyperlinked to the actual website.
- The website address for the American College of Radiology (ACR) Appropriateness Criteria® is <http://www.acr.org>.

Preface-7: Copyright Information

- ©2021 eviCore healthcare. All rights reserved. No part of these materials may be changed, reproduced, or transmitted in any form or by any means, electronic or mechanical, including photocopying or recording, or in any information storage or retrieval system, without the prior express written permission of eviCore.

Preface-8: Trademarks

- **CPT® (Current Procedural Terminology)** is a registered trademark of the American Medical Association (AMA). **CPT®** five digit codes, nomenclature, and other data are copyright 2019 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values, or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.