



CLINICAL GUIDELINES

CMM-211: Spinal Cord Stimulators

Version 20.0.2018

Effective October 22, 2018



CMM-211: Spinal Cord Stimulators	
CMM-211.1: Definitions	3
CMM-211.2: Indications	4
CMM-211.3: Replacement	5
CMM-211.4: Non-Indications	6
CMM-211.5: Procedure (CPT®) Codes	8
CMM-211.6: References	9

CMM-211.1: Definitions

Spinal cord stimulation, also known as dorsal column stimulation or neuromodulation, is a reversible therapy applied for neuropathic pain with techniques that include multi-output implanted pulse generators and a choice of electrodes, some of which can be placed percutaneously. The technical goal of this therapy is to achieve stimulation of paresthesia of the dorsal horn of the spinal cord at a subjectively comfortable level, overlapping an individual's topography of pain. The procedure initially involves a short-term trial (e. g., greater than 48 hours) of percutaneous (temporary) spinal cord stimulation, prior to the subcutaneous (permanent) implantation of the spinal cord stimulation device, to determine whether the spinal cord stimulator device will induce sufficient pain relief to render it medically necessary.

High frequency spinal cord stimulation, also referred to as kilohertz frequency spinal cord stimulation or HF10, is a type of spinal cord stimulation (SCS) providing a higher frequency than traditional spinal cord stimulator systems. The HF10 SCS uses low amplitude, high frequency, and short duration pulses. HF10 SCS does not generate paresthesia and operates at a frequency of 10,000 Hz to provide pain relief in comparison to traditional spinal cord stimulation systems, which operate at a frequency in the range of 40-60 Hz and do generate paresthesia. As an alternative to traditional dorsal spinal column stimulation, HF10 SCS is proven safe and effective for treatment of chronic, intractable low back and leg pain in patients with failed back surgery syndrome.

In contrast to spinal cord stimulation, **peripheral nerve stimulation** or dorsal root ganglion stimulation, involves implantation of electrodes near or on a peripheral nerve to reduce pain. **Peripheral nerve field stimulation** is a technology that involves placement of electrodes subcutaneously within an area of maximal pain, with the objective of stimulating a region of affected nerves to reduce pain. Depending on the targeted nerve, leads may be placed percutaneously just under the skin or via an open approach for larger deeper peripheral nerves. Similar to spinal cord stimulation, a short term trial is required prior to permanent implantation of a generator. The use of these technologies, used alone or in combination with spinal cord stimulation for treatment of pain conditions is under investigation.

CMM-211.2: Indications

The determination of medical necessity for implantation of a dorsal spinal cord stimulator is always made on a case-by-case basis.

Chronic Intractable Pain Secondary to Failed Back Surgery Syndrome

- A short-term trial (e.g., greater than 48 hours) spinal cord stimulation [i.e., low frequency, high-frequency (HF 10 SCS)] is considered **medically necessary** for the treatment of chronic, intractable pain secondary to failed back surgery syndrome (FBSS) with intractable neuropathic leg pain when ALL of the following criteria are met:
 - ◆ Failure of at least six consecutive months of physician-supervised conservative medical management (e.g., pharmacotherapy, physical therapy, cognitive therapy, and activity lifestyle modification)
 - ◆ Surgical intervention is not indicated or for patients who do not wish to proceed with spinal surgery
 - ◆ An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would impact perception of pain and/or negatively impact the success of a SCS or contraindicate placement of the device.
- Permanent implantation of a spinal cord stimulator (i.e., low frequency, HF 10 SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to failed back surgery syndrome (FBSS) with intractable neuropathic leg pain when at least a 50% reduction in pain has been demonstrated during a short-term trial of SCS.

Complex Regional Pain Syndrome (CRPS)/Reflex Sympathetic Dystrophy (RSD)

- A short-term trial (i.e., greater than 48 hours) of a low frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) when ALL of the following criteria are met:
 - ◆ Failure of at least six consecutive months of physician-supervised conservative medical management (e.g., pharmacotherapy, physical therapy, cognitive therapy, and activity lifestyle modification)
 - ◆ Surgical intervention is not indicated
 - ◆ An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would impact perception of pain

and/or negatively impact the success of SCS or contraindicate placement of the device.

- Permanent implantation of a low frequency dorsal column spinal cord stimulator is considered **medically necessary** for the treatment of chronic, intractable pain secondary to complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) when at least a 50% reduction in pain has been demonstrated during a short-term trial of SCS.

Chronic Critical Limb Ischemia (CLI)

- A short-term trial (e.g., greater than 48 hours) of a low frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to chronic critical limb ischemia (CLI) when BOTH of the following criteria are met:
 - ◆ Failure of available conventional multidisciplinary medical (e.g., pharmacological, physical therapy) and surgical management (e.g., revascularization)
 - ◆ An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or drug dependence, depression, psychosis) that would impact perception of pain and/or negatively impact the success of a SCS or contraindicate placement of the device.
- Permanent implantation of a low frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to chronic critical limb ischemia (CLI) when a beneficial clinical response from a temporarily implanted electrode has been demonstrated prior to consideration of permanent implantation.

Chronic Stable Angina Pectoris

- A short-term trial (e.g., greater than 48 hours) of a low frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to chronic stable angina pectoris/myocardial ischemia when all of the following criteria are met:
 - ◆ Angina pectoris is Canadian Cardiovascular Society (CCS) functional class III or class IV (see Appendix A)
 - ◆ Attestation that the individual's treating cardiologist confirms coronary artery disease (CAD) and the individual is not a suitable candidate for a revascularization procedure
 - ◆ Optimal pharmacological treatment using anti-anginal medications (e.g., long-acting nitrates, beta-adrenergic blockers, or calcium-channel antagonists) has failed to adequately improve anginal symptoms
 - ◆ An evaluation by a mental health provider (e.g., a face-to-face assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental health problem (e.g., alcohol or

drug dependence, depression, psychosis) that would impact perception of pain and/or negatively impact the success of a SCS or contraindicate placement of the device.

- Permanent implantation of a low frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to chronic stable angina pectoris for myocardial ischemia when a beneficial clinical response from a temporarily implanted electrode has been demonstrated prior to consideration of permanent implantation.

CMM-211.3: Replacement

- The replacement of an existing high frequency or low frequency dorsal column spinal cord stimulator (SCS) and/or battery/generator is considered **medically necessary** for an individual when the existing stimulator and/or battery/generator is malfunctioning, cannot be repaired, and is no longer under warranty.
- Replacement of a functioning low frequency dorsal column spinal cord stimulator with a high frequency spinal cord stimulator is considered **not medically necessary**.

CMM-211.4: Non-Indications

- A high frequency spinal cord stimulator is considered **experimental, investigational, or unproven** for ANY other indication, including CRPS/RSD.
- A low frequency dorsal column spinal cord stimulator (SCS) is considered **experimental, investigational or unproven** for any other indication including but not limited to:
 - ◆ Post-amputation pain (phantom limb pain)
 - ◆ Post-herpetic neuralgia
 - ◆ Peripheral neuropathy
 - ◆ Dysesthesias involving the lower extremities secondary to spinal cord injury.
- Peripheral nerve stimulation, dorsal root ganglion stimulation, including peripheral nerve field stimulation, is considered **experimental, investigational, or unproven** for treatment of acute or chronic pain conditions, including ANY of the following:
 - ◆ Failed back surgery syndrome (FBSS) with intractable neuropathic leg pain
 - ◆ Complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD)
 - ◆ Chronic Critical Limb Ischemia (CLI)
 - ◆ Chronic Stable Angina Pectoris
 - ◆ Post-amputation pain (phantom limb pain)
 - ◆ Post-herpetic neuralgia
 - ◆ Peripheral neuropathy
 - ◆ Dysesthesias involving the lower extremities secondary to spinal cord injury.

Appendix A

New York Heart Association and Canadian Cardiovascular Society Functional Classifications

Class	New York Heart Association Functional Classification	Canadian Cardiovascular Society Functional Classification
I	Patients with cardiac disease but without resulting limitations of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnea, or anginal pain.	Ordinary physical activity does not cause angina, such as walking and climbing stairs. Angina occurs with strenuous or rapid or prolonged exertion at work or recreation.
II	Patients with cardiac disease resulting in slight limitation of physical activity. They are comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnea, or anginal pain.	Slight limitation of ordinary activity. Walking or climbing stairs rapidly, walking uphill, walking or stair climbing after meals, in cold, in wind, or under emotional stress, or only during the few hours after awakening. Walking more than two blocks on the level and climbing more than one flight of ordinary stairs at a normal pace and in normal conditions.
III	Patients with cardiac disease resulting in marked limitation of physical activity. They are comfortable at rest. Less than ordinary physical activity causes fatigue, palpitation, dyspnea, or anginal pain.	Marked limitation of ordinary physical activity. Walking one to two blocks on the level and climbing one flight in normal conditions and at a normal pace.
IV	Patient with cardiac disease resulting in inability to carry on any physical activity without discomfort. Symptoms of cardiac insufficiency or of the anginal syndrome may be present even at rest. If any physical activity is undertaken, discomfort is increased.	Inability to carry on any physical activity without discomfort—anginal syndrome may be present at rest.

(Heart Failure Society of America [HFSA], 2006; Gibbons, et al., 2002; American Heart Association [AHA], 1994; Canadian Cardiovascular Society [CCS], 1976).

CMM-211.5: Procedure (CPT®) Codes

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.	
CPT®	Code Description/Definition
63650	Percutaneous implantation of neurostimulator electrode array, epidural
63655	Laminectomy for implantation of neurostimulator electrodes, plate/paddle, epidural
63661	Removal of spinal neurostimulator electrode percutaneous array(s), including fluoroscopy, when performed
63662	Removal of spinal neurostimulator electrode plate/paddle(s) placed via laminotomy or laminectomy, including fluoroscopy, when performed
63663	Revision including replacement, when performed, of spinal neurostimulator electrode
63664	Revision including replacement, when performed, of spinal neurostimulator electrode
63685	Insertion or replacement of spinal neurostimulator pulse generator or receiver, direct or inductive coupling
63688	Revision or removal of implanted spinal neurostimulator pulse generator or receiver
95970	Electronic analysis of implanted neurostimulator pulse generator system (e.g., rate, pulse, amplitude and duration, configuration of wave form, battery status, electrode selectability, output modulation, cycling, impedance and patient compliance measurements); simple or complex brain, spinal cord, or peripheral (i.e., cranial nerve, peripheral nerve, autonomic nerve, neuromuscular) neurostimulator pulse generator/transmitter, without reprogramming
95971	Electronic analysis of implanted neurostimulator pulse generator system (e.g., rate, amplitude and duration, configuration of wave form, battery status, electrode selectability, output modulation, cycling, impedance and patient compliance measurement(s); simple spinal cord, or peripheral (i.e., peripheral nerve, automatic nerve, neuromuscular) neurostimulator pulse generation/transmitter, with intraoperative or subsequent programming
95972	Electrode analysis of implanted neurostimulator pulse generator system (e.g., rate, pulse amplitude and duration, configuration of wave form, battery status, electrode selectability, output modulation, cycling, impedance and patient compliance measurement(s); complex spinal cord, or peripheral (except cranial nerve) neurostimulator pulse generator/transmitter, with intraoperative or subsequent programming
This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.	

CMM-211.6: References

1. Amann W, Berg P, Gersbach P, et al; European Peripheral Vascular Disease Outcome Study SCS- EPOS. Spinal cord stimulation in the treatment of non-reconstructable stable critical leg ischaemia: results of the European Peripheral Vascular Disease Outcome Study (SCS-EPOS). *Eur J Vasc Endovasc Surg.* 2003;26(3):280-286.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline, 2nd Ed.* 2008.
3. American Medical Association. *Current Procedural Terminology – 2014 Professional Edition.*
4. Bicket MC, Dunn RY, Ahmed SU. High-frequency spinal cord stimulation for chronic pain: Pre-clinical overview and systematic review of controlled trials. *Pain Med* 2016; 17(12): 2326-36.
5. Burns B, Watkins L, Goadsby PJ. Treatment of intractable chronic cluster headache by occipital nerve stimulation in 14 patients. *Neurology* 2009; 72: 341–345.
6. Burns B, Watkins L, Goadsby PJ. Treatment of hemicrania continua by occipital nerve stimulation with a bion device: long-term follow-up of a crossover study. *Lancet Neurol* 2008;7:1001–12.
7. Claeys L, Berg W, Jonas S. Spinal cord stimulation in the treatment of chronic critical limb ischemia. *Acta Neurochir Suppl.* 2007;97(Pt 1):259-265.
8. Cruccu G, Aziz T, Garcia-Larrea L, et al. EFNS guidelines on neurostimulation therapy for neuropathic pain. *Eur J Neurol.* 2007;14(9):952-970.
9. Daousi C, Benbow SJ, MacFarlane IA. Electrical spinal cord stimulation in the long-term treatment of chronic painful diabetic neuropathy. *Diabet Med* 2005;22:393–398.
10. Deer T, Pope J, Hayek S, et al. Neurostimulation for the treatment of axial back pain: A review of mechanisms, techniques, outcomes, and future advances. *Neuromodulation* 2014; 17(suppl 2): 52-68.
11. de Vos CC, Meier K, Zaalberg PB, et al. Spinal cord stimulation in patients with painful diabetic neuropathy: a multicenter randomized clinical trial. *Pain* 2014;155:2426–2431.
12. Fontaine D, Christophe Sol J, Raoul S, Fabre N, Geraud G, Magne C et al. Treatment of refractory chronic cluster headache by chronic occipital nerve stimulation. *Cephalalgia* 2011;31:1101–5.
13. Forouzanfar T, Kemler M, Weber W, et al. Spinal cord stimulation in complex regional pain syndrome: cervical and lumbar devices are comparably effective. *Br J Anaesth.* 2004;92(3):348-53.
14. Gersbach P, Argitis V, Gardaz Jpet al. Late outcome of spinal cord stimulation for unreconstructable and limb-threatening lower limb ischemia. *Eur J Vasc Endovasc Surg.* 2007;33(6):717-724.
15. Grider JS, Manchikanti L, Carayannopoulos A, et al. Effectiveness of spinal cord stimulation in chronic spinal pain: A systemic review. *Pain Physician* 2016; 19(1):E33-54.
16. Harke H, Gretenkort P, Ladleif H, Rahman S. Spinal cord stimulation in sympathetically maintained complex regional pain syndrome type I with severe disability. A prospective clinical study. *Eur J Pain.* 2005;9(4):363-373.
17. Jose De Andres, MD, PhD, FIPP, EDRA, Vicente Monsalve-Dolz, PhD, Gustavo Fabregat-Cid, MD, PhD, Vicente Villanueva-Perez, MD, PhD, Anushik Harutyunyan, Juan Marcos Asensio-Samper, M.D, Nerea Sanchis-Lopex, MD. Prospective, Randomized Blind Effect-on-Outcome Study of Conventional vs High-Frequency Spinal Cord Stimulation in Patients with Pain and Disability Due to Failed Back Surgery Syndrome. *Pain Medicine* 2017; 18: 2401-2421.
18. Jivegård L, Augustinsson L, Holm J, et al. Effects of spinal cord stimulation (SCS) in patients with inoperable severe lower limb ischaemia: a prospective randomised controlled study. *Eur J Vasc Endovasc Surg.* 1995;9(4):421-425.
19. Kapural L, Narouze S, Janicki TI, Mekhail N. Spinal cord stimulation is an effective treatment for the chronic intractable visceral pelvic pain. *Pain Med.* 2006;;7(5):440-443.
20. Kemler M, Barendse GA van Kleef M, et al. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. *N Engl J Med.* 2000;343(9):618-624.
21. Kemler M, de Vet H, Barendse G, et al. Effect of spinal cord stimulation for chronic complex regional pain syndrome Type I: five-year final follow-up of patients in a randomized controlled trial. *J Neurosurg.* 2008;108(2):292-298.

22. Kemler M, de Vet H, Barendse G, et al. Spinal cord stimulation for chronic reflex sympathetic dystrophy – five year follow-up. *N Engl J Med.* 2006;354(22):2394-2396.
23. Kemler M, de Vet H, Barendse G, et al. The effect of spinal cord stimulation in patients with chronic reflex sympathetic dystrophy: two years' follow-up of the randomized controlled trial. *Ann Neurol.* 2004;55:13-18.
24. Kemler M, Furnee C. Economic evaluation of spinal cord stimulation for chronic reflex sympathetic dystrophy. *Neurology.* 2002;59(8):1203-1209.
25. Klomp H, Spincemaille G, Steyerberg E, et al. Spinal-cord stimulation in critical limb ischaemia: a randomised trial. ESES Study Group. *Lancet.* 1999;353(9158):1040-1044.
26. Kumar K, Hunter G, Demeria D. Spinal cord stimulation in treatment of chronic benign pain: challenges in treatment planning and present status, a 22-year experience. *Neurosurgery.* 2006;58(3):481-496; discussion 481-96.
27. Kumar K, Taylor RS, Jacques L, et al. Spinal cord stimulation versus conventional medical management for neuropathic pain: A multicenter randomized controlled trial in patients with failed back surgery syndrome. *Pain* 2007; 132(1-2): 179-88.
28. Latif O, Nedeljkovic S, Stevenson L. Spinal cord stimulation for chronic intractable angina pectoris: a unified theory on its mechanism. *Clin Cardiol.* 2001;24:533-541.
29. Magis D, Allena M, Bolla M, De Pasqua V, Remacle JM, Schoenen J. Occipital nerve stimulation for drug-resistant chronic cluster headache: a prospective pilot study. *Lancet Neurol* 2007; 6: 314–321.
30. Magis D, Gerardy PY, Remacle JM, Schoenen J. Sustained effectiveness of occipital nerve stimulation in drug-resistant chronic cluster headache. *Headache* 2011;51:1191–201.
31. Magis D, Schoenen J. Advances and challenges in neurostimulation for headaches. *Lancet Neurol* 2012;11:708–19.
32. Lipton R, Goadsby P, Cady R, Aurora SK, Grosberg B, Freitag F et al. PRISM study: occipital nerve stimulation for treatment-refractory migraine. *Cephalalgia* 2009;29:30.
33. Mailis-Gagnon A, Furlan A, Sandoval J, Taylor R, Spinal cord stimulation for chronic pain,
34. *Cochrane Database Syst Rev.* 2004;3:CD003783.
35. Mannheimer C, Eliasson T, Augustinsson LE, et al. Electrical stimulation versus coronary artery bypass surgery in severe angina pectoris: the ESBY study. *Circulation.* 1998;97:1157-1163.
36. Marin JC, Goadsby P. Response of SUNCT (short-lasting unilateral neuralgiform headaches with conjunctival injection and tearing), SUNA (short-lasting unilateral neuralgiform headaches with autonomic symptoms) and primary stabbing headaches to occipital nerve stimulation (ONS). *Neurology* 2010;74:P04.006, abstract.
37. McGreevy K, Williams K. Contemporary insights into painful diabetic neuropathy and treatment with spinal cord stimulation. *Curr Pain Headache Rep* 2012;16:43–49.
38. National Institute for Health and Care Excellence (NICE). Pain (chronic neuropathic or ischaemic) - spinal cord stimulation: final appraisal determination. September 2008.
39. North R, Calkins S, Campbell D, et al. Automated, patient-interactive, spinal cord stimulator adjustment: a randomized controlled trial. *Neurosurgery.* 2003;52(3):572-580.
40. North RB, Campbell JN, James CS, et al. Failed back surgery syndrome: Five year followup in 102 patients undergoing repeated operation. *Neurosurgery* 1991; 28; 28: 685-91.
41. North R, Kidd D, Farrokhi F, Piantadosi S. Spinal cord stimulation versus repeated lumbosacral spine surgery for chronic pain: a randomized, controlled trial. *Neurosurgery.* 2005;56(1):98-106.
42. North R, Kidd D, Lee M, et al. A prospective, randomized study of spinal cord stimulation versus reoperation for failed back surgery syndrome: initial results. *Stereotact Funct Neurosurg.* 1994;62:267-272.
43. North R, Kidd H, Piantadosi S. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a prospective, randomized study design. *Acta Neurochir.* 1995;64(Suppl):106-8.
44. North R, Kidd D, Shipley J, Taylor R. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a cost effectiveness and cost utility analysis based on a randomized, controlled trial. *Neurosurgery.* 2007;61(2):361-368.

45. Pedrini L, Magnoni F. Spinal cord stimulation for lower limb ischemic pain treatment. *Interact Cardiovasc Thorac Surg*. 2007;6(4):495-500.
46. Perruchoud C, Eldabe S, Batterham AM, et al. Analgesic efficacy of high-frequency spinal cord stimulation: A randomized double-blind placebo controlled study. *Neuromodulation* 2013; 16(4): 363-9.
47. Reed K. Peripheral neuromodulation and headaches: history, clinical approach, and considerations on underlying mechanisms. *Curr pain Headache Rep* 2013;17:305.
48. Reed KL, Black SB, Banta CJ 2nd, Will KR. Combined occipital and supraorbital neurostimulation for the treatment of chronic migraine headaches: initial experience. *Cephalalgia* 2010;30:260–71.
49. Saper JR, Dodick DW, Silberstein SD, McCarville S, Sun M, Goadsby PJ. Occipital nerve stimulation for the treatment of intractable chronic migraine headache: ONSTIM feasibility study. *Cephalalgia* 2011; 31:271–85.
50. Schwedt TJ, Dodick DW, Trentman TL, Zimmerman RS. Response to occipital nerve block is not useful in predicting efficacy of occipital nerve stimulation. *Cephalalgia* 2007; 27: 271–274.
51. Shanahan P, Watkins L, Matharu M. Treatment of medically intractable short--- lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT) and short---lasting unilateral neuralgiform headache attacks with autonomic symptoms (SUNA) with occipital nerve stimulation (ONS) in 6 patients. *Cephalalgia* 2009;29:150.
52. Silberstein S, Dodick D, Saper J, Huh B, Slavin KV, Sharan A et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: Results from a randomized, multicenter, double---blinded, controlled study. *Cephalalgia* 2012;32(16):1165–79.
53. Skaribas I, Calvillo O, Delikanaki---Skaribas E. Occipital peripheral nerve stimulation in the management of chronic intractable occipital neuralgia in a patient with neurofibromatosis type 1: a case report. *Journal of Medical Case Reports* 2011;5:174.Lambru G, Matharu M. Occipital nerve stimulation in primary headache syndromes. *Ther Adv Neurol Disord* 2012;5(1):57---67.
54. Slangen R, Pluijms WA, Faber CG, Dirksen CD, Kessels AG, van Kleef M. Sustained effect of spinal cord stimulation on pain and quality of life in painful diabetic peripheral neuropathy. *Br J Anaesth* 2013;111:1030– 1031.
55. Slangen R, Schaper NC, Faber CG, et al. Spinal cord stimulation and pain relief in painful diabetic peripheral neuropathy: a prospective two---center randomized controlled trial. *Diabetes Care* 2014;37:3016–3024.
56. Stancák A, Kozák J, Vrba I, et al. Functional magnetic resonance imaging of cerebral activation during spinal cord stimulation in failed back surgery syndrome patients. *Eur J Pain*. 2008;12(2):137-48.
57. Stanton-Hicks M, Burton A, Bruehl S, et al. An updated interdisciplinary clinical pathway for CRPS: report of an expert panel. *Pain Pract*. 2005; 84(3):S4-S16.
58. Sundaraj S, Johnstone C, Noore F, et al. Spinal cord stimulation: a seven-year audit. *J Clin Neurosci*. 2005 Apr;12(3):264-270.
59. Taylor R, Buyten J, Buchser E. Spinal cord stimulation for complex regional pain syndrome: A systematic review of the clinical and cost-effectiveness literature and assessment of prognostic factors. *Eur J Pain*. 2006;10(2):91-101.
60. Taylor RS, Desai MJ, Rigoard P, Taylor RJ. Predictors of pain relief following spinal cord stimulation in chronic back and leg pain and failed back surgery syndrome: A systemic review and meta-regression analysis. *Pain Pract* 2014; 14(6): 489-505.
61. Taylor R, Taylor R. Spinal cord stimulation for failed back surgery syndrome: a decision-analytic model and cost-effectiveness analysis. *Int J Technol Assess Health Care*. 2005;21(3):351-358.
62. TenVaarwerk I, Jessurun G, DeJongste M, et al. Clinical outcome of patients treated with spinal cord stimulation for therapeutically refractory angina pectoris. *The Working Group on Neurocardiology. Heart*. 1999;82(1):82-88.
63. Tesfaye S, Watt J, Benbow S, et al. Electrical spinal-cord stimulation for painful diabetic peripheral neuropathy. *Lancet*. 1996;348:1698-1701.
64. Ubbink D, Vermeulen H. Spinal cord stimulation for critical leg ischemia: a review of effectiveness and optimal patient selection. *J Pain Symptom Manage*. 2006;31:S30-S35.

66. Ubbink D, Vermeulen H. Spinal cord stimulation for non-reconstructable chronic critical leg ischaemia. *Cochrane Database Syst Rev.* 2005;CD004001.
67. van Beek M, Slangen R, Schaper N, Faber C, Joosten E, Dirksen C, van Dongen R, Kessels A, van Kleef M. Sustained treatment effect of spinal cord stimulation in painful diabetic peripheral neuropathy: 24-month follow-up of a prospective two-center randomized controlled trial. *Diabetes Care* 2015;38:e132-e134.
68. VanBuyten JP, Al-Kaisy A, Smet I, Palmisani S, Smith T. High-frequency spinal cord stimulation for the treatment of chronic back pain patients: Results of a prospective multicenter European clinical study. *Neuromodulation* 2013; 16(1): 59-65.
69. Weiner RL, Reed KL (1999) Peripheral neurostimulation for control of intractable occipital neuralgia. *Neuromodulation* 2:217-22.
70. Workloss Data Institute. *Official Disability Guidelines* 2008.