



# CLINICAL GUIDELINES

---

## Interventional Pain Management Services

Version 2.0.2019

Effective August 1, 2019



<b>Interventional Pain Management</b>	
<b>CMM-200: Epidural Steroid Injections (ESI)</b>	<b>3</b>
<b>CMM-201: Facet Joint Injections/Medial Branch Blocks</b>	<b>17</b>
<b>CMM-202: Trigger Point Injections</b>	<b>24</b>
<b>CMM-203: Sacroiliac Joint Injections</b>	<b>32</b>
<b>CMM-204: Prolotherapy</b>	<b>38</b>
<b>CMM-207: Epidural Adhesiolysis</b>	<b>41</b>
<b>CMM-208: Radiofrequency Joint Ablations/Denervations</b>	<b>45</b>
<b>CMM-209: Regional Sympathetic Blocks</b>	<b>51</b>
<b>CMM-210: Implantable Intrathecal Drug Delivery Systems</b>	<b>57</b>
<b>CMM-211: Spinal Cord Stimulators</b>	<b>64</b>
<b>CMM-308: Thermal Intradiscal Procedures</b>	<b>65</b>
<b>CMM-310: Manipulation of the Spine Under Anesthesia</b>	<b>70</b>
<b>CMM-400: Anesthesia Services for Interventional Pain Procedures</b>	<b>73</b>

**CMM-200: Epidural Steroid Injections (ESI)**

<b>CMM-200.1: Definitions</b>	<b>4</b>
<b>CMM-200.2: General Guidelines</b>	<b>5</b>
<b>CMM-200.3: Indications: Selective Nerve Root Block (SNRB)</b>	<b>6</b>
<b>CMM-200.4: Indications: Epidural Steroid Injections (Transforaminal, Interlaminar, or Caudal)</b>	<b>6</b>
<b>CMM-200.5: Non-Indications: SNRB</b>	<b>7</b>
<b>CMM-200.6: Non-Indications: ESI</b>	<b>8</b>
<b>CMM-200.7: Procedure (CPT®) Codes</b>	<b>9</b>
<b>CMM-200.8: References</b>	<b>10</b>

## **CMM-200.1: Definitions**

- **Transforaminal epidural steroid injection (TFESI)** is a therapeutic injection of contrast (absent allergy to contrast) performed at a single or multiple spinal levels, followed by the introduction of a corticosteroid and possibly a local anesthetic by inserting a needle into the neuroforamen under fluoroscopic or computed tomography (CT) guidance.
- **Selective Nerve Root Block (SNRB)** is a diagnostic injection of contrast (absent allergy to contrast) of a single nerve root to assist with surgical planning, followed by the introduction of a local anesthetic by inserting a needle into the neuroforamen under fluoroscopic or computed tomography (CT) guidance. SNRBs are erroneously referred to as transforaminal epidural steroid injection (TFESI), although technically SNRBs involve the introduction of anesthetic only and are used for diagnostic purposes.
  - ◆ Selective nerve root blocks (SNRBs) performed for the purpose of treating pain (i.e., repeat SNRB at the same level) may be termed therapeutic selective nerve root blocks. There is insufficient evidence to support the clinical utility of therapeutic selective nerve root blocks (SNRBs).
- **Interlaminar epidural steroid injection (ILESI)** is an injection of contrast (absent allergy to contrast), followed by the introduction of a corticosteroid and possibly a local anesthetic into the epidural space of the spine either through a paramedian or midline interlaminar approach under fluoroscopic guidance.
- **Caudal epidural steroid injection (CESI)** is an injection of contrast (absent allergy to contrast), followed by the introduction of corticosteroids and possibly a local anesthetic into the epidural space of the spine by inserting a needle through the sacral hiatus under fluoroscopic guidance into the epidural space at the sacral canal.
- **Radiculopathy**, for the purpose of this policy, is defined as the presence of pain, dysaesthesia(s), or paraesthesia(s) reported by the individual in a specified dermatomal distribution of an involved named spinal root(s) causing significant functional limitations (i.e., diminished quality of life and impaired, age-appropriate activities of daily living), and **EITHER** of the following:
  - ◆ Documentation of **ONE or MORE** of the following, concordant with nerve root compression of the involved named spinal root(s) demonstrated on a detailed neurologic examination within the prior three (3) months:
    - Loss of strength of specific named muscle(s) or myotomal distribution(s)
    - Altered sensation to light touch, pressure, pin prick or temperature in the sensory distribution
    - Diminished, absent or asymmetric reflex(es)
  - ◆ Documentation of **EITHER** of the following performed within the prior 12 months:
    - A concordant radiologist's interpretation of an advanced diagnostic imaging study (MRI or CT) of the spine demonstrating compression of the involved named spinal nerve root(s)
    - Electrodiagnostic studies (EMG/NCV's) diagnostic of nerve root compression of the involved named spinal nerve root(s).

- **Radicular pain** is pain which radiates to the extremity along the course of a spinal nerve root, typically resulting from compression, inflammation and/or injury to the nerve root.
- **Radiculitis** is defined, for the purpose of this policy, as radicular pain without objective neurological findings on physical examination.
- **Spinal stenosis** refers to the narrowing of the spinal canal usually due to spinal degeneration that occurs with aging. It may also be the result of spinal disc herniation, osteoarthritis, or a tumor. Lumbar spinal stenosis results in low back pain as well as pain or abnormal sensations in the legs, thighs, feet or buttocks, or loss of bladder and bowel control. Neurogenic claudication is often a clinical condition that results from spinal stenosis.

### **CMM-200.2: General Guidelines**

- The determination of medical necessity for the performance of a selective nerve root block (SNRB) or a therapeutic epidural steroid injection is always made on a case-by-case basis.
- **Please note:** this guideline does not apply to epidural injections administered for obstetrical or surgical epidural anesthesia.
- An epidural steroid injection should be performed with the use of fluoroscopic or CT guidance and the injection of a contrast, with the exception of an emergent situation or when fluoroscopic/CT guidance or the injection of contrast is contraindicated (e.g., pregnancy).
- The use of an indwelling catheter to administer a continuous infusion/intermittent bolus should be limited to use in a hospital setting only. It is inappropriate to represent the use of a catheter for single episode injection(s) that is/are commonly performed in an outpatient setting as an indwelling catheter for continuous infusion/intermittent bolus.
- There is insufficient scientific evidence to support the scheduling of a “series-of-three” injection in either a diagnostic or therapeutic approach. The medical necessity of subsequent injections should be evaluated individually and be based on the response of the individual to the previous injection with regard to clinically relevant sustained reductions in pain, decreased need for medication and improvement in the individual’s functional abilities.
- When performing transforaminal epidural steroid injections (TFESIs) or selective nerve root blocks (SNRBs), no more than two (2) nerve root levels should be injected during the same session/procedure.
- When medical necessity criteria is met, a total of three (3) epidural steroid injections (ESIs) per episode of pain, per region may be performed in six (6) months, not to exceed four (4) epidural steroid injections (ESIs) per region in 12 months.
- Additionally, when medical necessity criteria are met for a cervical/thoracic interlaminar (ILESIs) and/or a cervical/thoracic transforaminal epidural steroid injection (TFESI), advanced diagnostic imaging should be performed within 12 months prior to the injection.

### **CMM-200.3: Indications: Selective Nerve Root Block (SNRB)**

- A diagnostic selective nerve root block (SNRB), performed at a single nerve root, involving the introduction of anesthetic only, is considered **medically necessary** when attempting to establish the diagnosis of radicular pain (including radiculitis) or radiculopathy when the diagnosis remains uncertain after standard evaluation (neurologic examination, radiological studies and electrodiagnostic studies) in **ANY** of the following clinical situations:
  - ◆ When the physical signs and symptoms differ from that found on imaging studies
  - ◆ When there is clinical evidence of multi-level nerve root pathology
  - ◆ When the clinical presentation is suggestive, but not typical for both nerve root and peripheral nerve or joint disease involvement
  - ◆ When the clinical findings are consistent with radiculopathy in a dermatomal distribution, but the imaging studies do not corroborate the findings (positive straight leg raise test)
  - ◆ When the individual has had previous spinal surgery
  - ◆ For the purposes of surgical planning.
- A diagnostic selective nerve root block (SNRB) at a spinal level other than the initial level is considered **medically necessary** when **ALL** of the following criteria are met:
  - ◆ A response to the prior block of less than 80% relief based on the injectate utilized
  - ◆ Evidence of multilevel pathology
  - ◆ It has been at least 7 days since the prior block

### **CMM-200.4: Indications: Epidural Steroid Injections (Transforaminal, Interlaminar, or Caudal)**

- An epidural steroid injection (ESI) is considered **medically necessary** for **ANY** of the following:
  - ◆ Treatment of presumed radiculopathy when there has been failure of at least six (6) weeks of conservative treatment (e.g., exercise, physical methods including physical therapy and/or chiropractic care, nonsteroidal anti-inflammatory drugs [NSAID's] and/or muscle relaxants).
  - ◆ Treatment of presumed radiculitis or radicular pain when **ALL** of the following criteria are met:
    - Radicular pain, with or without motor weakness, which follows a specified dermatomal distribution of an involved named spinal root(s)
    - A positive straight leg raise, crossed leg raise, and/or Spurling's
    - Failure of at least six (6) weeks of conservative treatment (e.g., exercise, physical methods including physical therapy and/or chiropractic care, NSAID's and/or muscle relaxants).
  - ◆ An initial trial when there is evidence of symptomatic spinal stenosis and **ALL** of the following criteria are met:
    - Diagnostic evaluation has ruled out other potential causes of pain
    - MRI or CT with or without Myelography within the past twelve (12) months demonstrates moderate to severe spinal stenosis at the level to be treated
    - Significant functional limitations resulting in diminished quality of life and impaired, age-appropriate activities of daily living.

- Failure of at least four (4) weeks of conservative treatment (e.g., exercise physical methods including physical therapy and/or chiropractic care, NSAIDS, and/or muscle relaxants)
- A transforaminal epidural steroid injection (TFESI) in addition to an intra-articular facet joint injection with synovial cyst aspiration is considered **medically necessary** when **BOTH** of the following criteria are met:
  - ◆ Advanced diagnostic imaging studies (e.g., MRI CT, CT myelogram) confirm compression or displacement of the corresponding nerve root by a facet joint synovial cyst
  - ◆ Clinical correlation with the individual's signs and symptoms of radicular pain or radiculopathy, based on history and physical examination.
- A repeat epidural steroid injection (ESI) is considered **medically necessary** when at least **TWO** of the following criteria are met for two or more week's duration:
  - ◆ 50% or greater pain relief
  - ◆ Increase in the level of function/physical activity (e.g., return to work)
  - ◆ Reduction in the use of pain medication and/or additional medical services such as physical therapy/chiropractic case

#### **CMM-200.5: Non-Indications: SNRB**

- Diagnostic selective nerve root blocks (SNRBs) are considered **not medically necessary** for any other indication.
- Therapeutic selective nerve root blocks (SNRBs) (i.e., a repeat SNRB at the same level) are considered **experimental, investigational, or unproven** for any indication.
- A diagnostic selective nerve root block (SNRB) at a spinal level other than the initial level is considered **not medically necessary** for **ALL** of the following:
  - ◆ An adequate response to the first block, as determined by the injectate utilized
  - ◆ An absence of multilevel pathology when the first injection is performed under fluoroscopy/CT guidance using contrast
  - ◆ Repeating diagnostic selective nerve root blocks (SNRBs) more frequently than every seven (7) days



### **CMM-200.6: Non-Indications: ESI**

- An epidural steroid injection (ESI) performed with ultrasound guidance is considered **experimental, investigational, or unproven**.
- An epidural steroid injection is considered **not medically necessary** for **ALL** of the following:
  - ◆ When performed without imaging guidance (i.e., CT, fluoroscopy)
  - ◆ Transforaminal epidural steroid injection (TFESI) performed at more than two (2) nerve root levels during the same session/procedure.
  - ◆ An interlaminar epidural steroid injection (ILES), performed at more than a single level during the same session/procedure
  - ◆ Epidural steroid injection (ESI) administered in the same region as other spinal injections on the same day of service with the exception of an epidural steroid injection performed with an intra-articular facet joint injection with synovial cyst aspiration in accordance with criteria in CMM-200.4 above.
  - ◆ Performed in isolation (i.e., without the individual participating in an active rehabilitation program/home exercise program/functional restoration program)
  - ◆ Repeating epidural steroid injections more frequently than every 14 days
  - ◆ More than three (3) epidural steroid injections (ESIs) per episode of pain, per region in 6 months
  - ◆ More than four (4) epidural steroid injections (ESIs) per region, per 12 months
  - ◆ For axial spinal pain (i.e., absence of radiculopathy, myelopathy, myeloradiculopathy)
  - ◆ A caudal epidural steroid injection (CESI) for levels above L4-L5 without supporting clinical rationale for use of alternative approaches (e.g., translaminar, transforaminal)



## CMM-200.7: Procedure (CPT®) Codes

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only.

Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
62320	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance
62321	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (i.e., fluoroscopy or CT)
62322	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance
62323	Injection(s), of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); with imaging guidance (i.e., fluoroscopy or CT)
62324	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance
62325	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (i.e., fluoroscopy or CT)
62326	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance
62327	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (e.g., anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); with imaging guidance (i.e., fluoroscopy or CT)
64479	Injection(s), anesthetic agent and/or steroid, transforaminal epidural; with imaging guidance (fluoroscopy or CT); cervical or thoracic, single level
+64480	Injection(s), anesthetic agent and/or transforaminal epidural with imaging guidance (fluoroscopy or CT); cervical or thoracic, each additional level (List separately in addition to code for primary procedure)
64483	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); lumbar or sacral, single level
+64484	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with imaging guidance (fluoroscopy or CT); lumbar or sacral, each additional level (List separately in addition to code for primary procedure)

Epidural Steroid Injections (ESI)

CPT®	Codes Considered Experimental, Investigational or Unproven
0228T	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with ultrasound guidance, cervical or thoracic; single level
0229T	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with ultrasound guidance, cervical or thoracic; each additional level (List separately in addition to code for primary procedure)
0230T	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with ultrasound guidance, lumbar or sacral; single level
0231T	Injection(s), anesthetic agent and/or steroid, transforaminal epidural, with ultrasound guidance, lumbar or sacral; each additional level (List separately in addition to code for primary procedure)

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.

### **CMM-200.8: References**

1. Ackerman WE 3rd, Ahmad M. The efficacy of lumbar epidural steroid injections in patients with lumbar disc herniations. *Anesth Analg* 2007; 104:1217-1222.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*, 2nd Ed. 2008.
3. Albert H, Manniche C. The Efficacy of Systematic Active Conservative Treatment for Patients with Severe Sciatica. A Single-Blind, Randomized, Clinical, Controlled Trial. *Spine*. Vol 37, 7. 2012
4. Akuthota V, Bogduk N, Easa J, O'Brien D, Patel A, Prather H, Sharma A, Standaert C, Summers J, Lumbar Transforaminal Epidural Steroid Injections Review & Recommendation Statement. North American Spine Society, January 2013, 28-30.
5. Allen T, Tatli Y, Lutz G. Fluoroscopic percutaneous lumbar zygapophyseal joint cyst rupture: a clinical outcome study. *Spine J*. 2009 May;9(5): 387-95.
6. Amoretti N, Huwart L, Foti P, Boileau P, Amoretti M, Pellegrin A, Marcy P, Hauger O. Symptomatic lumbar facet joint cysts treated by CT-guided ntracystic and intra-articular steroid injections. *Eur Radiol* 2012 Dec; (12): 2836-40
7. Amr YM. Effect of addition of epidural ketamine to steroid in lumbar radiculitis: One-year follow-up. *Pain Physician*. 2011;14:475-481.
8. Anderberg L, Annertz M, Persson L, et al. Transforaminal steroid injections for the treatment of cervical radiculopathy: a prospective and randomised study. *Eur Spine J*. 2007;16(3):321-328.
9. Arden N, Price C, Reading I, et al. A multicentre randomized controlled trial of epidural corticosteroid injections for sciatica: the WEST study. *Rheumatology (Oxford)*. 2005; 44:1399-1406.
10. Arden NK, Price C, Reading I, Stubbing J, Hazelgrove J, Dunne C, et al; WEST Study Group. A multicentre randomized controlled trial of epidural corticosteroid injections for sciatica: The WEST study. *Rheumatology (Oxford)* 2005;44:1399-1406.
11. Becker C, Heidersdorf S, Drewlo S, de Rodriguez SZ, Krämer J, Willburger RE. Efficacy of epidural perineural injections with autologous conditioned serum for lumbar radicular compression: An investigator-initiated, prospective, double-blind, reference controlled study. *Spine (Phila Pa 1976)* 2007;32:1803-1808.
12. Benzon HT, Huntoon MA, Rathmell JP. Improving the safety of epidural steroid injections. *JAMA*. 2015 May 5;313(17):1713-4. doi: 10.1001/jama.2015.2912.
13. Blankenbaker D, De Smet A, Stanczak J, Fine J. Lumbar radiculopathy: treatment with selective lumbar nerve blocks—comparison of effectiveness of triamcinolone and betamethasone injectable suspensions. *Radiology*. 2005;237:738-741.
14. Botwin K, Baskin M, Rao S. Adverse effects of fluoroscopically guided interlaminar thoracic epidural steroid injections. *Am J Phys Med Rehabil*. 2006;85:14-23.
15. Botwin K, Gruber R, Bouchlas C, et. al. Fluoroscopically guided lumbar transforaminal epidural steroid injections in degenerative lumbar stenosis: an outcome study. *Am J Phys Med Rehabil*. 2002;81(12):898-905.

16. Botwin K, Sakalkale D. Epidural steroid injections in the treatment of symptomatic lumbar spinal stenosis associated with epidural lipomatosis. *Am J Phys Med Rehabil.* 2004;83:926-993.
17. Buttermann G. Treatment of lumbar disc herniation: epidural steroid injection compared with discectomy. A prospective, randomized study. *J Bone Joint Surg Am.*2004;86-A(4):670-679.
18. Candido KD, Rana MV, Sauer R, Chupatanakul L, Tharian A, Vasic V, Knezevic NN. Concordant pressure paresthesia during interlaminar lumbar epidural steroid injections correlates with pain relief in patients with unilateral radicular pain. *Pain Physician.* 2013;16:497-511.
19. Carette S, Leclaire R, Marcoux S, Morin F, Blaise GA, St-Pierre A, Truchon R, Parent F, Levesque J, Bergeron V, Montminy P, Blanchette C. Epidural corticosteroid injections for sciatica due to herniated nucleus pulposus. *N Engl J Med.* 1997;336:1634-1640.
20. Chang Chien GC, Knezevic NN, McCormick Z, Chu SK, Trescot Am, Candido KD. Transforaminal versus interlaminar approaches to epidural steroid injections: A systematic review of comparative studies for lumbosacral radicular pain. *Pain Physician.* 2014;17:E509-E524.
21. Chou R, Loeser J, Owens D, Rosenquist R, Atlas S, Baisden J, Carragee E, Grabois M, Murphy D, Resnick D, Stanos S, Shafer W, Wall E. Interventional Therapies, Surgery, and Interdisciplinary Rehabilitation for Low Back Pain. An Evidence Based Clinical Practice Guideline From the American Pain Society. *Spine* 2009 34(10) 1066-77.
22. Cohen SP, Hanling S, Bicket MC, White RL, Veizi E, Kurihara C, Zhao Z, Hayek S, Guthmiller KB, Griffith SR, Gordin V, White MA, Vorobeychik Y, Pasquina PF. Epidural steroid injections compared with gabapentin for lumbosacral radicular pain: Multicenter randomized double blind comparative efficacy study. *BMJ* 2015; 350:h1748.
23. Cohen S, Hayek S, Semenov Y, Pasquina P, White R, Veizi E, Huang J, Kurihara C, Zhao Z, Guthmiller K, Griffith S, Verdun A, Giampetro D, Vorobeychik Y. Epidural steroid injections, conservative treatment, or combination treatment for cervical radicular pain: a multicenter, randomized, comparative-effectiveness study. *Anesthesiology.* 2014 Nov; 121(5): 1045 – 55.
24. Cooper G, Lutz G, Boachie-Adjei O, Lin J. Effectiveness of transforaminal epidural steroid injections in patients with degenerative lumbar scoliotic stenosis and radiculopathy. *Pain Physician.* 2004; 7:311-317.
25. Dashfield A, Taylor M, Cleaver J, Farrow D. Comparison of caudal steroid epidural with targeted steroid placement during spinal endoscopy for chronic sciatica: a prospective, randomized, double-blind trial. *Br J Anaesth.* 2005; 94:514-559.
26. Datta R, Upadhyay KK. A randomized clinical trial of three different steroid agents for treatment of low backache through the caudal route. *Med J Armed Forces India.* 2011;67:25-33.
27. Delitto A, George S, VanDillen L, Whitman J, Sowa G, Shekelle P, Denninger T, Godges J. Low Back Pain Clinical Practice Guidelines Linked to the International Classification of Functioning, Disability, and Health from the Orthopaedic Section of the American Physical Therapy Association. *J Orthop Sports Phys Ther.* 2012;42(4):A1 – A 57. Affirmed by the the American Academy of Physical Medicine and Rehabilitation Board of Governors.
28. DePalma M, Bhargava A, Slipman C. A critical appraisal of the evidence for selective nerve root injection in the treatment of lumbosacral radiculopathy. *Arch Phys Med Rehabil.* 2005;86(7):1477-1483.
29. Donelson R, Long A, Spratt K, Fung, T. Influence of directional preference on two clinical dichotomies: acute versus chronic pain and axial low back pain versus sciatica. *PM R.* 2012;4:667.
30. Dreyfuss P, Baker R, Bogduk N. Comparative effectiveness of cervical transforaminal injections with particulate and nonparticulate corticosteroid preparations for cervical radicular pain. *Pain Med.* 2006;7:237-242.
31. Eckel TS, Bartyński WS. Epidural steroid injections and selective nerve root blocks. *Tech Vasc Interv Radiol.* 2009 Mar 12(1): 11 – 21.
32. Friedly JL, Comstock BA, Turner JA, et al. A randomized trial of epidural glucocorticoid injections for spinal stenosis. *N Engl J Med* 2014; 371:11-21.
33. Fukusaki M, Kobayashi I, Hara T, Sumikawa K. Symptoms of spinal stenosis do not improve after epidural steroid injection. *Clin J Pain* 1998;14:148-151.
34. Furman M, Butler S, Kim R, Mehta A, Simon J, Patel R, Lee T, Reeves R (2012). Injectate volumes needed to reach specific landmarks in S1 transforaminal epidural steroid injections. *Pain Medicine.* 13(10):1265-1274.

35. Furman M, Mehta A, Kim R, Simon J, Patel R, Lee T, Reeves R (2010). Injectate volumes to reach specific landmarks in lumbar transforaminal epidural steroid injections. *PM&R*, 2(7), 625-635.
36. Furman MB, Lee TS, Mehta A, Simon JI, Cano WG (2008). Contrast flow selectivity during transforaminal epidural steroid injections. *Pain Physician*, 11(6), 855-861.
37. Ghai B, Bansal D, Kay JP, Vadaje KS, Wig J. Transforaminal versus parasagittal interlaminar epidural steroid injection in low back pain with radicular pain: A randomized, double-blind, active-control trial. *Pain Physician* 2014; 17:277-290.
38. Ghahreman A, Ferch R, Bogduk N. The efficacy of transforaminal injection of steroids for the treatment of lumbar radicular pain. *Pain Med* 2010; 11:1149-1168.
39. Ghai B, Kumar K, Bansal D, Dhatt SS, Kanukula R, Batra YK. Effectiveness of parasagittal interlaminar epidural local anesthetic with or without steroid in chronic lumbosacral pain: A randomized, double-blind clinical trial. *Pain Physician* 2015;18:237-248.
40. Hagen K, Hilde G, Jamtvedt G, Winnem M. The cochrane review of advice to stay as active as a single treatment for low back pain and sciatica. *Spine*. 2002 Aug 15; 27(16): 1736 – 41.
41. Hooten W, Cohen S. Evaluation and treatment of low back pain: a clinically focused review for primary care specialists. *Mayo Clin Proc*. 2015 Dec;90(12): 1699-718.
42. Huang R, Shapiro G, Lim M, Set al. Cervical epidural abscess after epidural steroid injection. *Spine*. 2004; 29:E7-E9.
43. Huda N, Bansal P, Gupta SM, Ruhela A, Rehman M, Afzal M. The efficacy of epidural depomethylprednisolone and triamcinolone acetate in relieving the symptoms of lumbar canal stenosis: A comparative study. *J Clin Diagn Res* 2010;4:2843-2847.
44. Jeong HS, Lee JW, Kim SH, Myung JS, Kim JH, Kang HS. Effectiveness of transforaminal epidural steroid injection by using a preganglionic approach: A prospective randomized controlled study. *Radiology* 2007;245:584-590.
45. Kennedy DJ, Plastaras C, Casey E, Visco CJ, Rittenberg JD, Conrad B, Sigler J, Dreyfuss P. Comparative effectiveness of lumbar transforaminal epidural steroid injections with particulate versus nonparticulate corticosteroids for lumbar radicular pain due to intervertebral disc herniation: a prospective, randomized, double-blind trial. *Pain Med*. 2014;15:548-55.
46. Kerezoudis P, Rinaldo L, Alvi MA, Hunt CL, Qu W, Maus TP, Bydon M. The Effect of Epidural Steroid Injections on Bone Mineral Density and Vertebral Fracture Risk: A Systematic Review and Critical Appraisal of Current Literature. *Pain Med*. 2018 Jan 2. doi: 10.1093/pm/pnx324.
47. Iversen T, Solberg TK, Romner B, Wilsgaard T, Twisk J, Anke A, Nygaard O, Hasvold T, Ingebrigtsen T. Effect of caudal epidural steroid or saline injection in chronic lumbar radiculopathy: multicentre, blinded, randomised controlled trial. *BMJ*. 2011;343:d5278.
48. Karppinen J, Malmivaara A, Kurunlahti M, Kyllönen E, Pienimäki T, Nieminen P, Ohinmaa A, Tervonen O, Vanharanta H. Periradicular infiltration for sciatica: A randomized controlled trial. *Spine (Phila Pa 1976)* 2001;26:1059-1067.
49. Kaye AD, Manchikanti L, Abdi S, et al. Efficacy of epidural injections in managing chronic spinal pain: A best evidence synthesis. *Pain Physician*. 2015; in press.
50. King W, Miller DC, Smith CC. Systemic Effects of Epidural Corticosteroid Injection. *Pain Med*. 2018 Feb 1;19(2):404-405.
51. Koh WU, Choi SS, Park SY, Joo EY, Kim SH, Lee JD, Shin JY, Suh JH, Leem JG, Shin JW. Transforaminal hypertonic saline for the treatment of lumbar lateral canal stenosis: A doubleblinded, randomized, active-control trial. *Pain Physician*. 2013;16:197-211.
52. Kreiner D, Hwang S, Easa J, Resnick D, Baisden J, Dougherty P, Fernand R, Ghiselli G, Haanna A, Lamer T, Lisi A, Mazanec D, Meagher R, Nucci R, Patel R, Sembrano J, Sharma A, Summers J, Taleghani C, Tontz W, Toton J. North American Spine Society Evidence Based Guideline for Multidisciplinary Spine Care. *Clinical Guidelines for Diagnosis and Treatment of Lumbar Disc Herniation with Radiculopathy*. 2012, page 30.
53. Kreiner D, Shaffer W, Baisden J, Gilbert T, Summers J, Toton J, Hwang S, Mendel R, Reitman C. Evidence – Based Clinical Guidelines for Multidisciplinary Spine Care. *Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis*. North American Spine Society 2011, 43-44. (43 references) Endorsed by the American Academy of Physical Medicine and Rehabilitation Board of Governors
54. Lee JH, An JH, Lee SH. Comparison of the effectiveness of interlaminar and bilateral transforaminal epidural steroid injections in treatment of patients with lumbosacral disc herniation and spinal stenosis. *Clin J Pain*. 2009;25:206-210.



55. Lee J, Kim S, Lee I, et al. Therapeutic effect and outcome predictors of sciatica treated using transforaminal epidural steroid injection. *AJR Am J Roentgenol.* 2006; 187:1427-1431. Lee K, Lin C, Hwang S, et al. Transforaminal periradicular infiltration guided by CT for unilateral sciatica—an outcome study. *Clin Imaging.* 2005; 29:211-214.
56. Lee JH, Kim DH, Shin KS, Park SJ, Lee GJ, Lee CH, Yang HS. Comparison of clinical efficacy of epidural injection with or without steroid in lumbosacral disc herniation: A systematic review and meta-analysis. *Pain Physician* 2018; in press.
57. Lipetz, J Pathophysiology of inflammatory degenerative and compressive radiculopathies. *PMR Clinics of North America.* 2002; Aug: 13(3): 439-49
58. Liu K, Liu P, Liu R, Wu X, Cai M. Steroid for epidural injection in spinal stenosis: A systematic review and meta-analysis. *Drug Des Devel Ther.* 2015;9:707–716
59. Liu J, Zhou H, Lu L, Li X, Jia J, Shi Z, Yao X, Wu Q, Feng S. The Effectiveness of Transforaminal Versus Caudal Routes for Epidural Steroid Injections in Managing Lumbosacral Radicular Pain: A Systematic Review and Meta-Analysis. *Medicine (Baltimore).* 2016 May;95(18):e3373.
60. Lutz G, Shen T. Fluoroscopically guided aspiration of a symptomatic lumbar zygapophyseal joint cyst: a case report. *Arch Phys Med Rehabil* 2002 Dec; 83(12); 1789-91.
61. Martha J, Swaim B, Wang D, Kim D, Hill D, Bode R, Schwartz C. Outcome of percutaneous rupture of lumbar synovial cysts: a case series of 101 patients. *Spine J* 2009 Nov 9(11) 899-904.
62. Machado LA, de Souza MS, Ferreira PH, Ferreira ML. The McKenzie method for low back pain: a systematic review of the literature with a meta-analysis approach. *Spine (Phila Pa 1976).* 2006;31(9):E254–E262.
63. Machado LA, Maher CG, Herbert RD, Clare H, McAuley JH. The effectiveness of the McKenzie method in addition to first-line care for acute low back pain: a randomized controlled trial. *BMC Med.* 2010;8:10.
64. Macvicar J, King W, Landers MH, Bogduk N. The effectiveness of lumbar transforaminal injection of steroids: A comprehensive review with systematic analysis of the published data. *Pain Med.* 2013; 14:14-28.
65. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain: Part II: Guidance and recommendations. *Pain Physician.* 2013;16:S49-S283.
66. Manchikanti L, Benyamin RM, Falco FJ, Kaye AD, Hirsch JA. Do epidural injections provide short- and long-term relief for lumbar disc herniation? A systematic review. *Clin Orthop Relat Res.* 2015;473:1940-1956.
67. Manchikanti L, Cash KA, McManus CD, Damron KS, Pampati V, Falco FJE. A randomized, double-blind controlled trial of lumbar interlaminar epidural injections in central spinal stenosis: 2-year follow-up. *Pain Physician.* 2015;18:79-92.
68. Manchikanti L, Cash KA, McManus CD, Pampati V. Fluoroscopic caudal epidural injections in managing chronic axial low back pain without disc herniation, radiculitis or facet joint pain. *J Pain Res.* 2012;5:381-390.
69. Manchikanti L, Cash KA, McManus CD, Pampati V, Benyamin RM. A randomized, double-blind, active-controlled trial of fluoroscopic lumbar interlaminar epidural injections in chronic axial or discogenic low back pain: Results of a 2-year follow-up. *Pain Physician.* 2013;16:E491- E504.
70. Manchikanti L, Cash KA, McManus CD, Pampati V, Benyamin R. Fluoroscopic lumbar interlaminar epidural injections in managing chronic lumbar axial or discogenic pain. *J Pain Res.* 2012; 5:301-311.
71. Manchikanti L, Cash KA, McManus CD, Pampati V, Fellows B. Results of 2-year follow-up of a randomized, double-blind, controlled trial of fluoroscopic caudal epidural injections in central spinal stenosis. *Pain Physician.* 2012;15:371-384.
72. Manchikanti L, Cash KA, Pampati V, Falco FJE. Transforaminal epidural injections in chronic lumbar disc herniation: A randomized, double-blind, active-control trial. *Pain Physician.* 2014;17:E489-E501.
73. Manchikanti L, Falco FJE, Pampati V, Hirsch JA. Lumbar interlaminar epidural injections are superior to caudal epidural injections in managing lumbar central spinal stenosis. *Pain Physician* 2014;17:E691-E702.
74. Manchikanti L, Cash KA, Pampati V, Malla Y. Fluoroscopic cervical epidural injections in chronic axial or disc-related neck pain without disc herniation, facet joint pain, or radiculitis. *J Pain Res.* 2012;5:227-236.

75. Manchikanti L, Cash KA, Pampati V, Malla Y. Two-year follow-up results of fluoroscopic cervical epidural injections in chronic axial or discogenic neck pain: A randomized, double-blind, controlled trial. *Int J Med Sci.* 2014;11:309-320.
76. Manchikanti L, Hirsch JA. An update on the management of chronic lumbar discogenic pain. *Pain Manag.* 2015;5:373-386.
77. Manchikanti L, Hirsch JA. Clinical management of radicular pain. *Expert Rev Neurother.* 2015;15:681-693.
78. Manchikanti L, Knezevic NN, Boswell MV, Kaye AD, Hirsch JA. Epidural injections for lumbar radiculopathy and spinal stenosis: A comparative systematic review and meta-analysis. *Pain Physician* 2016; E365-E410.
79. Manchikanti L, Nampiaparampil DE, Manchikanti KN, et al. Comparison of the efficacy of saline, local anesthetics, and steroids in epidural and facet joint injections for the management of spinal pain: A systematic review of randomized controlled trials. *Surg Neurol Int.* 2015;6:S194- S235.
80. Manchikanti L, Pampati V, Benyamin RM, Boswell MV. Analysis of efficacy differences between caudal and lumbar interlaminar epidural injections in chronic lumbar axial discogenic pain: Local anesthetic alone vs. local combined with steroids. *Int J Med Sci.* 2015;12:214-222.
81. Manchikanti L, Singh V, Cash KA, Pampati V, Damron KS, Boswell MV. Effect of fluoroscopically guided caudal epidural steroid or local anesthetic injections in the treatment of lumbar disc herniation and radiculitis: A randomized, controlled, double blind trial with a two year follow-up. *Pain Physician* 2012;15:273-286.
82. Manchikanti L, Singh V, Cash KA, Pampati V, Falco FJE. A randomized, double-blind, active control trial of the effectiveness of lumbar interlaminar epidural injections in disc herniation. *Pain Physician.* 2014; 7:E61-E74.
83. Manchikanti L, Singh V, Pampati V, Falco FJE, Hirsch JA. Comparison of the efficacy of caudal, interlaminar, and transforaminal epidural injections in managing lumbar disc herniation: Is one method superior to the other? *Korean J Pain.* 2015;28:11-21.
84. Manchikanti L, Staats PS, Nampiaparampil DE, Hirsch JA. What is the role of epidural injections in the treatment of lumbar discogenic pain: A systematic review of comparative analysis with fusion and disc arthroplasty. *Korean J Pain.* 2015;28:75-87. Matz P, Meagher R, Lamet T, Tontz W. Evidence – Based Clinical Guidelines for Multidisciplinary Spine Care. *Diagnosis and Treatment of Degenerative Lumbar Spondylolisthesis.* North American Spine Society 2014, 39. (87 references) Endorsed by the American Academy of Physical Medicine and Rehabilitation Board of Governors
85. McCormick Z, Cushman D, Casey E, Garvan C, Kennedy D, Plastares C. Factors associated with pain reduction after transforaminal epidural steroid injection for lumbosacral radicular pain. *Arch Phys Med Rehabil* 2014 Dec; 95(12): 2350-6.
86. Melfi R, Aprill C. Percutaneous puncture of zygapophyseal joint synovial cyst with fluoroscopic guidance. *Pain Med* 2005 Mar-0Apr;6(2):122-8.
87. Meng H, Fei Q, Wang B, Yang Y, Li D, Li J, Su N. Epidural injections with or without steroids in managing chronic low back pain secondary to lumbar spinal stenosis: a meta-analysis of 13 randomized controlled trials. *Drug Des Devel Ther* 2015; 9:4657-67.
88. Murakibhavi VG, Khemka AG. Caudal epidural steroid injection: A randomized controlled trial. *Evid Based Spine Care J.* 2011;2:19-26.
89. Ng L, Chaudhary N, Sell P. The efficacy of corticosteroids in periradicular infiltration for chronic radicular pain: A randomized, double-blind, controlled trial. *Spine (Phila Pa 1976).* 2005;30:857- 862.
90. North American Spine Society (NASS). NASS Coverage Policy Recommendation.: Cervical Epidural Injections and Diagnostic Spinal Nerve Blocks. Lumbar Epidural Injections Facet Joint Intervention. Copyright © 2016 North American Spine Society
91. Park KD, Lee J, Jee H, Park Y. Kambin triangle versus the supraneural approach for the treatment of lumbar radicular pain. *Am J Phys Med Rehabil.* 2012;91:1039-1050.
92. Park Y, Lee JH, Park KD, Ahn JK, Park J, Jee H. Ultrasound-guided vs. fluoroscopy-guided caudal epidural steroid injection for the treatment of unilateral lower lumbar radicular pain: A prospective, randomized, single-blind clinical study. *Am J Phys Med Rehabil.* 2013;92:575-586.
93. Parr AT, Diwan S, Abdi S. Lumbar interlaminar epidural injections in managing chronic low back pain and lower extremity pain: a systematic review. *Pain Physician* 2009 Jan-Feb; 12(1): 163 – 88.
94. Patel J, Schneider B, Smith C on behalf of SIS Patient Safety Committee. Intra-articular Corticosteroid Injections and hyperglycemia. 10/4/17.

95. Peng B, Wu, Li Z, Guo JWang X. Chemical Radiculitis. *Pain*. 2007; Jan (1-2): 11-6.
96. Pinto RZ, Maher CG, Ferreira ML, Hancock M, Oliveira VC, McLachlan AJ, Koes B, Ferreira PH. Epidural corticosteroid injections in the management of sciatica: A systematic review and meta-analysis. *Ann Intern Med*. 2012;157:865-877.
97. Pirbudak L, Karakurum G, Oner U, Gulec A, Karadasli H. Epidural corticosteroid injection and amitriptyline for the treatment of chronic low back pain associated with radiculopathy. *Pain Clinic*. 2003;15:247-253.
98. Rados I, Sakic K, Fingler M, Kapural L. Efficacy of interlaminar vs transforaminal epidural steroid injection for the treatment of chronic unilateral radicular pain: prospective, randomized study. *Pain Med*. 2011;12:1316-1321.
99. Riew KD, Yin Y, Gilula L, et al.. *J Bone Joint Surg Am*. 2000;82:1589-1593.
100. Riew K, Park J, Cho Y, et al. Nerve root blocks in the treatment of lumbar radicular pain. A minimum five-year follow-up. *J Bone Joint Surg Am*. 2006;88(8):1722-1725.
101. Sabers S, Ross S, Grogg B, Lauder T. Procedure-based nonsurgical management of lumbar zygapophyseal joint cyst-induced radicular pain. *Arch Phys Med Rehabil* 2005 Sep;86(9): 1767-71.
102. Saifuddin A, Mitchel R, Taylor B. Extradural inflammation associated with annular tears: Demonstration with gadolinium-enhanced lumbar spine MRI. *Eur Spine J*. 1999; 8 (1): 34-9.
103. Sasso R, Macadaeg K, Nordmann D, Smith M. Selective nerve root injections can predict surgical outcome for lumbar and cervical radiculopathy: comparison to magnetic resonance imaging. *J Spinal Disord Tech*. 2005;18:471-478.
104. Sayegh FE, Kenanidis EI, Papavasiliou KA, Potoupnis ME, Kirkos JM, Kapetanios GA. Efficacy of steroid and nonsteroid caudal epidural injections for low back pain and sciatica: A prospective, randomized, double-blind clinical trial. *Spine (Phila Pa 1976)*. 2009;34:1441-1447.
105. Schaufele M, Hatch L, Jones W. Interlaminar versus transforaminal epidural injections for the treatment of symptomatic lumbar intervertebral disc herniations. *Pain Physician*. 2006;9:361-366.
106. Shah RD, Cappiello D, Suresh S. Interventional procedures for chronic pain in children and adolescents: a review of the current evidence. *World Institute of Pain*. 2016: 359-369.
107. Shanthanna H, Busse JW, Thabane L, Paul J, Couban R, Choudhary H, Kaushal A, Suzumura E, Kim I, Harsha P. Local anesthetic injections with or without steroid for chronic non-cancer pain: a protocol for a systematic review and meta-analysis of randomized controlled trials. *Syst Rev* 2016; 5:18.
108. Slipman C, Lipetz J, DePalma M, Jackson H. Therapeutic selective nerve root block in the nonsurgical treatment of traumatically induced cervical spondylotic radicular pain. *Am J Phys Med Rehabil*. 2004; 83:446-454.
109. Slipman C, Lipetz J, Wakeshima Y, Jackson H. Nonsurgical treatment of zygapophyseal joint cyst-induced radicular pain. *Arch Phys Med Rehabil* 2000 Jul; 81(7): 973-7.
110. Southern D, Lutz GE, Cooper G, Barre L. Are fluoroscopic caudal epidural steroid injections effective for managing chronic low back pain? *Pain Physician* 2003 Apr;6(2): 167-72.
111. Spijker-Huiges A, Vermeulen K, Winters JC, van Wijhe M, van der Meer K. Epidural steroids for lumbosacral radicular syndrome compared to usual care: quality of life and cost utility in general practice. *Arch Phys Med Rehabil*. 2015 Mar;96(3):381.
112. Staal JB, de Bie R, de Vet HCW, Hildebrandt J, Nelemans P. Injection therapy for subacute and chronic low back pain: an updated Cochrane review. *Spine* 2009; 34(1):49.
113. Stout A. Epidural Steroid injection for low back pain. *Phys Med Rehabil Clin N Am*. 2010 Nov 21(4): 825 – 34.
114. Tafazal S, Ng L, Chaudhary N, Sell P. Corticosteroids in peri-radicular infiltration for radicular pain: A randomised double blind controlled trial: one year results and subgroup analysis. *Eur Spine J*. 2009;18:1220-1225.
115. Thomas E, Cyteval C, Abiad L, et al. Efficacy of transforaminal -versus interspinous corticosteroid injection in discal radiculalgia – A prospective, randomised, double-blind study. *Clin Rheumatol*.2003; 22:299-304.
116. Vad VB, Bhat AL, Lutz GE, Cammisa F. Transforaminal epidural steroid injections in lumbosacral radiculopathy: A prospective randomized study. *Spine (Phila Pa 1976)* 2002; 27:11-16.
117. Valat J, Genevay S, Marty M, Rozenberg S, Koes B. Sciatica. *Best Pract Res Clin Rheumatol*. 2010 Apr;24(2): 241-52.



118. Valat J, Giraudeau B, Rozenberg S, et al. Epidural corticosteroid injections for sciatica: a randomised, double blind, controlled clinical trial. *Ann Rheum Dis*. 2003;62:639-643.
119. Van Helvoirt H, Apeldoorn A, Ostelo R, Knol D, Arts M, Kamper S, van Tulder M. Transforaminal epidural steroid injections followed by mechanical diagnosis and therapy to prevent surgery for lumbar disc herniation. *Pain Med*. 2014 Jul; 15(7): 1100-8.
120. Wei G, Liang J, Chen B, Zhou C, Ru N, Chen J, Zhang F. Comparison of transforaminal versus interlaminar epidural steroid injection in low back pain with lumbosacral radicular pain: a meta-analysis of the literature. *Int Orthop*. 2016 Dec;40(12):2533-2545. Epub 2016 May 20.
121. Wilson-MacDonald J, Burt G, Griffin D, Glynn C. Epidural steroid injection for nerve root compression. A randomised, controlled trial. *J Bone Joint Surg Br* 2005; 87:352-355.
122. Woodward J, Herring S, Windsor R, ed. Lennard T. *Epidural Procedures in Spine Pain Management. Pain Procedures in Clinical Practice*. Second Edition. Hanley and Belfus Inc. Philadelphia. 2000, page 359.
123. Zhai J, Zhang L, Li M, Tian Y, Zheng W, Chen J, Huang T, Li X, Tian Z. Epidural injection with or without steroid in managing chronic low back and lower extremity pain: A meta-analysis of ten randomized controlled trials. *Int J Clin Exp Med*. 2015;8(6):8304-8316.
124. Zhai J, Zhang L, Li M, Tian Y, Zheng W, Chen J, Huang T, Li X, Tian Z. Epidural injection with or without steroid in managing chronic low back and lower extremity pain: A meta-analysis of ten randomized controlled trials. *Am J Ther*. 2017 May;24(3):e259-e269.

**CMM-201: Facet Joint Injections/Medial Branch Blocks**

<b>CMM-201.1: Definitions</b>	<b>18</b>
<b>CMM-201.2: General Guidelines</b>	<b>18</b>
<b>CMM-201.3: Indications</b>	<b>19</b>
<b>CMM-201.4: Non-Indications</b>	<b>20</b>
<b>CMM-201.5: Procedure (CPT®) Codes</b>	<b>20</b>
<b>CMM-201.6: References</b>	<b>21</b>

## **CMM-201.1: Definitions**

- **Facet Joint Injections/medial branch blocks** refer to the injection of local anesthetic and possibly a corticosteroid in the facet joint capsule or along the nerves supplying the facet joints from C2-3 to L5-S1. The injection/block applies directly to the facet joint(s) blocked and not to the number of nerves blocked that innervate the facet joint(s). Even though either procedure can be used to diagnose facet joint pain, a medial branch block is generally considered more appropriate. A diagnostic facet joint injection/medial branch block is considered positive when there is at least 80% relief of facet mediated pain for at least the expected minimum duration of the effect of the local anesthetic used.

## **CMM-201.2: General Guidelines**

- The determination of medical necessity for the performance of facet joint injections/medial branch blocks is always made on a case-by-case basis.
- Facet joint injections/medial branch blocks should only be performed for neck pain or low back pain in the absence of an untreated radiculopathy (with the exception of radiculopathy caused by a facet joint synovial cyst).
- A diagnostic facet joint injection/medial branch block may be performed to determine whether spinal pain originates in the facet joint or nerves innervating the facet joint. A second facet joint injection/medial branch block must be performed to confirm the validity of the clinical response of the initial injection and should only be performed with the intent that if successful, a radiofrequency joint denervation/ablation procedure (facet neurotomy, facet rhizotomy) would be considered as an option at the diagnosed level(s).
- More than two facet injections/medial branch blocks at the same level are considered to be therapeutic rather than diagnostic. Following a spinal fusion, a diagnostic facet joint injection/medial branch block may be performed immediately above or below the fused level if a prior injection/block was negative. There is a paucity of published scientific evidence supporting the use of therapeutic facet joint injections/medial branch blocks. Although limited, some anecdotal evidence supports a facet joint injection/medial branch block as an alternative treatment to a radiofrequency ablation/neurotomy for a subset of individuals when the initial facet joint injection/medial branch blocks has resulted in significant pain relief (i.e., > 50%) for at least 12 weeks following the facet joint injection/medial branch block and the individual is not a candidate for a radiofrequency joint denervation/ablation procedure. For this specific subset of individuals a repeat facet joint injection may be considered appropriate, although no sooner than six months from when the prior diagnostic injection was performed.
- It may be necessary to perform the facet joint injection/medial branch block at the same facet joint level(s) bilaterally, however, no more than three (3) facet joint levels should be injected during the same session/procedure.

- Facet joint injections/medial branch blocks are not without risk and can expose patients to potential complications that may be increased when a patient is sedated. As a result, when performing facet joint injections/medial branch blocks, the use of supplemental sedation in addition to local anesthesia is not required and not recommended.

### **CMM-201.3: Indications**

- An initial diagnostic facet joint injection/medial branch block is considered **medically necessary** to determine whether chronic neck or back pain is of facet joint origin when **ALL** of the following criteria are met:
  - ◆ Pain is exacerbated by facet loading maneuvers on physical examination
  - ◆ Pain has persisted despite at least four weeks of appropriate conservative treatment (e.g., physical methods including physical therapy, chiropractic care and exercise, nonsteroidal anti-inflammatory drugs (NSAIDs), and/or analgesics) unless contraindicated and the reason(s) for contraindication(s) is/are documented in the medical record
  - ◆ Clinical findings and imaging studies suggest no other obvious cause of the pain (e.g., central spinal stenosis with neurogenic claudication/myelopathy, foraminal stenosis or disc herniation with concordant radicular pain/radiculopathy, infection, tumor, fracture, pseudoarthrosis, pain related to spinal instrumentation).
  - ◆ The spinal motion segment is not posteriorly fused.
- A second diagnostic facet joint injection/medial branch block, performed to confirm the validity of the clinical response to the initial facet joint injection, is considered **medically necessary** when **ALL** of the following criteria are met:
  - ◆ Administered at the same level as the initial block
  - ◆ The initial diagnostic facet joint injection produced a positive response (i.e., at least 80% relief of facet mediated pain for at least the expected minimum duration of the effect of the local anesthetic)
  - ◆ A radiofrequency joint denervation/ablation procedure is being considered
- An intra-articular facet joint injection performed with synovial cyst aspiration, in addition to a transforaminal epidural steroid injection, is considered **medically necessary** when the following criteria are met:
  - ◆ Advanced diagnostic imaging studies (e.g., MRI, CT, CT myelogram) confirm compression or displacement of the corresponding nerve root by a facet joint synovial cyst
  - ◆ Clinical correlation with the individual's signs and symptoms of radicular pain or radiculopathy, based on history and physical examination.

### **CMM-201.4: Non-Indications**

- Performance of a facet joint injection/medial branch block is considered **not medically necessary** when performed for **ANY** of the following indications:
  - ◆ Without the use of fluoroscopic or CT guidance
  - ◆ In the presence of an untreated radiculopathy (with the exception of radiculopathy caused by a facet joint synovial cyst)
  - ◆ When a radiofrequency joint denervation/ablation procedure (i.e., facet neurotomy, facet rhizotomy) is not being considered
  - ◆ The facet joint injection is performed at a fused posterior spinal motion segment (with the exception of patients with clinically suspected pseudoarthrosis)
  - ◆ On the same day of service when performing other injections (e.g., epidural steroid, sacroiliac) in the same region
  - ◆ Performance of injections/blocks on more than three (3) contiguous spinal joint levels (with the exception of an intervening fused segment)
  - ◆ Additional diagnostic facet joint injection/medial branch blocks at the same level(s) as a prior successful radiofrequency denervation/ablation procedure
- Performance of a facet joint injection/medial branch block is considered **experimental, investigational, or unproven** when performed for **ANY** of the following indications:
  - ◆ Unless performed as a second confirmatory block, all injections subsequent to the initial injection (i.e., therapeutic injections)
  - ◆ When performed under ultrasound guidance

### **CMM-201.5: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
64490	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), cervical or thoracic, single level
+64491	Injection(s), of diagnostic or therapeutic substance(s) (including anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, includes contrast for localization when performed, epidural or subarachnoid; lumbar or sacral (caudal), second level (List separately)
+64492	Injection(s), of diagnostic or therapeutic substance(s) (including anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, includes contrast for localization when performed, epidural or subarachnoid; lumbar or sacral (caudal), third and any additional level(s) (List separately)
64493	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), lumbar or sacral, single level
+64494	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), lumbar or sacral, second level (List separately)
64495	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with image guidance (fluoroscopy or CT), lumbar or sacral, third and any additional level(s) (List separately)

CPT®	Codes Considered Experimental, Investigational or Unproven
0213T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; single level
0214T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; second level (List separately in addition to code for primary procedure)
0215T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, cervical or thoracic; third and any additional level(s) (List separately in addition to code for primary procedure)
0216T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; single level
0217T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; second level (List separately in addition to code for primary procedure)
0218T	Injection(s), diagnostic or therapeutic agent, paravertebral facet (zygapophyseal) joint (or nerves innervating that joint) with ultrasound guidance, lumbar or sacral; third and any additional level(s) (List separately in addition to code for primary procedure)
This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.	

### CMM-201.6: References

1. Airaksinen O, Brox J, Cedraschi C, et al. On behalf of the COST B13 Working Group on Guidelines for Chronic Low Back Pain. Chapter 4 European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J.* 2006;15(Supplement 2):s192-s300.
2. Allen TL, Tatli Y, Lutz, GE. Fluoroscopic percutaneous lumbar zygoapophyseal joint cysts rupture: a clinical outcome study. *Spine.* 2009 May;9(5): 387-95.
3. American Medical Association. *Current Procedural Terminology: CPT 2008, Professional Edition.* AMA Press, 2007.
4. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline, 2nd Ed.* 2008.
5. Bogduk N. A narrative review of intra-articular corticosteroid injections for low back pain. *Pain Med.* 2005;6(4):287-296.
6. Boswell M, Colson J, Sehgal N, et al. A systematic review of therapeutic facet joint interventions in chronic spinal pain. *Pain Physician.* 2007;10:229-253.
7. Boswell M, Colson J, Spillane W. Therapeutic Facet Joint Interventions in Chronic Spinal Pain: A Systematic Review of Effectiveness and Complications. *Pain Physician.* 2005;8:101-114.
8. Boswell MV, Manchikanti L, Kaye AD, et al. A best-evidence systematic appraisal of the diagnostic accuracy and utility of facet (zygapophysial) joint injections in chronic spinal pain. *Pain Physician* 2015;18:E497-E533.
9. Boswell M, Shah R, Everett C, et al. Interventional techniques in the management of chronic spinal pain: evidence-based practice guidelines. *Pain Physician.* 2005;8(1):1-47.
10. Boswell M, Trescot A, Datta S, et al. American Society of Interventional Pain Physicians. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician.* 2007;10(1):7-111.
11. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg.* 2012;22:200-206.
12. Cohen S, Raja S. Pathogenesis, diagnosis, and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology.* 2007;106:591-614.
13. Dreyfuss P, Dreyer S. NASS. Lumbar zygapophysial (facet) joint injections. *Spine J.* 2003;3(3 Suppl):50S-59S.
14. Dreyfus P, Kaplan M, Dreyer S, ed Lennard T. *Zygapophyseal Joint Injection Techniques in the Spinal Axis. Procedures in Clinical Practice.* Second Edition. Hanley and Belfus Inc. Philadelphia. 2000, page 276.



15. Friedly J, Chan L, Deyo R. Increases in lumbosacral injections in the Medicare population: 1994 to 2001. *Spine*. 2007;32(16):1754-1760.
16. Friedrich K, Nemecek S, Peloschek P, et al. The prevalence of lumbar facet joint edema in patients with low back pain. *Skeletal Radiol*. 2007;36(8):755-760.
17. Fritz J, Niemeyer T, Clasen S, et al. Management of chronic low back pain: rationales, principles, and targets of imaging-guided spinal injections. *Radiographics*. 2007;27(6):1751-1771.
18. Fuchs S, Erbe T, Fischer H, Tibesku C. Intraarticular hyaluronic acid versus glucocorticoid injections for nonradicular pain in the lumbar spine. *J Vasc Interv Radiol*. 2005;16:1493-1498.
19. Fuchs S, Erbe T, Fischer HL, Tibesku CO. Intraarticular hyaluronic acid versus glucocorticoid injections for nonradicular pain in the lumbar spine. *J Vasc Interv Radiol*. 2005;16:1493-1498.
20. Geilhorn A, Katz J, Suri P. Osteoarthritis of the spine; the facet joints. *Nat Rev Rheum* 2013; 9(4): 216 – 224.
21. Hancock M, Maher C, Latimer J, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J*. 2007;16(10):1539-1550.
22. IPM Guidelines. An Update of Comprehensive Evidence-based Guidelines For Interventional Techniques in Chronic Spinal Pain. Part 2: Guidance and Recommendations. *Pain Physician* 2013; 16:S49 – S283.
23. Kirpalani D, Mitra R. Cervical facet joint dysfunction: a review. *Arch Phys Med Rehabil*. 2008;89(4):770-774.
24. Lakemeier S, Lind M, Schultz W, Fuchs-Winkelmann S, Timmesfeld N, Foelsch C, Peterlein CD. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: A randomized, controlled, double-blind trial. *Anesth Analg*. 2013;117:228-235.
25. Laslett M, McDonald B, Aprill C, et al. Clinical predictors of screening lumbar zygapophyseal joint blocks: development of clinical prediction rules. *Spine J*. 2006;6(4):370-379.
26. Laslett M, Oberg B, April C, McDonald B. Zygapophysial joint blocks in chronic low back pain: a test of Revel's model as a screening test. *BMC Musculoskelet Disord*. 2004;16:5:43.
27. Lord SM, Barnsley L, Bogduk N. Percutaneous radiofrequency neurotomy in the treatment of cervical zygapophysial joint pain: a caution. *Neurosurgery*. 1995;36:732-9.
28. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain: Part II: Guidance and recommendations. *Pain Physician*. 2013;16:S49-S283.
29. Manchikanti L, Cash K, Pampati V, Fellows B. Influence of psychological variables on the diagnosis of facet joint involvement in chronic spinal pain. *Pain Physician*. 2008;11(2):145-160.
30. Manchikanti L, Damron K, Cash K, et al. Therapeutic cervical medial branch blocks in managing chronic neck pain: a preliminary report of a randomized, double-blind, controlled trial. *Pain Physician*. 2006;9(4):333-346.
31. Manchikanti L, Kaye AD, Boswell MV, et al. A systematic review and best evidence synthesis of the effectiveness of therapeutic facet joint interventions in managing chronic spinal pain. *Pain Physician*. 2015; 18:E535-E582.
32. Manchikanti L, Manchikanti K, Cash K, et al. Age-related prevalence of facet-joint involvement in chronic neck and low back pain. *Pain Physician*. 2008;11(1):67-75.
33. Manchikanti L, Manchikanti K, Manchukonda R, et al. Evaluation of lumbar facet joint nerve blocks in the management of chronic low back pain: preliminary report of a randomized, double-blind controlled trial. *Pain Physician*. 2007;10(3):425-40.
34. Manchikanti L, Manchukonda R, Pampati V, et al. Prevalence of facet joint pain in chronic low back pain in postsurgical patients by controlled comparative local anesthetic blocks. *Arch Phys Med Rehabil*. 2007;88(4):449-55.
35. Manchikanti L, Singh V, Falco F, et al. Lumbar facet joint nerve blocks in managing chronic facet joint pain: one-year follow-up of a randomized, double-blind controlled trial. *Pain Physician*. 2008;11(2):121-132.
36. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V. Evaluation of lumbar facet joint nerve blocks in managing chronic low back pain: A randomized, double-blind, controlled trial with a 2-year follow-up. *Int J Med Sci*. 2010;7:124-135.



37. Manchikanti L, Singh V, Falco FJE, Cash KA, Fellows B. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: A randomized, double-blind controlled trial. *Pain Physician*. 2010;13:437-450.
38. Manchikanti L, Pampati V, Bakhit C, et al. Effectiveness of lumbar facet joint nerve blocks in chronic low back pain: A randomized clinical trial. *Pain Physician*. 2001;4:101-117.
39. Manchikanti L, Singh V, Falco FJE, Cash KA, Pampati V, Fellows B. The role of thoracic medial branch locks in managing chronic mid and upper back pain: A randomized, double-blind, active control trial with a 2-year follow-up. *Anesthesiol Res Pract*. 2012;2012:585806.
40. Manchukonda R, Manchikanti K, Cash K, et al. Facet joint pain in chronic spinal pain: an evaluation of prevalence and false-positive rate of diagnostic blocks. *J Spinal Disord Tech*. 2007;20(7):539-545.
41. Nordin M, Carragee E, Hurwitz L, et al; Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. Treatment of neck pain: injections and surgical interventions: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine*. 2008;15;33(4 Suppl):S153-S169.
42. North American Spine Society (NASS). NASS Coverage Policy Recommendation. Facet Joint Intervention. Copyright © 2016 North American Spine Society.
43. Park KD, Jee H, Nam HS, Cho SK, Kim HS, Park Y, Lim OK. Effect of medial branch block in chronic facet joint pain for osteoporotic compression fracture: one year retrospective study. *Ann Rehabil Med*. 2013 Apr;37(2):191-201. doi: 10.5535/arm.2013.37.2.191. Epub 2013 Apr30.
44. Park SC, Kim KH. Effect of adding cervical facet joint injections in a multimodal treatment program for long-standing cervical myofascial pain syndrome with referral pain patterns of cervical facet joint syndrome. *J Anesth*. 2012;26:738-745.
45. Patel J, Schneider B, Smith C on behalf of SIS Patient Safety Committee. Intraarticular Corticosteroid Injections and hyperglycemia. 10/4/17.
46. Resnick D, Choudhri T, Dailey A, et al. American Association of Neurological Surgeons/Congress of Neurological Surgeons. Guidelines for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 13: injection therapies, low-back pain, and lumbar fusion. *J Neurosurg Spine*. 2005;2(6):707-715.
47. Ribeiro LH, Furtado RN, Konai MS, et al. Effect of facet joint injection versus systemic steroids in low back pain: A randomized controlled trial. *Spine (Phila Pa 1976)*. 2013;38:1995-2002.
48. Schneider G, Jull G, Smith A, Emery C, Faris P, Cook C, Frizzell B, Salo P. Derivation of a clinical decision guide in the diagnosis of cervical facet joint pain. *Arch Phys Med Rehabil*, 2014; 95(9): 1695-701.
49. Sehgal N, Dunbar E, Shah R, Colson J. Systematic review of diagnostic utility of facet (zygapophysial) joint injections in chronic spinal pain: an update. *Pain Physician*. 2007;10(1):213-228.
50. Shah RD, Cappiello D, Suresh S. Interventional procedures for chronic pain in children and adolescents: a review of the current evidence. *World Institute of Pain*. 2016: 359-369.
51. van Tulder M, Koes B, Seitsalo S, Malmivaara A. Outcome of invasive treatment modalities on back pain and sciatica: an evidence-based review. *Eur Spine J*. 2006; Suppl 1:S82-92.
52. Workloss Data Institute. Official Disability Guidelines. 2015.
53. Yun DH, Kim HS, Yoo SD, Kim DH, Chon JM, Choi SH, Hwang DG, Jung PK. Efficacy of ultrasonography-guided injections in patients with facet syndrome of the low lumbar spine. *Ann Rehabil Med*. 2012;36:66-71.

**CMM-202: Trigger Point Injections**

<b>CMM-202.1: Definitions</b>	<b>25</b>
<b>CMM-202.2: General Guidelines</b>	<b>25</b>
<b>CMM-202.3: Indications</b>	<b>25</b>
<b>CMM-202.4: Non-indications</b>	<b>26</b>
<b>CMM-202.5: Procedure (CPT®) Codes</b>	<b>26</b>
<b>CMM-202.6: References</b>	<b>27</b>

### CMM-202.1: Definitions

- **Trigger point injections** are defined as an injection of a local anesthetic with or without the addition of a corticosteroid into clinically identified myofascial trigger points.
- **Myofascial trigger point** is defined as a discrete, focal, hyperirritable spot found within a taught band of skeletal muscle or its fascia which when provocatively compressed causes local pain or tenderness as well as characteristic referred pain, tenderness and/or autonomic phenomena. Digital palpation, as well as needle insertion into the trigger point, can often lead to a local twitch response. A local twitch response is a transient visible or palpable contraction of the muscle. The presence of characteristic referred pain, tenderness, muscle shortening and/or autonomic phenomena (e.g., vasomotor changes, pilomotor changes, muscle twitches, etc.) is necessary to render the diagnosis of a myofascial trigger point. Tender points within a muscle or its fascia, which do not refer pain, tenderness and/or autonomic phenomena and lack a local twitch response, cannot be considered a myofascial trigger point.

### CMM-202.2: General Guidelines

- Trigger point injections are not without risk, and can expose patients to potential complications.
- The determination of medical necessity for the use of trigger point injections is always made on a case-by-case basis.

### CMM-202.3: Indications

- Trigger point injections are considered **medically necessary** when **BOTH** of the following criteria are met:
  - ◆ A myofascial trigger point has been identified by the presence of **ONE or MORE** of the following on physical examination:
    - Characteristic referred pain
    - Tenderness
    - Muscle shortening
    - Autonomic phenomena (e.g., vasomotor changes, pilomotor changes, muscle twitches, etc.)
  - ◆ Performed using a local anesthetic with or without steroid (e.g., saline or glucose)
- Repeat trigger point injections are considered **medically necessary** when **BOTH** of the following are documented:
  - ◆ At least 50% pain relief with evidence of functional improvement for a minimum of six (6) weeks following the prior injection(s)
  - ◆ Adequate instruction or supervision in self-management strategies (i.e., therapeutic exercise, ergonomic advice, ADL training, etc.)

### **CMM-202.4: Non-indications**

- Trigger point injections are considered **not medically necessary** for any of the following:
  - ◆ When performed with any substance other than local anesthetic with or without steroid (e.g., saline or glucose)
  - ◆ When performed on the same day of service as other treatments in the same region
  - ◆ When requested for any of the following:
    - Acupuncture
    - Electro-Acupuncture
    - Acupoint injections, aka Biopuncture (saline, sugar, herbals, homeopathic substances)
    - Dry needling
    - Image-guided injection over spinal hardware
- Repeat trigger point injections are considered **not medically necessary** for any of the following:
  - ◆ An isolated treatment modality
  - ◆ An interval of less than two (2) months
  - ◆ More than four (4) trigger point injection sessions per body region per year

### **CMM-202.5: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required. Pre- authorization requirements vary by individual payor.

CPT®	Code Description/Definition
20552	Injection(s); single or multiple trigger point(s), 1 or 2 muscle(s)
20553	Injection(s); single or multiple trigger point(s), 3 or more muscle(s)

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the individual payor (health insurance company, etc.) and is based on the member/patient/client/beneficiary's policy or benefit entitlement structure as well as any third party payor guidelines and/or claims processing rules. Providers are strongly urged to contact each payor for individual requirements if they have not already done so.

## CMM-202.6: References

1. Al-Shenqiti A, Oldham J. Test-retest reliability of myofascial trigger point detection in patients with rotator cuff tendonitis. *Clinical Rehabilitation*. 19(5):482-7, 2005 Aug.
2. Alvarez D, Rockwell P. Trigger points: diagnosis and management. *American Family Physician*. 65(4):653-60, 2002 Feb 15.
3. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*, 2nd Ed. 2008. Accessed 10/1/08.
4. Audette J, Wang F, Smith H. Bilateral activation of motor unit potentials with unilateral needle stimulation of active myofascial trigger points. *American Journal of Physical Medicine & Rehabilitation*. 83(5):368-74, quiz 375-7, 389, 2004 May.
5. Bajaj P, Bajaj P, Graven-Nielsen T, et al: Trigger points in patients with lower limb osteoarthritis. *J Musculoskeletal Pain* 2001;9:17–33.
6. Baldry P. Management of myofascial trigger point pain. *Acupunct Med* 2002;20:2–10 25. Chu J: Does EMG (dry needling) reduce myofascial pain symptoms due to cervical nerve root irritation? *Electromyogr Clin Neurophysiol* 1997;37:259–72.
7. Bron C, Wensing M, Franssen JL, Oostendorp R. Treatment of myofascial trigger points in common shoulder disorders by physical therapy: a randomized controlled trial. *BMC Musculoskeletal Disorders*. 8:107, 2007.
8. Carlsson C. Acupuncture mechanisms for clinically relevant long-term effects—reconsideration and a hypothesis. *Acupunct Med* 2002;20:82–99.
9. Casimiro L, Brosseau L, Milne S, et al: Acupuncture and electroacupuncture for the treatment of RA. *Cochrane Database Syst Rev* 2002;(3):CD003788.
10. Chen J, Chung K, Hou C, et al: Inhibitory effect of dry needling on the spontaneous electrical activity recorded from myofascial trigger spots of rabbit skeletal muscle. *Am J Phys Med Rehabil* 2001;80:729–35.
11. Chen Q, Bensamoun S, Basford J, et al . Identification and quantification of myofascial taut bands with magnetic resonance elastography. *Arch Phys Med Rehabil* 2007;88(12): 1658–61.
12. Chen S, Chen J, Wu Y, et al: Myofascial trigger point in intercostal muscles secondary to herpes zoster infection to the intercostal nerve. *Arch Phys Med Rehabil* 1998;79:336–8.
13. Chu J. Twitch-obtaining intramuscular stimulation: observation in the management of radiculopathic chronic low back pain. *J Musculoskeletal Pain* 1999;7:131–46.
14. Cole T, Edgerton V. Musculoskeletal disorders. In: Cole T, Edgerton V, eds. Report of the Task Force on Medical Rehabilitation Research: June 28-29, 1990, Hunt Valley Inn, Hunt Valley, Md. Bethesda: National Institutes of Health, 1990:61-70.
15. Coupe C, Torelli P, Fuglsang-Frederiksen A, et al. Myofascial trigger points are very prevalent in patients with chronic tension-type headache: a double-blinded controlled study. *Clinical Journal of Pain*. 23(1):23-7, 2007 Jan.
16. Cummings T, White A: Needling therapies in the management of myofascial trigger point pain: a systemic review. *Arch Phys Med Rehabil* 2001;82:986–92.
17. Davies C, Davies A. *The Trigger Point Therapy Workbook (2nd Ed.)*. USA: New Harbinger Publication, Inc., 2004;323.
18. Ettlin T, Schuster C, Stoffel R, et al. A distinct pattern of myofascial findings in patients after whiplash injury. *Archives of Physical Medicine & Rehabilitation*. 89(7):1290-3, 2008 Jul.
19. Fernandez-Carnero J, Fernandez-de-Las-Penas C, de la Llave-Rincon AI, et al. Prevalence of and referred pain from myofascial trigger points in the forearm muscles in patients with lateral epicondylalgia. *Clinical Journal of Pain*. 23(4):353-60, 2007 May.
20. Fernandez-de-Las-Penas C, Alonso-Blanco C, Cuadrado ML. Et al. Myofascial trigger points and their relationship to headache clinical parameters in chronic tension-type headache. *Headache*. 46(8):1264-72, 2006 Sep.
21. Fernandez-de-Las-Penas C, Cuadrado M, Arendt-Nielsen L, et al. Association of cross-sectional area of the rectus capitis posterior minor muscle with active trigger points in chronic tension-type headache: a pilot study. *American Journal of Physical Medicine & Rehabilitation*. 87(3):197-203, 2008 Mar.

23. Fernandez de las Penas C, Cuadrado M, Gerwin R, Pareja J. Referred pain from the trochlear region in tension-type headache: a myofascial trigger point from the superior oblique muscle. *Headache*. 45(6):731-7, 2005 Jun.
24. Fernandez-de-Las-Penas C, Cuadrado M, Pareja JA. Myofascial trigger points, neck mobility, and forward head posture in episodic tension-type headache. *Headache*. 47(5):662-72, 2007 May. Fernandez-de-Las-Penas C, Ge H, Arendt-Nielsen L, et al. Referred pain from trapezius muscle trigger points shares similar characteristics with chronic tension type headache. *European Journal of Pain: Ejp*. 11(4):475-82, 2007 May.
25. Fernandez-de-Las-Penas C, Simons D, Cuadrado M, Pareja J. The role of myofascial trigger points in musculoskeletal pain syndromes of the head and neck. *Current Pain & Headache Reports*. 11(5):365-72, 2007 Oct.
26. Ferrante F, Bearn L, Rothrock R, King L. Evidence against trigger point injection technique for the treatment of cervicothoracic myofascial pain with botulinum toxin type A. *Anesthesiology*. 103(2):377-83, 2005 Aug.
27. Fischer A. Injection techniques in the management of local pain. *J Back Musculoskeletal Rehabil* 1996;7:107-17.
28. Fischer A. New approaches in treatment of myofascial pain. *Phys Med Rehabil Clin North Am* 1997;8:153-69.
29. Fischer A. Pressure threshold measurement for diagnosis of myofascial pain and evaluation of treatment results. *Clin J Pain* 1987;2:207-14.
30. Fischer A. Pressure threshold meter: its use for quantification of tender spots. *Arch Phys Med Rehabil* 1986;67:836-8.
31. Friction J, Kroening R, Haley D, Siegert R. Myofascial pain syndrome of the head and neck: a review of clinical characteristics of 164 patients. *Oral Surg Oral Med Oral Pathol* 1985;60:615-23.
32. Ga H, Choi J, Park C, Yoon H. Acupuncture needling versus lidocaine injection of trigger points in myofascial pain syndrome in elderly patients--a randomised trial. *Acupuncture in Medicine*. 25(4):130-6, 2007 Dec.
33. Ga H, Choi J, Park C, Yoon H. Dry needling of trigger points with and without paraspinal needling in myofascial pain syndromes in elderly patients. *Journal of Alternative & Complementary Medicine*. 13(6):617-24, 2007 Jul-Aug.
34. Ga H, Koh H, Choi J, Kim C. Intramuscular and nerve root stimulation vs. lidocaine injection to trigger points in myofascial pain syndrome. *Journal of Rehabilitation Medicine*. 39(5):374-8, 2007 May.
35. Gam A, Warming S, Larsen L, Jet al, Treatment of myofascial trigger-points with ultrasound combined with massage and exercise--a randomised controlled trial. *Pain*. 77(1):73-9, 1998 Jul.
36. Ge H, Zhang Y, Boudreau S, et al. Induction of muscle cramps by nociceptive stimulation of latent myofascial trigger points. *Experimental Brain Research*. 187(4):623-9, 2008 Jun.
37. Gerwin R, Shannon S, Hong C, et al. Interrater reliability in myofascial trigger point examination. *Pain*. 1997 Jan;69(1-2):65-73.
38. Giamberardino M, Tafuri E, Savini A, et al. Contribution of myofascial trigger points to migraine symptoms. *Journal of Pain*. 8(11):869-78, 2007 Nov.
39. Gunn C. Treatment of Chronic Pain. Intramuscular Stimulation for Myofascial Pain of Radiculopathic Origin. London, Churchill Livingstone, 1996.
40. Han S, Harrison P. Myofascial pain syndrome and trigger-point management. *Reg Anesth* 1997;22: 89-101.
41. Hanten W, Olson S, Butts N, Nowicki A. Effectiveness of a home program of ischemic pressure followed by sustained stretch for treatment of myofascial trigger points. *Physical Therapy*. 80(10):997-1003, 2000 Oct. Ho K, Tan K. Botulinum toxin A for myofascial trigger point injection: a qualitative systematic review. *European Journal of Pain: Ejp*. 11(5):519-27, 2007 Jul.
42. Hoheisel U, Mense S, Simons D, et al: Appearance of new receptive fields in rat dorsal horn neurons following noxious stimulation of skeletal muscle: a model for referral of muscle pain? *Neurosci Lett* 1993;153:9-12.
43. Hong C. Algometry in evaluation of trigger points and referred pain. *J Musculoskeletal Pain* 1998;6:47-59.



44. Hong C, Chen J, Chen S, et al: Histological findings of responsive loci in a myofascial trigger spot of rabbit skeletal muscle from where localized twitch responses could be elicited. *Arch Phys Med Rehabil* 1996;77:962.
45. Hong C, Chen Y, Twehous D, et al: Pressure threshold for referred pain by compression on the trigger point and adjacent areas. *J Musculoskeletal Pain* 1996;4:61–79.
46. Hong C. Consideration and recommendation of myofascial trigger point injection. *J Musculoskeletal Pain* 1994;2:29–59.
47. Hong C. Current research on myofascial trigger points—pathophysiological studies. *J Musculoskeletal Pain* 1999;7:121–9.
48. Hong C, Hsueh T. Difference in pain relief after trigger point injections in myofascial pain patients with and without fibromyalgia. *Arch Phys Med Rehabil* 1996;77:1161-6.
49. Hong C, Kuan T, Chen J, Chen S. Referred pain elicited by palpation and by needling of myofascial trigger points: a comparison. *Archives of Physical Medicine & Rehabilitation*. 78(9):957-60, 1997 Sep.
50. Hong C. Lidocaine injection versus dry needling to myofascial trigger point. The importance of the local twitch response. *Am J Phys Med Rehabil* 1994;73:256-63.
51. Hong C. Myofascial trigger points: pathophysiology and correlation with acupuncture points. *Acupunct Med* 2000;18:41–7.
52. Hong C. New trends in myofascial pain syndrome. *Zhonghua Yi Xue Za Zhi (Taipei)* 2002;65:501–12.
53. Hong C. Pathophysiology of myofascial trigger point. *J Formos Med Assoc* 1996;95:93–104.
54. Hong C. Persistence of local twitch response with loss of conduction to and from the spinal cord. *Arch Phys Med Rehabil* 1994;75:12–6.
55. Hong C, Simons D. Response to treatment for pectoralis minor myofascial pain syndrome after whiplash. *J Musculoskeletal Pain* 1993;1:89–131.
56. Hong C, Simons D. Pathophysiologic and electrophysiologic mechanisms of myofascial trigger points. *Archives of Physical Medicine & Rehabilitation*. 79(7):863-72, 1998 Jul.
57. Hong C, Torigoe Y. Electrophysiologic characteristics of localized twitch responses in responsive bands of rabbit skeletal muscle fibers. *J Musculoskeletal Pain* 1994;2:17–43.
58. Hong C, Torigoe Y, Yu J. The localized twitch responses in responsive bands of rabbit skeletal muscle fibers are related to the reflexes at spinal cord level. *J Musculoskeletal Pain* 1995;3:15–33.
59. Hong C. Treatment of myofascial pain syndrome. *Current Pain & Headache Reports*. 10(5):345-9, 2006 Oct.
60. Hopwood M, Abram S. Factors associated with failure of trigger point injections. *Clin J Pain* 1994;10:227-34.
61. Hou C, Tsai L, Cheng K, et al. Immediate effects of various physical therapeutic modalities on cervical myofascial pain and trigger-point sensitivity. *Archives of Physical Medicine & Rehabilitation*. 83(10):1406-14, 2002 Oct.
62. Hsueh T, Cheng P, Kuan T, Hong C. The Immediate Effectiveness of Electrical Nerve Stimulation and Electrical Muscle Stimulation on Myofascial Trigger Points. *American Journal of Physical Medicine & Rehabilitation*. 1997;76(6):471-476.
63. Hsueh T, Yu S, Kuan T, et al: Association of active myofascial trigger points and cervical disc lesion. *J Formos Med Assoc* 1998;97:174–80.
64. Hubbard D, Berkoff G. Myofascial trigger points show spontaneous needle EMG activity. *Spine* 1993;18:1803–7.
65. Imamura S, Fischer A, Imamura M, Teixeira, et al. Pain management using myofascial approach when other treatment failed. *Phys Med Rehabil Clin North Am* 1997;8:179-96.
66. Kuan T, Chang Y, Hong C. Distribution of active loci in rat skeletal muscle. *J Musculoskeletal Pain* 1999;7:45–54.
67. Kuan T, Chen J, Chen S, et al. Effect of botulinum toxin on endplate noise in myofascial trigger spots of rabbit skeletal muscle. *American Journal of Physical Medicine & Rehabilitation*. 81(7):512-20; quiz 521-3, 2002 Jul.
68. Lavelle E, Lavelle W, Smith H. Myofascial trigger points. *Anesthesiology Clinics*. 25(4):841-51, vii-iii, 2007 Dec.



69. Lee S, Chen C, Lee C, et al. Effects of needle electrical intramuscular stimulation on shoulder and cervical myofascial pain syndrome and microcirculation. *Journal of the Chinese Medical Association: JCMA*. 71(4):200-6, 2008 Apr.
70. Lewis J, Tehan P. A blinded pilot study investigating the use of diagnostic ultrasound for detecting active myofascial trigger points. *Pain*. 79(1):39-44, 1999 Jan.
71. Lewit K. The needle effect in relief of myofascial pain. *Pain* 1979;6:83-90.
72. Ling F, Slocumb J. Use of trigger point injections in chronic pelvic pain. *Obstet Gynecol Clin North Am* 1993;20:809-15.
73. Majlesi J, Unalan H. High-power pain threshold ultrasound technique in the treatment of active myofascial trigger points: a randomized, double-blind, case-control study. *Archives of Physical Medicine & Rehabilitation*. 85(5):833-6, 2004 May.
74. Melzack R. Myofascial trigger points: relation to acupuncture and mechanism of pain. *Arch Phys Med Rehabil* 1981;62:114-7.
75. Mense S, Schmit R. Muscle pain: which receptors are responsible for the transmission of noxious stimuli? In: Rose F, ed. *Physiological aspects of clinical neurology*. Oxford: Blackwell Scientific Publications, 1977:265-78.
76. Mense S. Considerations concerning the neurobiological basis of muscle pain. *Can J Physiol Pharmacol* 1991;69:610-6.
77. Mense S. Nociception from skeletal muscle in relation to clinical muscle pain. *Pain* 1993;54:241-89.
78. Mense S. Peripheral mechanisms of muscle nociception and local muscle pain. *J Musculoskeletal Pain* 1993;1:133-70.
79. Mense S. Referral of muscle pain: new aspects. *Am Pain Soc J* 1994;3:1-9.
80. Meyer H. Myofascial pain syndrome and its suggested role in the pathogenesis and treatment of fibromyalgia syndrome. *Current Pain & Headache Reports*. 6(4):274-83, 2002 Aug.
81. Myburgh C, Larsen AH, Hartvigsen J. A systematic, critical review of manual palpation for identifying myofascial trigger points: evidence and clinical significance. *Archives of Physical Medicine & Rehabilitation*. 89(6):1169-76, 2008 Jun.
82. Ohrbach R, Gale E. Pressure pain thresholds in normal muscles: reliability, measurement effects and topographic differences. *Pain* 1989;37:257-63.
83. Ohrbach R, Gale E. Pressure pain thresholds, clinical assessment, and differential diagnosis: reliability and validity in patients with myofascial pain. *Pain* 1989;39:157-69.
84. Rachlin E. History and physical examination for regional myofascial pain syndrome. In: Rachlin ES, ed. *Myofascial pain and fibromyalgia: trigger point management*. St. Louis: Mosby, 1994:159-72.
85. Rachlin E. Trigger points. In: Rachlin E, ed. *Myofascial pain and fibromyalgia: trigger point management*. St. Louis: Mosby, 1994:145-57.
86. Reeves J, Jaeger B, Graff-Radford S. Reliability of the pressure algometer as a measure of myofascial trigger point sensitivity. *Pain* 1986;24:313-21.
87. Ruoff G. Technique of trigger point injection. In: Pfenninger J, Fowler G, eds. *Procedures for primary care physicians*. St. Louis: Mosby, 1994:164-7.
88. Sciotti V, Mittak V, DiMarco L, Fet al. Clinical precision of myofascial trigger point location in the trapezius muscle. *Pain*. 93(3):259-66, 2001 Sep.
89. Shah J, Danoff J, Desai M, et al. Biochemicals associated with pain and inflammation are elevated in sites near to and remote from active myofascial trigger points. *Archives of Physical Medicine & Rehabilitation*. 89(1):16-23, 2008 Jan.
90. Simons D, Hong C, Simons L. Endplate potentials are common to midfiber myofascial trigger points. *Am J Phys Med Rehabil* 2002;81:212-22.
91. Simons D, Hong C, Simons L. Prevalence of spontaneous electrical activity at trigger spots and at control sites in rabbit skeletal muscle. *J Musculoskeletal Pain* 1995;3:35-48.
92. Simons D. Cardiology and Myofascial Trigger Points: Janet G. Travell's Contribution. *Tex Heart Inst J*. 2003; 30(1): 3-7.
93. Simons D. Clinical and etiological update of myofascial pain from trigger points. *J Musculoskeletal Pain* 1996;4:93-121.
94. Simons D. New views of myofascial trigger points: etiology and diagnosis. *Arch Phys Med Rehabil* 2008;89(1): 157-9.

95. Simons D, Travell J, Simons L. Travell & Simons' *Myofascial pain and dysfunction: the trigger point manual*. 2d ed. Baltimore: Williams & Wilkins, 1999.
96. Smith-Fassler M, Lopez-Bushnell K: Acupuncture as complementary therapy for back pain. *Holist Nurs Pract* 2001;15:35–44.
97. Sola A, Bonica J. Myofascial pain syndromes. In: Bonica JJ, ed. *The management of pain*. 2d ed. Philadelphia: Lea & Febiger, 1990:352-67.
98. Srbely J, Dickey J. Randomized controlled study of the antinociceptive effect of ultrasound on trigger point sensitivity: novel applications in myofascial therapy?. *Clinical Rehabilitation*. 21(5):411-7, 2007 May.
99. Trinh K, Phillips S, Ho E, et al: Acupuncture for the alleviation of lateral epicondyle pain: a systematic review. *Rheumatology (Oxford)* 2004;43:1085–90.
100. Workloss Data Institute. Official Disability Guidelines.
101. Yunus M. Fibromyalgia syndrome and myofascial pain syndrome: clinical features, laboratory tests, diagnosis, and pathophysiologic mechanisms. In: Rachlin E, ed. *Myofascial pain and fibromyalgia: trigger point management*. St. Louis: Mosby, 1994:3-29.
102. Zaralidou A, Amaniti E, Maidatsi P, et al. Comparison between newer local anesthetics for myofascial pain syndrome management. *Methods & Findings in Experimental & Clinical Pharmacology*. 29(5):353-7, 2007 Jun.
103. Zohn D, Mennell J. Musculoskeletal pain: diagnosis and physical treatment. Boston: Little, Brown, 1976:126-9,190-3.

**CMM-203: Sacroiliac Joint Injections**

<b>CMM-203.1: Definitions</b>	<b>33</b>
<b>CMM-203.2: General Guidelines</b>	<b>33</b>
<b>CMM-203.3: Indications</b>	<b>34</b>
<b>CMM-203.4: Non-Indications</b>	<b>35</b>
<b>CMM-203.5: Procedure Codes</b>	<b>35</b>
<b>CMM-203.6: References</b>	<b>36</b>

### **CMM-203.1: Definitions**

- The presence of pain over the sacroiliac joint in the absence of radicular findings in and of itself does not substantiate the diagnosis of sacroiliac joint pain. There must also be clinical evidence as described below.
- **Intra-articular sacroiliac joint injection** refers to the injection of contrast (absent allergy to contrast), followed by the introduction of a corticosteroid and/or a local anesthetic into the sacroiliac joint under fluoroscopic guidance.
- **Peri-articular injection** refers to the introduction of a corticosteroid and/or a local anesthetic to one or more sections of the posterior ligamentous structures of the sacroiliac joint.
- **Sacral lateral nerve block** refers to an injection of corticosteroid and/or local anesthetic adjacent to the sacral lateral nerve resulting in the temporary interruption of conduction of impulses for analgesia. Sacral lateral nerve blocks attempt to block pain signals and theoretically provide relief from pain. The duration of the block depends on the dose, concentration and type of pharmacological agent injected.
- **Sacroiliac joint pain** is defined as pain originating from the sacroiliac joint and/or its supporting ligamentous structures as a result of injury, disease or surgery.

### **CMM-203.2: General Guidelines**

- The determination of medical necessity for the performance of sacroiliac joint injections is always made on a case-by-case basis.
- Intra-articular sacroiliac joint injections should be performed using fluoroscopy with injection of contrast (absent allergy to contrast) for guidance, as it is considered the standard of care.
- Peri-articular sacroiliac joint injections may be performed with or without the use of fluoroscopic guidance.
- When sacroiliac joint injections are performed (anesthetic only) for the purpose of diagnosing sacroiliac joint pain, a positive diagnostic response is defined as  $\geq 75\%$  pain relief for the duration of the local anesthetic.
- Sacroiliac injections performed for the purpose of treating sacroiliac pain are termed therapeutic sacroiliac injections. When medical necessity criteria is met, a total of four therapeutic sacroiliac injections for the treatment of sacroiliac pain may be performed per joint during a 12 month period of time, with a minimum of two months duration between each injection, for the recurrence of pain.
- The performance of interventional pain procedures such as a sacroiliac joint injection does not require the need for supplemental anesthesia in addition to local anesthesia.

### **CMM-203.3: Indications**

- The performance of a diagnostic sacroiliac joint injection for localized, sacroiliac joint pain resulting from disease, injury or surgery is considered **medically necessary** when **ALL** of the following criteria are met:
  - ◆ Pain primarily experienced between the upper level of the iliac crests and the gluteal fold (the pain can refer distally, even below the knee)
  - ◆ Clinical findings and imaging studies suggest no other obvious cause of the pain (e.g., central spinal stenosis with neurogenic claudication/myelopathy, foraminal stenosis or disc herniation with concordant radicular pain/radiculopathy, infection, tumor, fracture, pseudoarthrosis, pain related to spinal instrumentation).
  - ◆ Reproduction of pain using at least three (3) of the following provocative tests:
    - Distraction or “Gapping” or FABER/Patrick’s Test
    - Thigh Thrust or Posterior Pelvic Pain Provocational Test
    - Gaenslan’s Test
    - Sacroiliac Joint Compression Test
    - Sacral Thrust or Yeoman’s Test.
- Pain persists despite **BOTH** of the following:
  - ◆ A minimum of four (4) weeks of noninvasive conservative therapy (e.g., exercise, physical therapy, chiropractic care, nonsteroidal anti-inflammatory drugs [NSAIDs] and analgesics)
  - ◆ Ongoing, active participation in rehabilitative/therapeutic exercise program
- A therapeutic sacroiliac joint injection for the treatment of sacroiliac joint pain is considered **medically necessary** following a diagnostic injection with  $\geq 75\%$  reduction in the reported pain.
- A repeat therapeutic sacroiliac joint injection for the treatment of sacroiliac joint pain is considered **medically necessary** following a therapeutic injection with  $\geq 75\%$  reduction in the reported pain and **BOTH** of the following are met:
  - ◆ **EITHER** of the following:
    - Increase in the individual’s level of function (i.e., return to work)
    - Reduction in the use of pain medication and/or additional medical services such as physical therapy/chiropractic care
  - ◆ A minimum of two months since the prior injection

### **CMM-203.4: Non-Indications**

- Ultrasound guidance for a sacroiliac joint injection, for any indication, is considered **experimental, investigational, or unproven**.
- Sacral lateral nerve branch blocks and/or ablations/neurotomies for the diagnosis and/or treatment of sacroiliac joint mediated pain is considered **experimental, investigational, or unproven**.
- A sacroiliac joint injection is considered **not medically necessary** for **ANY** of the following:
  - ◆ Sacroiliac joint injections performed without fluoroscopic or other alternative guidance, with the exception of ultrasound as noted above
  - ◆ When performed on the same of service as a facet joint block, epidural steroid injection, or lumbar sympathetic chain block
  - ◆ When performed in isolation (i.e., without the individual participating in an active rehabilitation program, home exercise program, or functional restoration program)
  - ◆ As a subsequent diagnostic block when the initial diagnostic block does not produce a positive response of  $\geq 75\%$  pain reduction
  - ◆ Therapeutic sacroiliac joint injections performed at a frequency greater than once every two (2) months for the treatment of sacroiliac pain
  - ◆ More than four (4) injections per SI joint performed within a 12 month period

### **CMM-203.5: Procedure Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

<b>CPT®</b>	<b>Code Description/Definition</b>
<b>27096</b>	Injection procedure for sacroiliac joint, anesthetic/steroid, with image guidance (fluoroscopy or CT) including arthrography when performed
<b>G0259</b>	Injection procedure for sacroiliac joint; arthrography
<b>G0260</b>	Injection procedure for sacroiliac joint; provision of anesthetic, steroid and/or other therapeutic agent, with or without arthrography

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.



## CMM-203.6: References

1. American College of Occupational and Environmental Medicine. Occupational Medicine Practice Guideline, 2nd Ed. 2008.
2. American Medical Association. *Current Procedural Terminology*. 2016 Professional Edition.
3. American Society of Anesthesiologists. Statement on Anesthetic Care During Interventional Pain Procedures for Adults. October 22, 2005, amended October 26, 2016.
4. Appropriate Use Criteria for Fluoroscopically-Guided Diagnostic and Therapeutic Sacroiliac Interventions: Results from the Spine Intervention Society-Convended Multispecialty Collaborative
5. Berthelot J, Labat J, Le Goff B, et al. Provocative sacroiliac joint maneuvers and sacroiliac joint block are unreliable for diagnosing sacroiliac joint pain. *Joint Bone Spine*. 2006;73(1):17-23.
6. Cheng J, Pope JE, Dalton JE, Cheng O, Bensitel A. Comparative outcomes of cooled versus traditional radiofrequency ablation of the lateral branches for sacroiliac joint pain. *Clin J Pain*.2013; 29:132-137.
7. Cohen SP, Hurley RW, Buckenmaier CC 3rd, et al. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology*.2008; 109:279-288.
8. Foley B, Buschbacher R: Sacroiliac joint pain: anatomy, biomechanics, diagnosis and treatment. *Am J Phys Med Rehabil*. 2006;85:997–1006.
9. Forst S, Wheeler M, Fortin J, Vilensky J. The sacroiliac joint: anatomy, physiology and clinical significance. *Pain Physician*. 2006;9(1):61-67.
10. Gunaydin I, Pereira P, Fritz J, et al. Magnetic resonance imaging guided corticosteroid injection of sacroiliac joints in patients with spondylarthropathy. Are multiple injections more beneficial?. *Rheumatology International*. 2006; 26(5):396-400.
11. Karabacakoglu A, Karakose S, Ozerbil O, Odev K. Fluoroscopy-guided intraarticular corticosteroid injection into the sacroiliac joints in patients with ankylosing spondylitis. *Acta Radiologica*. 2002;43(4):425-427.
12. Kennedy DJ, Engel AJ, Kreiner DS, Nampiaparampil D, Duszynski B, MacVicar J. Fluoroscopically guided diagnostic and therapeutic sacroiliac joint injections: a systematic review. *Pain Med* 2015; 16: 1500-1518.
13. King W, Ahmed SU, Baisden J, Patel N, Kennedy DJ, MacVicar J, Duszynski B. Diagnosis and treatment of posterior sacroiliac complex pain: a systematic review with comprehensive analysis of the published data. *Pain Med* 2015 Feb; 16(2): 257.
14. Laslett M. Evidence-Based Diagnosis and Treatment of the Painful Sacroiliac Joint. *J Man Manip Ther*. 2008; 16(3): 142–152.
15. Ling B, Lee J, Man H, et al. Transverse morphology of the sacroiliac joint: effect of angulation and implications for fluoroscopically guided sacroiliac joint injection. *Skeletal Radiology*. 2006;35(11):838-846.
16. Luukkainen R, Nissila M, Asikainen E, Set al. Periarticular corticosteroid treatment of the sacroiliac joint in patients with seronegative spondylarthropathy. *Clinical & Experimental Rheumatology*. 1999;17(1):88-90.
17. Luukkainen R, Wennerstrand P, Kautiainen H, et al. Efficacy of periarticular corticosteroid treatment of the sacroiliac joint in non-spondylarthropathic patients with chronic low back pain in the region of the sacroiliac joint. *Clinical & Experimental Rheumatology*. 2002;20(1):52-54.
18. McKenzie-Brown A, Shah R, Sehgal N, Everett C. A Systematic Review of Sacroiliac Joint Interventions. *Pain Physician*. 2005;8;115-125.
19. MacVicar J, Kreiner DS, Duszynski B, et. al. Appropriate Use Criteria for Fluoroscopically Guided Diagnostic and Therapeutic Sacroiliac Interventions: Results from the Spine Intervention Society Convended Multispecialty Collaborative, *Pain Medicine*, Volume 18, Issue 11, 1 November 2017, Pages 2081–2095
20. Manchikanti L, Staats P, Singh V, et al. Evidence-based practice guidelines for interventional techniques in the management of chronic spinal pain. *Pain Physician*. 2003;6:3-81.
21. Mitchell B, MacPhail T, Vivian D, Verrills P, Barnard A. Radiofrequency neurotomy for sacroiliac joint pain: A prospective study. *Surgical Science*. 2015;6:265-272.
22. Murakami E, Tanaka Y, Aizawa T, et al. Effect of periarticular and intraarticular lidocaine injections for sacroiliac joint pain: prospective comparative study. *Journal of Orthopaedic Science*. 2007;12(3):274-280.
23. Nelemans P, de Bie R, de Vet H, Sturmans F, Injection therapy for subacute and chronic benign low back pain. *Cochrane Database Syst Rev*. 2000;(2):CD001824.
24. Patel J, Schneider B, Smith C on behalf of SIS Patient Safety Committee. Intrarticular Corticosteroid Injections and hyperglycemia. 10/4/17.
25. Patel N, Gross A, Brown L, Gekht G. A randomized, placebo-controlled study to assess the efficacy of lateral branch neurotomy for chronic sacroiliac joint pain. *Pain Med*.2012; 13:383-398.
26. Pekkafehli M, Kiralp M, Basekim C, et al. Sacroiliac joint injections performed with sonographic guidance. *Journal of Ultrasound in Medicine*. 2003;22(6):553-559.
27. Rosenberg J, Quint T, de Rosayro A. Computerized tomographic localization of clinically-guided sacroiliac joint injections. *Clinical Journal of Pain*. 2000;16(1):18-21.



28. Schmidt GL, Bhandutia AK, Altman DT. Management of sacroiliac joint pain. *J Am Acad Orthop Surg*. 2018;26:610-616. doi: 10.5435/JAAOS-D-15-00063.
29. Slipman C, Lipetz J, Plastaras C, et al. Fluoroscopically guided therapeutic sacroiliac joint injections for sacroiliac joint syndrome. *American Journal of Physical Medicine & Rehabilitation*. 2001;80(6):425-432.
30. Simopoulos TT, Manchikanti L, Gupta S, et al. Systematic review of the diagnostic accuracy and therapeutic effectiveness of sacroiliac joint interventions. *Pain Physician*. 2015;18:E713-E756.
31. Stelzer W, Aiglesberger M, Stelzer D, Stelzer V. Use of cooled radiofrequency lateral branch neurotomy for the treatment of sacroiliac joint-mediated low back pain: A large case series. *Pain Med*. 2013;14:29-35.
32. Vallejo R, Benyamin R, Kramer J, et al. Pulsed radiofrequency denervation for the treatment of sacroiliac joint syndrome. *Pain Med*. 2006;7:429-434.
33. van der Wurff P, Buijs E, Groen G. A multitest regimen of pain provocation tests as an aid to reduce unnecessary minimally invasive sacroiliac joint procedures. *Archives of Physical Medicine & Rehabilitation*. 2006; 87(1):10-14.
34. Workloss Data Institute. Official Disability Guidelines. [www.worklossdata.com](http://www.worklossdata.com).
35. Young S, Aprill C, Laslett M. Correlation of clinical examination characteristics with three sources of chronic low back pain. *Spine Journal: Official Journal of the North American Spine Society*. 2003;3(6):460-465.
36. Zelle B, Gruen G, Brown S, George S. Sacroiliac joint dysfunction: evaluation and management. *Clin J Pain*. 2005;21(5):446-455.

**CMM-204: Prolotherapy**

<b>CMM-204.1: Definition</b>	<b>39</b>
<b>CMM-204.2: General Guidelines</b>	<b>39</b>
<b>CMM-204.3: Procedure (CPT®) Codes</b>	<b>39</b>
<b>CMM-204.4: References</b>	<b>40</b>

### **CMM-204.1: Definition**

**Prolotherapy** is defined as an injection or a series of injections designed to strengthen weak or lax ligaments, tendons or joints by injecting various proliferating agents (sclerosing solutions) directly into the proposed damaged or stretched ligaments or tendons or into a joint or its adjacent structures to create scar tissue in an effort to stabilize the joint or tendon. Agents used with prolotherapy have included zinc sulfate, psyllium seed oil, combinations of dextrose, glycerin and phenol, or dextrose alone.

### **CMM-204.2: General Guidelines**

Based on the lack of conclusive scientific evidence demonstrating the clinical efficacy of prolotherapy combined with the potential to expose individuals to adverse side effects or complications, the use of prolotherapy in the treatment of musculoskeletal pain and/or instability (e.g., laxity, weakness) is considered **experimental, investigational or unproven**.

### **CMM-204.3: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Codes Considered Experimental, Investigational , or Unproven
<b>M0076</b>	Prolotherapy

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.

## CMM-204.4: References

1. Alderman, D Prolotherapy for Low Back Pain *Practical Pain Management* 2007;7(4):58-63.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*, 2nd Ed. 2008.
3. Blue Cross Blue Shield. Medicine Section - Prolotherapy. Policy No: 40. Effective Date: 07/11/06.
4. Dagenais S, Haldeman S, Wooley J. Intraligamentous injection of sclerosing solutions (prolotherapy) for spinal pain: a critical review of the literature. *Spine J*. 2005;5(3):310-28.
5. Dagenais S, Ogunseitani O, Haldeman S, et al. Side effects and adverse events related to intraligamentous injection of sclerosing solutions (prolotherapy) for back and neck pain: A survey of practitioners. *Arch Phys Med Rehabil*. 2006;87(7):909-913.
6. Dagenais S, Yelland M, Del Mar C, Schoene M. Prolotherapy injections for chronic low-back pain. *Cochrane Database of Systematic Reviews*, 2007, Issue 2.
7. Dechow E, Davies R, Carr A, Thompson P. A randomized, double-blind, placebo-controlled trial of sclerosing injections in patients with chronic low back pain. *Rheumatology (Oxford)*. 1999; 38(12):1255-1259.
8. Fullerton B. High-resolution ultrasound and magnetic resonance imaging to document tissue repair after prolotherapy: a report of 3 cases. *Archives of Physical Medicine & Rehabilitation*. 2008; 89(2):377-385.
9. Hooper R, Ding M. Retrospective case series on patients with chronic spinal pain treated with dextrose prolotherapy. *J Altern Complement Med*. 2004;10(4):670-674.
10. Kim S, Stitik T, Foye P, et al, Critical Review of Prolotherapy for Osteoarthritis, Low Back Pain, and other Musculoskeletal Conditions: A Physiatric Perspective, *Am J Phys Med Rehabil*. 2004;83(5):379-89.
11. Manchikanti L, Damron K, Cash K, et al. Therapeutic cervical medial branch blocks in managing chronic neck pain: A preliminary report of a randomized, double-blind, controlled trial: Clinical trial NCT0033272. *Pain Physician*. 2006;9(4):333-346.
12. Manchikanti L, Manchikanti K, Manchukonda R, et al. Evaluation of lumbar facet joint nerve blocks in the management of chronic low back pain: Preliminary report of a randomized, double-blind controlled trial: clinical trial NCT00355914. *Pain Physician*. 2007;10(3):425-440.
13. Manchikanti L, Pampati V, Fellows B, et al. The diagnostic validity and therapeutic value of lumbar facet joint nerve blocks with or without adjuvant agents. *Curr Rev Pain*. 2000;4(5)337-344.
14. Nelemans P, de Bie R, de Vet H, Sturmans F. Injection therapy for subacute and chronic benign low back pain. *Cochrane Database of Systematic Reviews*. 2003, Issue 4.
15. Ohberg L, Alfredson H. Sclerosing therapy in chronic Achilles tendon insertional pain-results of a pilot study.
16. *Knee Surgery, Sports Traumatology, Arthroscopy*. 2003;11(5):339-343.
17. Rabago D, Best T, Beamsley M, Patterson J. A systematic review of prolotherapy for chronic musculoskeletal pain. *Clin J Sport Med*. 2005;15(5):376-380.
18. Reeves K, Hassanein K. Randomized prospective double-blind placebo-controlled study of dextrose prolotherapy for knee osteoarthritis with or without ACL laxity. *Altern Ther Health Med*. 2000;6(2):68-80.
19. Reeves K, Hassanein K. Randomized, prospective, placebo-controlled double-blind study of dextrose prolotherapy for osteoarthritic thumb and finger (DIP, PIP, and trapeziometacarpal) joints: evidence of clinical efficacy. *J Altern Complement Med*. 2000;6(4):311-320.
20. Reisner L. Biologic poisons for pain. *Curr Pain Headache Rep*. 2004;8(6):427-434.
21. Workloss Data Institute. Official Disability Guidelines.
22. Yelland M, Del Mar C, Pirozzo S, Schoene ML. Prolotherapy injections for chronic low back pain: a systematic review. *Spine*. 2004; 29(19):2126-2133.
23. Yelland M, Glasziou P, Bogduk N, et al. Prolotherapy injections, saline injections, and exercises for chronic low-back pain: a randomized trial. *Spine*. 2004;29(1):9-16.

**CMM-207: Epidural Adhesiolysis**

<b>CMM-207.1: Definition</b>	<b>42</b>
<b>CMM-207.2: General Guidelines</b>	<b>42</b>
<b>CMM-207.3: Procedure (CPT®) Codes</b>	<b>42</b>
<b>CMM-207.4: References</b>	<b>43</b>

### **CMM-207.1: Definition**

**Epidural adhesiolysis** is also known as epidural neurolysis, epidural decompressive neuroplasty, and Racz neurolysis. It is defined as a treatment for back pain that involves disruption, reduction, and/or elimination of fibrous tissue from the epidural space, which is carried out by either catheter manipulation or the injection of saline or other adhesiolytic agents. A catheter is used to enter the epidural space through a caudal, interlaminar, or transforaminal approach. The goal is to free the nerve root of adhesions and allow introduction of medications to the affected nerve root. An anesthetic along with a glucocorticosteroid may also be injected as part of the procedure. These procedures may also involve spinal endoscopy to visually address the adhesions.

### **CMM-207.2: General Guidelines**

There is insufficient scientific evidence to support the use of epidural adhesiolysis, performed by catheter or endoscopically, as a treatment for back pain. It is considered **experimental, investigational or unproven**.

### **CMM-207.3: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code’s inclusion on this list does not necessarily indicate prior authorization is required.	
<b>CPT®</b>	<b>Codes Considered Experimental, Investigational or Unproven</b>
<b>62263</b>	Percutaneous lysis of epidural adhesions using solution injection (eg, hypertonic saline, enzyme) or mechanical means (e.g., catheter) including radiologic localization (includes contrast when administered), multiple adhesiolysis sessions; 2 or more days
<b>62264</b>	Percutaneous lysis of epidural adhesions using solution injection (e.g., hypertonic saline, enzyme) or mechanical means (e.g., catheter) including radiologic localization (includes contrast when administered), multiple adhesiolysis sessions; 1 day
<b>62280</b>	Injection/infusion of neurolytic substance (e.g., alcohol, phenol, iced saline solutions), with or without other therapeutic substance; subarachnoid
<b>62281</b>	Injection/infusion of neurolytic substance (e.g., alcohol, phenol, iced saline solutions), with or without other therapeutic substance; epidural, cervical or thoracic
<b>62282</b>	Injection/infusion of neurolytic substance (e.g., alcohol, phenol, iced saline solutions), with or without other therapeutic substance; epidural, lumbar, sacral (caudal)
This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual’s policy or benefit entitlement structure as well as claims processing rules.	



## **CMM-207.4: References**

1. American College of Occupational and Environmental Medicine (ACOE). *Occupational Medicine Practice Guideline*. 2nd Ed. 2008.
2. American Medical Association (AMA). *Current Procedural Terminology – 2014 Professional Edition*.
3. Belozer M, Wang G. Epidural Adhesiolysis for the Treatment of Back Pain, Health Technology Assessment. *Washington State Department of Labor and Industries*. July 13, 2004.
4. Boswell M, Shah R, Everett C, et al. Interventional Techniques: Evidence-based Practice Guidelines in the Management of Chronic Spinal Pain. *Pain Physician*. 2007;10:7-111.
5. Boswell M, Shah R, Everett C, et al. Interventional techniques in the management of chronic spinal pain: evidence-based practice guidelines. *Pain Physician*. 2005;8(1):1-47.
6. Cahana A, Mavrocordatos P, Geurts J, Groen G. Do minimally invasive procedures have a place in the treatment of chronic low back pain? *Expert Rev Neurother*. 2004;4(3):479-90.
7. Choi E, Nahm FS, Lee PB. Evaluation of prognostic predictors of percutaneous adhesiolysis using a Racz catheter for post lumbar surgery syndrome or spinal stenosis. *Pain Physician*. 2013; 16:E531-E536.
8. Chopra P, Smith H, Deer T, Bowman R. Role of Adhesiolysis in the Management of Chronic Spinal Pain: A Systematic Review of Effectiveness and Complications. *Pain Physician*. 2005;8(1):87- 100.
9. Hammer M, Doleys D, Chung O. Transforaminal ventral epidural adhesiolysis. *Pain Physician*. 2001;4:273-279.
10. Heavner J, Racz G, Raj P. Percutaneous epidural neuroplasty: prospective evaluation of 0.9% NaCl versus 10% NaCl with or without hyaluronidase. *Reg Anesth Pain Medicine*. 1999;24(3):202-207.
11. Gerdesmeyer L, Wagenpfeil S, Birkenmaier C, et al. Percutaneous epidural lysis of adhesions in chronic lumbar radicular pain: A randomized double-blind placebo controlled trial. *Pain Physician*. 2013; 16:185-196.
12. Helm II S, Benyamin RM, Chopra P, Deer TR, Justiz R. Percutaneous adhesiolysis in the management of chronic low back pain in post lumbar surgery syndrome and spinal stenosis: A systematic review. *Pain Physician*. 2012; 15:E435-E462.
13. Helm II S, Benyamin RM, Falco FJE. Refinement in evidence synthesis of percutaneous adhesiolysis. *Pain Physician*. 2013; 16:177-184.
14. Helm II S, Hayek SM, Colson J, et al. Spinal endoscopic adhesiolysis in post lumbar surgery syndrome: An update of the assessment of the evidence. *Pain Physician*. 2013; 16:SE125-SE150.
15. Igarashi T, Hirabayashi Y, Seo N, et al. Lysis of adhesions and epidural injection of steroid/local anesthetic during epiduroscopy potentially alleviate low back and leg pain in elderly patients with lumbar spine stenosis. *Br J Anaesth*. 2004;93:181-187.
16. Kim SH, Choi WJ, Suh JH, et al. Effects of transforaminal balloon treatment in patients with lumbar foraminal stenosis: A randomized, controlled, double-blind trial. *Pain Physician*. 2013; 16:213-224.
17. Koh WU, Choi SS, Park WY, et al. Transforaminal hypertonic saline for the treatment of lumbar lateral canal stenosis: A double-blinded, randomized, active-control trial. *Pain Physician*. 2013; 197-211.
18. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain: Part II: Guidance and recommendations. *Pain Physician*. 2013;16:S49-S283.
19. Manchikanti L, Bakhit C. Percutaneous lysis of epidural adhesions. *Pain Physician*. 2000;3(1):46-64.
20. Manchikanti L, Boswell M, Rivera J, et al. A randomized, controlled trial of spinal endoscopic adhesiolysis in chronic refractory low back and lower extremity pain. *BMC Anesthesiol*. 2005;5:10.
21. Manchikanti L, Cash KA, McManus CD, Pampati V. Assessment of effectiveness of percutaneous adhesiolysis in managing chronic low back pain secondary to lumbar central spinal canal stenosis. *Int J Med Sci*. 2013; 10:50-59.
22. Manchikanti L, Manchikanti KN, Gharibo CG, Kaye AD. Efficacy of percutaneous adhesiolysis in the treatment of lumbar post surgery syndrome. *Anesth Pain Med*. 2015; in press.

23. Manchikanti L, Pampati V, Fellows B, et al. Role of one day epidural adhesiolysis in management of chronic low back pain: A randomized clinical trial. *Pain Physician*. 2001;4:153-166.
24. Manchikanti L, Pampati V, Rivera J, et al. Effectiveness of percutaneous adhesiolysis and hypertonic saline neurolysis in refractory spinal stenosis. *Pain Physician*. 2001;4:366-373.
25. Manchikanti L, Rivera J, Pampati Vet al. One day lumbar epidural adhesiolysis and hypertonic saline neurolysis in treatment of chronic low back pain: A randomized, double-blind trial. *Pain Physician*. 2004;7:177-186.
26. Manchikanti L, Singh V, Cash KA, Pampati V, Datta S. Assessment of effectiveness of percutaneous adhesiolysis and caudal epidural injections in managing lumbar post surgery syndrome: A 2-year follow-up of randomized, controlled trial. *J Pain Res*. 2012; 5:597-608.
27. Trescot A, Chopra P, Abdi S, et al. Systematic review of effectiveness and complications of adhesiolysis in the management of chronic spinal pain: an update. *Pain Physician*. 2007;10(1):129- 146.
28. Veihelmann A, Devens C, Trouillier H, et al. Epidural neuroplasty versus physiotherapy to relieve pain in patients with sciatica: a prospective randomized blinded clinical trial. *J Orthop Sci*. 2006;11(4):365-369.
29. Workloss Data Institute. *Official Disability Guidelines*. 2008.

<b>CMM-208: Radiofrequency Joint Ablations/Denervations</b>	
<b>CMM-208.1: Definitions</b>	<b>46</b>
<b>CMM-208.2: General Guidelines</b>	<b>46</b>
<b>CMM-208.3: Indications</b>	<b>46</b>
<b>CMM-208.4: Non-Indications</b>	<b>47</b>
<b>CMM-208.5: Procedure (CPT®) Codes</b>	<b>48</b>
<b>CMM-208.6: References</b>	<b>48</b>

### CMM-208.1: Definitions

- **Radiofrequency joint denervation/ablation** (i.e., facet neurotomy, facet rhizotomy) refers to the insertion of a radiofrequency probe towards the medial branch of the posterior primary rami, which supplies the innervation to the facet joints under fluoroscopic guidance. The radiofrequency electrode is then utilized to create a “continuous” heat lesion by coagulating the nerve supplying the joint with the intention of providing pain relief by denervating the painful facet joint. The injection/block applies directly to the facet joint(s) blocked/ablated and not to the number of nerves blocked/ablated that innervate the facet joint(s).

### CMM-208.2: General Guidelines

- The determination of medical necessity for the performance of radiofrequency joint denervations/ablations is always made on a case-by-case basis.
- When performing radiofrequency joint denervations/ablations, it may be necessary to perform the procedure at the same level(s) bilaterally; however, no more than three (3) levels should be performed during the same session/procedure.
- When performing a repeat radiofrequency joint denervation/ablation at the same spinal level(s) as a prior successful denervation/ablation procedure, further diagnostic facet joint injections/medial branch blocks at that spinal level(s) are not necessary.

### CMM-208.3: Indications

- A radiofrequency joint denervation/ablation is considered **medically necessary** for facet mediated pain resulting from disease, injury, or surgery when **ALL** of the following are met:
  - ◆ Clinical findings and imaging studies suggest no other obvious cause of the pain (e.g., central spinal stenosis with neurogenic claudication/myelopathy, foraminal stenosis or disc herniation with concordant radicular pain/radiculopathy, infection, tumor, fracture, pseudoarthrosis, pain related to spinal instrumentation).
  - ◆ Failure of at least three (3) months of conservative therapy (e.g., exercise, physical methods including physical therapy, chiropractic care, nonsteroidal anti-inflammatory drugs [NSAID’s] and/or analgesics) unless contraindicated and the reason(s) for the contraindication(s) is/are documented in the medical record
  - ◆ Two positive diagnostic facet joint injections/medial branch blocks as evidenced by at least 80% relief of facet mediated pain for at least the expected minimum duration of the effect of the local anesthetic used.
- For an individual with a prior spinal fusion, radiofrequency joint denervation/ablation is considered **medically necessary** when the above criteria are met and the procedure is performed at an unfused spinal segment located either above or below the fused spinal segment.

- A repeat radiofrequency joint denervation/ablation when **BOTH** of the following criteria are met:
  - ◆ There is documented pain relief of at least 50% which has lasted for a minimum of 12 weeks
  - ◆ The procedure is performed at a minimum of six months following the prior denervation/ablation.

#### **CMM 208.4: Non-Indications**

- Performance of a radiofrequency joint denervation/ablation for **ANY** of the following indications is considered **not medically necessary**:
  - ◆ When performed without the use of fluoroscopic guidance
  - ◆ Performing more than two procedures at the same level(s) during a 12 month period of time
  - ◆ In the absence of two sequential positive diagnostic facet joint injections/medial branch blocks at the same level(s) for an initial radiofrequency treatment, or for a repeat radiofrequency treatment in the absence of at least 50% relief of facet mediated pain for at least 6 months from a previous radiofrequency treatment at the same level(s).
  - ◆ When performed for neck pain or low back pain in the presence of an untreated radiculopathy
  - ◆ When performed at a posteriorly fused spinal motion segment (with the exception of patients with clinically suspected pseudarthrosis)
  - ◆ When performed on more than three (3) contiguous spinal joint levels during the same session/procedure
  - ◆ When performed to treat pain arising from above C2-3 and below L5-S1 spinal levels
- Performance of radiofrequency joint denervation/ablations for **ANY** of the following indications is considered **experimental, investigational, or unproven**:
  - ◆ Pulsed radiofrequency ablation for chronic pain syndromes
  - ◆ Endoscopic radiofrequency denervation/endoscopic dorsal ramus rhizotomy
  - ◆ Cryoablation/cryoneurolysis/cryodenervation
  - ◆ Chemical ablation (e.g., alcohol, phenol, glycerol)
  - ◆ Laser ablation
  - ◆ Ablation by any method for sacroiliac (SI) joint pain
  - ◆ Cooled radiofrequency ablation

## CMM-208.5: Procedure (CPT®) Codes

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
64633	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT), cervical or thoracic, single facet joint
64634	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT), cervical or thoracic, each additional facet joint (List separately in addition to code for primary procedure)
64635	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT), lumbar or sacral, single facet joint
64636	Destruction by neurolytic agent, paravertebral facet joint nerve(s), with imaging guidance (fluoroscopy or CT), lumbar or sacral, each additional facet joint (List separately in addition to code for primary procedure)

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.

## CMM-208.6: References

1. American College of Occupational and Environmental Medicine. Occupational Medicine Practice Guideline, 2nd Ed. 2008.
2. American Medical Association. *Current Procedural Terminology* – Professional Edition.
3. Abejon D, Garcia-del-Valle S, Fuentes M, et al. Pulsed radiofrequency in lumbar radicular pain: clinical effects in various etiological groups. *Pain Pract.* 2007 7(1):21-26.
4. Barnsley L. Percutaneous radiofrequency neurotomy for chronic neck pain: outcomes in a series of consecutive patients. *Pain Med.* 2005;6(4):282-286.
5. Boswell M, Colson J, Sehgal N, et al. A systematic review of therapeutic facet joint interventions in chronic spinal pain. *Pain Physician* 2007;10:229-253.
6. Boswell MV, Manchikanti L, Kaye AD, et al. A best-evidence systematic appraisal of the diagnostic accuracy and utility of facet (zygapophysial) joint injections in chronic spinal pain. *Pain Physician.* 2015;18:E497-E533.
7. Boswell M, Shah R, Everett C, et al. Interventional techniques in the management of chronic spinal pain: evidence-based practice guidelines. *Pain Physician.* 2005;8(1):1-47.
8. Buijs E, van Wijk R, Geurts J, Weeseman et al. Radiofrequency lumbar facet denervation: A comparative study of the reproducibility of lesion size after 2 current radiofrequency techniques. *Reg Anesth Pain Med.* 2004;29(5):400-407.
9. Cahana A, Van Zundert J, Macrea L, et al. Pulsed radiofrequency: current clinical and biological literature available. *Pain Med.* 2006 t;7(5):411-423.
10. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg.* 2012; 22:200-206.
11. Cohen S, Hurley R, Buckenmaier C, et. al. Randomized placebo-controlled study evaluating lateral branch radiofrequency denervation for sacroiliac joint pain. *Anesthesiology.* 2008; 109(2): 279–288.
12. Cohen S, Hurley R, Christo P, Winkley et al. Clinical predictors of success and failure for lumbar facet radiofrequency denervation. *Clin J Pain.* 2007;23:45-52.
13. Cohen S, Raja S. Pathogenesis, diagnosis, and treatment of lumbar zygapophysial (facet) joint pain. *Anesthesiology* 2007;106:591-614.
14. Cohen SP, Williams KA, Kurihara C, et al. Multicenter, randomized, comparative costeffectiveness study comparing 0, 1, and 2 diagnostic medial branch (facet joint nerve) block treatment paradigms before lumbar facet radiofrequency denervation. *Anesthesiology.* 2010; 113:395-405.



15. Conlin A, Bhogal S, Sequeira K, Teasell R. Treatment of whiplash-associated disorders -part II: Medical and surgical interventions. *Pain Res Manag*. 2005 10(1):33-40.
16. Dobrogowski J, Wrzosek A, Wordliczek J. Radiofrequency denervation with or without addition of pentoxifylline or methylprednisolone for chronic lumbar zygapophysial joint pain. *Pharmacol Rep*. 2005;57:475-480.
17. Dreyfuss P, Halbrook B, Pauza K, et al. Efficacy and validity of radiofrequency neurotomy for chronic lumbar zygapophysial joint pain. *Spine*. 2000;25(10):1270-1277.
18. Geurts J, van Wijk R, Stolker R, Groen G. Efficacy of radiofrequency procedures for the treatment of spinal pain: a systematic review of randomized clinical trials. *Reg Anesth Pain Med*. 2001; 26(5):394-400.
19. Gofeld M, Jitendra J, Faclier G. Radiofrequency denervation of the lumbar zygapophysial joints: 10-year prospective clinical audit. *Pain Physician*. 2007;10:291-300.
20. Haldeman S, Carroll L, Cassidy JD, et al. Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. The Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders: executive summary. *Spine*. 2008 33(4 Suppl):S5-A7.
21. Hancock M, Maher C, Latimer J, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J*. 2007;16(10):1539-1550.
22. Hooten W, Martin D, Huntoon M. Radiofrequency neurotomy for low back pain: evidence-based procedural guidelines. *Pain Med*. 2005;6(2):129-138.
23. King W, Ahmed SU, Baisden J, Patel N, Kennedy DJ, MacVicar J, Duszynski B. Diagnosis and treatment of posterior sacroiliac complex pain: a systematic review with comprehensive analysis of the published data. *Pain Med* 2015 Feb; 16(2): 257.
24. Koizuka S, Saito S, Kawauchi C, et al. Percutaneous radiofrequency lumbar facet rhizotomy guided by computed tomography fluoroscopy. *J Anesth*. 2005;19(2):167-169.
25. Joo YC, Park JY, Kim KH. Comparison of alcohol ablation with repeated thermal radiofrequency ablation in medial branch neurotomy for the treatment of recurrent thoracolumbar facet joint pain. *J Anesth*. 2013; 27:390-395.
26. Lakemeier S, Lind M, Schultz W, et al. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: A randomized, controlled, double-blind trial. *Anesth Analg*. 2013;117:228-235.
27. Laslett M, McDonald B, Aprill C, et al. Clinical predictors of screening lumbar zygapophysial joint blocks: development of clinical prediction rules. *Spine J*. 2006 ;6(4):370-379.
28. Laslett M, Oberg B, Aprill C, McDonald B. Zygapophysial joint blocks in chronic low back pain: a test of Revel's model as a screening test. *BMC Musculoskelet Disord*. 2004;5:43.
29. Leclaire R, Fortin L, Lambert R, et al. Radiofrequency facet joint denervation in the treatment of low back pain: a placebo-controlled clinical trial to assess efficacy. *Spine*. 2001;26(13):1411-1416.
30. Lee CH, Chung CK, Kim CH. The efficacy of conventional radiofrequency denervation in patients with chronic low back pain originating from the facet joints: a meta-analysis of randomized controlled trials. *Spine J*. 2017 Nov;17(11):1770-1780.
31. Lord S, Barnsley L, Wallis B, McDonald G, Bogduk N. Percutaneous radio-frequency neurotomy for chronic cervical zygapophysial-joint pain. *N Engl J Med*. 1996; 335:1721-1726.
32. Maas ET, Ostelo RW, Niemisto L, Jousimaa J, Hurri H, Malmivaara A, van Tulder MW. Radiofrequency denervation for chronic low back pain. *Cochrane Database Syst Rev*. 2015 Oct 23;(10):CD008572. doi: 10.1002/14651858.CD008572.pub2.
33. MacVicar J, Borowczyk JM, MacVicar AM, Loughnan BM, Bogduk N. Cervical medial branch radiofrequency neurotomy in New Zealand. *Pain Med*. 2012;13:647-654.
34. MacVicar J, Kreiner DS, Duszynski B, Kennedy DJ. Appropriate use criteria for fluoroscopically guided diagnostic and therapeutic sacroiliac interventions: results from the spine intervention society convened multispecialty collaborative. *Pain Med*. 2017;18(11):2081-2095.
35. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques of chronic spinal pain: Part II: Guidance and recommendations. *Pain Physician*. 2013;16:S49-S283.
36. Manchikanti L, Kaye AD, Boswell MV, et al. A systematic review and best evidence synthesis of the effectiveness of therapeutic facet joint interventions in managing chronic spinal pain. *Pain Physician*. 2015;18:E535-E582.

37. Manchikanti L, Cash K, Pampati V, Fellows B. Influence of psychological variables on the diagnosis of facet joint involvement in chronic spinal pain. *Pain Physician*. 2008 ;11(2):145-160.
38. Manchikanti L, Manchukonda R, Pampati V, et al. Prevalence of facet joint pain in chronic low back pain in postsurgical patients by controlled comparative local anesthetic blocks. *Arch Phys Med Rehabil*. 2007;88(4):449-455.
39. Manchikanti L, Staats P, Singh V, et al. Evidence-based practice guidelines for interventional techniques in the management of chronic spinal pain. *Pain Phys*. 2003;6:3-81.
40. Manchukonda R, Manchikanti K, Cash K, et al. Facet joint pain in chronic spinal pain: an evaluation of prevalence and false-positive rate of diagnostic blocks. *J Spinal Disord Tech*. 2007;20(7):539-545.
41. McDonald G, Lord S, Bogduk N. Long-term follow-up of patients treated with cervical radiofrequency neurotomy for chronic neck pain. *Neurosurgery* 1999 ;45(1):61-68.
42. Mikeladze G, Espinal R, Finnegan R, et al. Pulsed radiofrequency application in treatment of chronic zygapophyseal joint pain. *Spine J*. 2003;3(5):360-362.
43. Moon JY, Lee PB, Kim YC, Choi SP, Sim WS. An alternative distal approach for the lumbar medial branch radiofrequency denervation: A prospective randomized comparative study. *Anesth Analg*. 2013;116:1133-1140.
44. Nath S, Nath C, Pettersson K. Percutaneous lumbar zygapophysial (Facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: a randomized double-blind trial. *Spine*. 2008;33(12):1291-1298.
45. Niemisto L, Kalso E, Malmivaara A, et al. Radiofrequency denervation for neck and back pain. *The Cochrane Database of Systematic Reviews*. 2006; Issue 3.
46. Oh W, Shim J. A randomized controlled trial of radiofrequency denervation of the ramus communicans nerve for chronic discogenic low back pain. *Clin J Pain* 2004;20(1):55-60.
47. Patel N, Gross A, Brown L, Gekht G. A randomized, placebo-controlled study to assess the efficacy of lateral branch neurotomy for chronic sacroiliac joint pain. *Pain Medicine*. 2012;13:383-398.
48. Sanders M, Zuurmond W. Percutaneous intra-articular lumbar facet joint denervation in the treatment of low back pain: a comparison with percutaneous extra-articular lumbar facet denervation. *Pain Clinic*. 1999;11(4):329-335.
49. Sapir DA, Gorup JM. Radiofrequency medial branch neurotomy in litigant and non-litigant patients with cervical whiplash. *Spine (Phila Pa 1976)*. 2001;26:E268-E273.
50. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine*. 2004;29(21):2471-2473.
51. Slipman C, Bhat A, Gilchrist R, et al. A critical review of the evidence for the use of zygapophysial injections and radiofrequency denervation in the treatment of low back pain, *Spine J*. 2003;3(4):310-316.
52. Speldewinde GC. Outcomes of percutaneous zygapophysial and sacroiliac joint neurotomy in a community setting. *Pain Med*. 2011;12:209-218.
53. Tekin I, Mirzai H, Ok G, Erbuyun et al. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain*. 2007;23(6):524-529.
54. Tzaan W, Tasker R. Percutaneous radiofrequency facet rhizotomy--experience with 118 procedures and reappraisal of its value. *Canadian Journal of Neurological Sciences*. 2000;27(2):125-130.
55. Vallejo R, Benyamin R, Kramer J, et al. Pulsed radiofrequency denervation for the treatment of sacroiliac joint syndrome. *Pain Med*. 2006;7(5):429-434.
56. van Kleef M, Barendse G, Kessels A, et al. Randomized trial of radiofrequency lumbar facet denervation for chronic low back pain. *Spine*. 1999;24(18):1937-1942.
57. van Wijk R, Geurts J, Wynne H, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain*. 2005; 21(4):335-344.
58. Van Zundert J, Patihh J, Kessels A, et al. Pulsed radiofrequency adjacent to the cervical dorsal root ganglion in chronic cervical radicular pain: a double-blind sham controlled randomized clinical trial. *Pain*. 2007;127:173-182
59. Washington State Department of Labor and Industries. Guideline on diagnostic facet medial nerve branch blocks and facet neurotomy. *Provider Bull*. 2005;(PB 05-11):1-6.
60. Workloss Data Institute. Official Disability Guidelines 2008.

<b>CMM-209: Regional Sympathetic Blocks</b>	
<b>CMM-209.1: Definitions</b>	<b>52</b>
<b>CMM-209.2: General Guidelines</b>	<b>53</b>
<b>CMM-209.3: Indications</b>	<b>53</b>
<b>CMM-209.4: Non-Indications</b>	<b>54</b>
<b>CMM-209.5: Procedure (CPT®) Codes</b>	<b>54</b>
<b>CMM-209.6: References</b>	<b>55</b>

## **CMM-209.1: Definitions**

- **Regional sympathetic blocks** (i.e., Stellate Ganglion Blocks and Lumbar Sympathetic Blocks) refer to the injection of local anesthetic along the sympathetic ganglia of the under fluoroscopy to reduce sympathetic nervous system activity. A diagnostic regional sympathetic block is considered positive when there is significant reduction in pain and improvement in function for the duration of the local anesthetic used.
- **Complex Regional Pain Syndrome (CRPS)** is defined by the International Association for the Study of Pain (IASP) as a variety of painful conditions following injury which appear regionally having a distal predominance of abnormal findings, exceeding in both magnitude and duration the expected clinical course of the inciting event and often resulting in significant impairment of motor function, and showing variable progression over time. In addition to injury, CRPS can also occur as a result of various medical disorders or illnesses. The diagnostic criteria for CRPS are as follows:
  - ◆ Continuing pain that is disproportionate to any inciting event
  - ◆ Must report at least one (1) of the symptoms in the following categories:
    - Sensory: reports of hyperesthesia
    - Vasomotor: reports of temperature asymmetry, skin color changes, and/or skin color asymmetry
    - Sudomotor/edema: reports of edema, sweating changes, and/or sweating asymmetry
    - Motor/trophic: reports of decreased range of motion, motor dysfunction (weakness, tremor, dystonia), and/or trophic changes (hair, nail, skin).
  - ◆ Must display at least one (1) of the signs on physical examination in **TWO or MORE** the following categories:
    - Sensory: evidence of hyperalgesia (to pinprick) and/or allodynia (to light touch)
    - Vasomotor: evidence of temperature asymmetry, skin color changes, and/or asymmetry
    - Sudomotor/edema: evidence of edema, sweating changes, and/or sweating asymmetry
    - Motor/trophic: evidence of decreased range of motion, motor dysfunction (weakness, tremor, dystonia). and/or trophic changes (hair, nail, skin).

## **CMM-209.2: General Guidelines**

- The determination of medical necessity for the performance of regional sympathetic blocks is always made on a case-by-case basis.
- **Please note:** this guideline does not apply to injections/blocks of other autonomic nerves (e.g. sphenopalatine ganglion, carotid sinus, superior hypogastric plexus, celiac plexus, Gasserian ganglion [trigeminal nerve], splanchnic nerve, Ganglion of Impar, rami communicans).
- Regional sympathetic blocks should be performed using fluoroscopy.
- Due to insufficient evidence that regional sympathetic blocks (Stellate Ganglion Blocks and Lumbar Sympathetic Chain Blocks) performed as an isolated treatment alter the long term outcome of CRPS, all regional sympathetic blocks in recalcitrant cases of CRPS should be performed with the intent of facilitating involvement and advancement in an active rehabilitation/functional restoration program.

## **CMM-209.3: Indications**

- The performance of an initial diagnostic regional sympathetic block is considered **medically necessary** to establish the presence or absence of sympathetically mediated complex regional pain syndrome. A positive response is defined as at least 50% reduction in pain and improvement in function for the duration of the local anesthetic used.
- Following a successful initial diagnostic block, three (3) additional regional sympathetic blocks, performed within the first two (2) weeks of the initial block, may be considered **medically necessary** to diagnose the individual's pain and obtain a therapeutic response.
- Additional therapeutic regional sympathetic blocks are considered **medically necessary** when provided as part of a comprehensive pain management program and **ALL** of the following criteria are met:
  - ◆ Decreased use of pain medication
  - ◆ Increased functional ability (e.g., increased range of motion, strength, and use of the extremity in activities of daily living)
  - ◆ Increased tolerance to touch (e.g., decreased allodynia)
  - ◆ Ongoing participation in an active rehabilitation program
  - ◆ Performed at a frequency of no more than one time per week
  - ◆ No more than six (6) total blocks

### **CMM 209.4: Non-Indications**

Regional sympathetic blocks are considered **not medically necessary** for each of the following:

- When the individual is not capable of participating or is not involved in an ongoing active rehabilitation program
- Without the use of fluoroscopic guidance
- No significant reduction in pain and no improvement in function for the duration of the local anesthetic following the diagnostic block
- A repeat therapeutic block when there is no decrease in use of pain medication, increase in functional ability, and increase of tolerance to touch

### **CMM-209.5: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
<b>64510</b>	Injection, anesthetic agent; stellate ganglion(cervical sympathetic)
<b>64520</b>	Injection, anesthetic agent; lumbar or thoracic(paravertebral sympathetic)

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.



## **CMM-209.6: References**

1. Ackerman W, Zhang J. Efficacy of stellate ganglion blockade for the management of type 1 complex regional pain syndrome. *Southern Medical Journal*. 2006;99(10):1084-1088.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*, 2nd Ed. 2008.
3. American Medical Association. *Current Procedural Terminology – Professional Edition*.
4. Cepeda M, Carr D, Lau J. Local anesthetic sympathetic blockade for complex regional pain syndrome. *Cochrane Database Syst Rev*. 2005 Oct 19;4:CD004598.
5. Cepeda M, Lau J, Carr D. Defining the therapeutic role of local anesthetic sympathetic blockade in complex regional pain syndrome: a narrative and systematic review. *Clin J Pain*. 2002;18:216-233.
6. Chou R, Huffman L. American Pain Society. American College of Physicians. Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Annals of Internal Medicine*. 2007;147(7):505-14.
7. Chou R, Huffman LH American Pain Society. American College of Physicians. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Annals of Internal Medicine*. 2007;147(7):492-504.
8. Chou R, Qaseem A, Snow V, et al. Clinical Efficacy Assessment Subcommittee of the American College of Physicians. American College of Physicians. American Pain Society Low Back Pain Guidelines Panel. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Annals of Internal Medicine*. 2007;147(7):478-491.
9. Chou R. Using evidence in pain practice: Part I: Assessing quality of systematic reviews and clinical practice guidelines. *Pain Medicine*. 2008; 9(5):518-530.
10. Forouzanfar T, Köke A, van Kleef M, Weber W. Treatment of complex regional pain syndrome type I. *Eur J Pain*. 2002;6:105-122.
11. Furlan A, Mailis A, Papagapiou M. Are we paying a high price for surgical sympathectomy? A systemic literature review of late complications. *J Pain*. 2000;1:245-257.
12. Galer B, Bruehl S, Harden R. IASP diagnostic criteria for complex regional pain syndrome: a preliminary empirical validation study. *Clin J Pain*. 1998;14:48–54.
13. Harden R, Bruehl S, Galer B, et al. Complex regional pain syndrome: are the IASP diagnostic criteria valid and sufficiently comprehensive? *Pain* 1999;83:211-219.
14. Harden R, Bruehl S, Perez R, et al. Validation of proposed diagnostic criteria (the “Budapest criteria”) for complex regional pain syndrome. *PAIN* 2010;150:268 – 74.
15. Harden R, Bruehl S, Stanton-Hicks M, Wilson P. Proposed new diagnostic criteria for complex regional pain syndrome. *Pain Med*. 2007;8(4):326-331.
16. Hartrick C, Kovan J, Naismith P. Outcome prediction following sympathetic block for complex regional pain syndrome. *Pain Pract*. 2004;4:222-228.
17. Harvey AM. Classification of chronic pain – descriptions of chronic pain syndromes and definitions of pain terms. *Clin J Pain* 1995; 11(2); 179.
18. Janig W, Baron R. Complex regional pain syndrome: mystery explained? *Lancet Neurol*. 2003;2:687-697.
19. Leis S, Weber M, Schmelz M, Birklein F. Facilitated neurogenic inflammation in unaffected limbs of patients with complex regional pain syndrome. *Neurosci Lett*. 2004;359:163-166.
20. Ozturk E, Mohur H, Arslan N, et al. Quantitative three-phase bone scintigraphy in the evaluation of intravenous regional blockade treatment in patients with stage-I reflex sympathetic dystrophy of upper extremity. *Annals of Nuclear Medicine*. 2004;18(8):653-658.
21. Paraskevas K, Michaloglou A, Briana D, Samara M. Treatment of complex regional pain syndrome type I of the hand with a series of intravenous regional sympathetic blocks with guanethidine and lidocaine. *Clin Rheumatol*. 2005;7:1-7.
22. Perez R, Kwakkel G, Zuurmond W, de Lange J. Treatment of reflex sympathetic dystrophy (CRPS type 1): a research synthesis of 21 randomized clinical trials. *J Pain Sympt Management*. 2001;21:511-526.
23. Schurmann M, Gradl G, Wizgal I, et al. Clinical and physiologic evaluation of stellate ganglion blockade for complex regional pain syndrome type I. *Clin J Pain*. 2001;17:94-100.

24. Severens J, Oerlemans H, Weegels A, et al. Cost-effectiveness analysis of adjuvant physical or occupational therapy for patients with reflex sympathetic dystrophy. *Arch Phys Med Rehabil.* 1999;80:1038–1043
25. Sharma A, Williams K, Raja S. Advances in treatment of complex regional pain syndrome: recent insights on a perplexing disease. *Curr Opin Anaesthesiol.* 2006;19:566-72.
26. Stanton-Hicks M. A report on the 2nd IASP Research Symposium, Cardiff, Wales. Complex regional pain syndrome: current research on mechanisms and diagnosis. In: International Association for the Study of Pain. Special Interest Group on Pain and the Sympathetic Nervous System. 2000; 1– 2.
27. Stanton-Hicks M. In: Wakefield CA, Bajwa JH. *Principles and Practice of Pain Medicine.* 2nd ed. 2004.
28. Stanton-Hicks M, Baron R, et al. Consensus report: complex regional pain syndromes: guidelines for therapy. *Clin J Pain.* 1998;14:155–166
29. Stanton-Hicks M, Burton A, Bruhl S, et al. An updated interdisciplinary clinical pathway for CRPS: report of an expert panel. *Pain Practice.* 2005;84(3):S4-S16.
30. Stanton-Hicks M. Complex regional pain syndrome: manifestations and the role of neurostimulation in its management. *J Pain Symptom Manage.* 2006;(4 Suppl):S20-S4.
31. Suresh S, Wheeler M, Patel A. Case series: IV regional anesthesia with ketorolac and lidocaine: is it effective for the management of complex regional pain syndrome 1 in children and adolescents? *Anesth Analg.* 2003;96:694-695.
32. Turner J, Loeser J, Deyo R, Sanders S. Spinal cord stimulation for patients with failed back surgery syndrome or complex regional pain syndrome: a systematic review of effectiveness and complications. *Pain.* 2004;108:137-147.
33. Varrassi G, Paladini A, Marinangeli F, Racz G. Neural modulation by blocks and infusions. *Pain Pract.* 2006;6:34-38.
34. Wang L, Chen H, Chang P, et al. Axillary brachial plexus block with patient controlled analgesia for complex regional pain syndrome type I: a case report. *Reg Anesth Pain Med.* 2001;26(1):68-71.
35. Workloss Data Institute. *Official Disability Guidelines* 2009.

<b>CMM-210: Implantable Intrathecal Drug Delivery Systems</b>	
<b>CMM-210.1: Definitions</b>	<b>58</b>
<b>CMM-210.2: General Guidelines</b>	<b>58</b>
<b>CMM-210.3: Indications</b>	<b>58</b>
<b>CMM-210.4: Non-indications</b>	<b>60</b>
<b>CMM-210.5: Replacement</b>	<b>60</b>
<b>CMM-210.6: Procedure (CPT®) Codes</b>	<b>60</b>
<b>CMM-210.7: References</b>	<b>62</b>

### **CMM-210.1: Definitions**

- **An implantable intrathecal drug delivery system** (Pain pump or Baclofen pump) is a device used for the continuous infusion of a drug directly into the cerebrospinal fluid via a catheter placed in the intrathecal or epidural space. A pump is placed in the subcutaneous tissue of the abdomen and connected to the catheter. The pump reservoir holds the medication(s), and the pump is programmed to give a set dose of medication over time. For most individuals, it should be used as part of a program to facilitate restoration of function and return to activity, and not just for pain reduction. An intrathecal drug delivery trial can be accomplished by either a single intrathecal bolus injection or an intrathecal catheter infusion.

### **CMM-210.2: General Guidelines**

- **Please note:** this guideline does not apply to epidural injections administered for obstetrical or surgical epidural anesthesia.
- The determination of medical necessity for the performance of an implantable intrathecal or epidural drug delivery system is always made on a case-by-case basis.

### **CMM-210.3: Indications**

- The use of an implantable intrathecal or epidural drug delivery system is considered **medically necessary** for **ANY** of the following indications when the associated criteria are met:
  - ◆ Nonmalignant, chronic intractable pain (e.g., failed back surgery syndrome with low back pain and/or radicular pain, complex regional pain syndrome [i.e., reflex sympathetic dystrophy], post-herpetic neuralgia)
  - ◆ Severe, refractory spasticity or chronic intractable dystonia in individuals who are unresponsive to or cannot tolerate oral anti-spasticity agents (i.e., Baclofen [Lioresal<sup>®</sup>]) (i.e., intrathecal injection of Baclofen)
  - ◆ Cancer-related pain

### **Nonmalignant, Chronic Intractable Pain**

- A trial with a percutaneous intrathecal or epidural drug delivery system for nonmalignant chronic intractable pain is considered **medically necessary** when **ALL** of the following criteria have been met:
  - ◆ There is a documented pathology (i.e., an objective basis for the pain complaint)
  - ◆ Failure of a sufficient trial of at least six (6) months of provider-directed noninvasive pain management, including active rehabilitative exercise and fixed schedule dosing of opioids or other analgesics unless contraindicated and the reason(s) for the contraindication(s) is/are documented in the medical record
  - ◆ Further surgical intervention or other treatment is not indicated or likely to be effective
  - ◆ Statement from a primary care physician, neurologist, physiatrist, psychiatrist, psychologist, or other licensed behavioral and/or medical health care provider attesting to the absence of untreated, underlying mental health conditions/issues (e.g., depression, drug, alcohol abuse) as a major contributor to chronic pain.

- ◆ Individual agrees to a 50% reduction in systemic opiates prior to undergoing an intrathecal opiate trial.
- A permanent implantable intrathecal or epidural drug delivery system for the above listed pain conditions is considered **medically necessary** if the individual has met the above criteria for a preliminary trial and has experienced > 50% reduction in pain for 8 hours and concomitant increase in function during an appropriate trial.

### **Severe, Refractory Spasticity/Chronic Intractable Dystonia**

- A trial with a percutaneous intrathecal drug delivery system for severe, refractory spasticity or chronic intractable dystonia is considered **medically necessary** for **EITHER** of the following indications:
  - ◆ There is failure, contraindication or intolerance to at least a six-week trial of oral antispasmodic drugs and physical therapy.
  - ◆ Individual has a baseline average Ashworth score of at least 3 (or a Modified Ashworth score of 2) and a Spasm Frequency score of at least 2.
    - An Ashworth score of 3 represents a considerable increase in muscle tone when testing resistance to passive movement about a joint with varying degrees of velocity.
    - A Modified Ashworth score of 2 represents a slight increase in muscle tone followed by minimal resistance of the range of motion.
    - A Spasm Frequency score of 2 represents a patient's self-report of between 1 to 5 spasms per day.
- A permanent implantable infusion for the treatment of chronic intractable spasticity or chronic intractable dystonia is considered **medically necessary** when a preliminary trial of intrathecal antispasmodic drug administration, that meets the above medical necessity criteria, demonstrates a beneficial clinical response (e.g., demonstrates at least a 2-point reduction in the Ashworth or Spasm Frequency score for 4 hours following an intrathecal trial bolus of Baclofen)

### **Cancer-Related Pain**

- A trial with a percutaneous intrathecal or epidural drug delivery system for cancer-related pain is considered **medically necessary** when there is failure, intolerance, or contraindication to noninvasive methods of pain control, including systemic opioids.
- A permanent implantable intrathecal or epidural drug delivery system for the above listed pain conditions is considered **medically necessary** if the individual has met the above criteria for a preliminary trial and has experienced at least a 50% reduction in pain during an appropriate trial.
- **Please Note:** A trial with a percutaneous intrathecal or epidural drug delivery system for cancer-related pain is not required in the presence of advanced disease, when survival time is limited, and when the individual is considered at high risk for procedures.

### **CMM-210.4: Non-indications**

- An intrathecal or epidural drug delivery system is considered **experimental, investigational or unproven** for **ANY** other indication, including the following:
  - ◆ Cancer-related pain, spastic/dystonic, or other pain conditions that do not meet the above criteria
  - ◆ Administration of insulin for diabetes
  - ◆ Administration of antibiotics for osteomyelitis
  - ◆ Administration of heparin for thromboembolic disease

### **CMM-210.5: Replacement**

- Replacement of an implanted intrathecal or epidural drug infusion system is considered **medically necessary** when **BOTH** of the following criteria have been met:
  - ◆ The existing device is documented to be nearing end of battery life, will no longer be beneficial and cannot be repaired, or a built-in component provides notification of impending failure
  - ◆ There is no evidence to suggest the device has been abused or neglected
- Replacement of an implantable intrathecal infusion pump is considered **not medically necessary** when the existing infusion pump and/or components remain functional.

### **CMM-210.6: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior	
CPT®	Code Description/Definition
62320	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance.
62321	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (ie, fluoroscopy or CT)
62322	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance.
62323	Injection(s), of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, including needle or catheter placement, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); with imaging guidance (ie, fluoroscopy or CT)
62324	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, cervical or thoracic; without imaging guidance.



<b>62325</b>	Injection(s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, cervical or thoracic; with imaging guidance (ie, fluoroscopy or CT)
<b>62326</b>	Injection (s), including indwelling catheter placement, continuous infusion or intermittent bolus, of diagnostic or therapeutic substance(s) (eg, anesthetic, antispasmodic, opioid, steroid, other solution), not including neurolytic substances, interlaminar epidural or subarachnoid, lumbar or sacral (caudal); without imaging guidance.
<b>62350</b>	Implantation, revision or repositioning of tunneled intrathecal or epidural catheter, for long-term medication administration via an external pump or implantable reservoir/infusion pump; without laminectomy
<b>62351</b>	Implantation, revision or repositioning of tunneled intrathecal or epidural catheter, for long-term medication administration via an external pump or implantable reservoir/infusion pump; with laminectomy
<b>62355</b>	Removal of previously implanted intrathecal or epidural catheter
<b>62360</b>	Implantation or replacement of device for intrathecal or epidural drug infusion; subcutaneous reservoir
<b>62361</b>	Implantation or replacement of device for intrathecal or epidural drug infusion; nonprogrammable pump
<b>62362</b>	Implantation or replacement of device for intrathecal or epidural drug infusion; programmable pump, including preparation of pump with or without programming
<b>62365</b>	Removal of subcutaneous reservoir or pump, previously implanted for intrathecal or epidural infusion
<b>62367</b>	Electronic analysis of programmable, implanted pump for intrathecal or epidural drug infusion (includes evaluation of reservoir status, alarm status, drug prescription status); without reprogramming
<b>62368</b>	Electronic analysis of programmable, implanted pump for intrathecal or epidural drug infusion (includes evaluation of reservoir status, alarm status, drug prescription status); with reprogramming
<b>95990</b>	Refilling and maintenance of implantable pump or reservoir for drug delivery, spinal (intrathecal, epidural) or brain (intraventricular), includes electronic analysis of pump, when performed
<b>95991</b>	Refilling and maintenance of implantable pump or reservoir for drug delivery, spinal (intrathecal, epidural) or brain (intraventricular), includes electronic analysis of pump when performed; requiring skill of a physician or other qualified health care professional
This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.	

## CMM-210.7: References

1. Ackerman L, Follett K, Rosenquist R. Long-term outcomes during treatment of chronic pain with intrathecal clonidine or clonidine/opioid combinations. *J Pain Symptom Manage*. 2003;;26(1):668-677.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*, 2nd Ed. 2008.
3. American Medical Association. *Current Procedural Terminology – 2014 Professional Edition*.
4. Anderson V, Burchiel K. A prospective study of long term intrathecal morphine in the management of chronic nonmalignant pain. *Neurosurgery*. 1999;44:289-300.
5. Anderson V, Cooke B, Burchiel K. Intrathecal hydromorphone for chronic nonmalignant pain: a retrospective study. *Pain Med*. 2001;2(4):287-297.
6. Angel I, Gould H Jr, Carey M. Intrathecal morphine pump as a treatment option in chronic pain of nonmalignant origin. *Surg Neurol*. 1998;49(1):92-98.
7. Boswell M, Shah R, Everett C, et al. Interventional Techniques: Evidence-based Practice Guidelines in The Management of Chronic Spinal Pain: Evidence-Based Practice Guidelines. *Pain Physician*. 2005;8:1-47.
8. Brown J, Klapow J, Doleys D, et al. Disease-specific and generic health outcomes: a model for the evaluation of long-term intrathecal opioid therapy in noncancer low back pain patients. *Clin J Pain*. 1999;15:122-131.
9. Dahm P, Nitescu P, Appelgren L, Curelaru I. Efficacy and technical complications of long-term continuous intraspinal infusions of opioid and/or bupivacaine in refractory nonmalignant pain: a comparison between the epidural and the intrathecal approach with externalized or implanted catheters and infusion pumps. *Clin J Pain*. 1998;14:4-16.
10. Dario A, Scamoni C, Picano M, et al. The infection risk of intrathecal drug infusion pumps after multiple refill procedures. *Neuromodulation*. 2005;8(1):36-39.
11. Deer T, Krames ES, Hassenbusch SJ, Burton A, Caraway D, Dupen S, et al. Polyanalgesic consensus conference 2007: Recommendations for the management of pain by intrathecal (intraspinale) drug delivery: Report of an interdisciplinary expert panel. *Neuromodulation*. 2007;10(4):300-328.
12. Deer TR, Hayek SM, Pope JE, et al. The polyanalgesic consensus conference (PACC): recommendations for training of intrathecal drug delivery infusion therapy. *Neuromodulation*. 2017;20:133-154. DOI: 10.1111/ner.12543.
13. Deer TR, Kim C, Bowman R, Tolentino D, Stewart C, Tolentino W. Intrathecal ziconotide and opioid combination therapy for noncancer pain: an observational study. *Pain Physician*. 2009;12(4):E291-E296.
14. Deer TR, Smith HS, Burton AW, et al. Comprehensive Consensus Based Guidelines on Intrathecal Drug Delivery Systems in the Treatment of Pain Caused by Cancer Pain. *Pain Physician*. 2011; 14:E283-E312.
15. Deer TR, Prager J, Levy R, Rathmell J, Buchser E, Burton A, et al. Polyanalgesic Consensus Conference 2012: recommendations for the management of pain by intrathecal (intraspinale) drug delivery: report of an interdisciplinary expert panel. *Neuromodulation*. 2012a;15(5):436-464.
16. Deer TR, Prager J, Levy R, Burton A, Buchser E, Caraway D, et al. Polyanalgesic Consensus Conference--2012: recommendations on trialing for intrathecal (intraspinale) drug delivery: report of an interdisciplinary expert panel. *Neuromodulation*. 2012b;15(5):420-35.
17. Deer T, Chapple I, Classen A, et al. Intrathecal drug delivery for treatment of chronic low back pain: report from the National Outcomes Registry for Low Back Pain. *Pain Med*. 2004;5(1):6-13.
18. Deer T. Current and future trends in spinal cord stimulation for chronic pain. *Curr Pain Headache Rep*. 2001;5(6):503-509.
19. Du Pen S, Du Pen A, Hillyer J. Intrathecal hydromorphone for intractable nonmalignant pain: a retrospective study. *Pain Med*. 2006;7(1):10-15.
20. Guillaume D, Van Havenbergh A, Vloerberghs M, et al. A clinical study of intrathecal baclofen using a programmable pump for intractable spasticity. *Arch Phys Med Rehabil*. 2005;86:2165-2171.

21. Hamza M, Doleys DM, Saleh IA, Medvedovsky A, Verdolin MH, Hamza M. A Prospective, Randomized, Single-Blinded, Head-to-Head Long-Term Outcome Study, Comparing Intrathecal (IT) Boluses With Continuous Infusion Trialing Techniques Prior to Implantation of Drug Delivery Systems (DDS) for the Treatment of Severe Intractable Chronic Nonmalignant Pain. *Neuromodulation*. 2015 Oct;18(7):636-48.
22. Hassenbusch S, Portenoy R, Cousins M, et al. Polyanalgesic Consensus Conference 2003: an update on the management of pain by intraspinal drug delivery--report of an expert panel. *J Pain Symptom Manage*. 2004;27(6):540-563.
23. Krames E. Intraspinal opioid therapy for chronic nonmalignant pain: current practice and clinical guidelines. *J Pain Symptom Manage*. 1996;11(6):333-352.
24. Kumar K, Hunter G, Demeria D. Treatment of chronic pain by using intrathecal drug therapy compared with conventional pain therapies: a cost-effectiveness analysis. *J Neurosurg*. 2002;97(4):803-810.
25. Miele V, Price K, Bloomfield S, et al. A review of intrathecal morphine therapy related granulomas. *Eur J Pain*. 2006;10(3):251-261.
26. Modified Ashworth scale and spasm frequency score in spinal cord injury: reliability and correlation. *Spinal Cord* (2016) 54, 702–708 (2016).
27. North American Spine Society. NASS Coverage Policy Recommendations: Intrathecal Drug Delivery Systems. Burr Ridge, IL: North American Spine Society. March 2017.
28. Nguyen H, Garber J, Hassenbusch S. Spinal analgesics. *Anesth Clin of NA*. 2003;21(4).
29. Osenbach R, Harvey S. Neuraxial infusion in patients with chronic intractable cancer and noncancer pain. *Curr Pain Headache Rep*. 2001;5(3):241-9.
30. Penn RD et al. Intrathecal baclofen for severe spinal spasticity. *N Engl J Med* 1989; 320: 1517–1521.
31. Raffaelli W, Marconi G, Fanelli G, et al. Opioid-related side-effects after intrathecal morphine: a prospective, randomized, double-blind dose-response study. *Eur J Anaesthesiology*. 2006;23:605-10.
32. Rauck R, Wallace M, Leong M, et al; Ziconotide 301 Study Group. A randomized, double-blind, placebo-controlled study of intrathecal ziconotide in adults with severe chronic pain. *J Pain Symptom Manage*. 2006;31(5):393-406.
33. Staal C, Arends A, Ho S. A self-report of quality of life of patients receiving intrathecal baclofen therapy. *Rehabil Nurs*. 2003 Sep-Oct;28(5):159-63.
34. Thimineur M, Kravitz E, Vodapally M. Intrathecal opioid treatment for chronic non-malignant pain: a 3-year prospective study. *Pain*. 2004;109(3):242-249.
35. Turner J, Sears J, Loeser J. Programmable intrathecal opioid delivery systems for chronic noncancer pain: a systematic review of effectiveness and complications. *Clin J Pain*. 2007;23(2):180-95.
36. van Hilten B, van de Beek W, Hoff J, et al. Intrathecal baclofen for the treatment of dystonia in patients with reflex sympathetic dystrophy. *N Engl J Med*. 2000;343(9):625-630.
37. Waara-Wolleat K, Hildebrand K, Stewart G. A review of intrathecal fentanyl and sufentanil for the treatment of chronic pain. *Pain Med*. 2006;7:251-259.
38. Winkelmuller M, Winkelmuller W. Long-term effects of continuous intrathecal opioid treatment in chronic pain of nonmalignant etiology. *J Neurosurg*. 1996;85:458-467.
39. Workloss Data Institute. *Official Disability Guidelines*.
40. Yoshida G, Nelson R, Capen D, Nagelberg et al. Evaluation of continuous intraspinal narcotic analgesia for chronic pain from benign causes. *Am J Orthop*. 1996;25(10):693-694.

## **CMM-211: Spinal Cord Stimulators**

### **Prior Authorization Requirements:**

For Spinal Cord Stimulators, please refer to **Regence SUR Policy No. 45 Spinal Cord and Dorsal Root Ganglion Stimulation.**

**CMM-308: Thermal Intradiscal Procedures**

<b>CMM-308.1: Definitions</b>	<b>66</b>
<b>CMM-308.2: Non-indications</b>	<b>66</b>
<b>CMM-308.3: Procedure (CPT®) Codes</b>	<b>67</b>
<b>CMM-308.4: References</b>	<b>68</b>

### **CMM-308.1: Definitions**

- **Thermal intradiscal procedures** are minimally invasive surgical procedures which involve the percutaneous placement of an intradiscal probe into the suspected painful disc(s) and through the use of radiofrequency energy or electrothermal energy, produce heat to either coagulate and/or disrupt (shrink) type I collagen within the disc for decompression of the disc material. The goal of thermal intradiscal procedures is to treat symptomatic patients with annular disruption of contained herniated disc, to seal annular tears or fissures, or destroy nociceptors for the purpose of relieving pain. These techniques include those that use single or multiple probes/catheters, which utilize a resistance coil or other delivery system technology, are flexible or rigid, and are placed within the nucleus, the nuclear-annular junction or the annulus.
- **Thermal intradiscal procedures** include, but are not limited to:
  - ◆ Annulo-nucleoplasty (The Disc-FX procedure)
  - ◆ Cervical intradiscal radiofrequency lesioning
  - ◆ Coblation percutaneous disc decompression
  - ◆ Intradiscal biacuplasty (IDB)/intervertebral disc biacuplasty/cooled radiofrequency
  - ◆ Intradiscal electrothermal annuloplasty (IEA)
  - ◆ Intradiscal electrothermal therapy (IDET)
  - ◆ Intradiscal thermal annuloplasty (IDTA)
  - ◆ Nucleoplasty (also known as percutaneous radiofrequency thermomodulation or percutaneous plasma discectomy)
  - ◆ Percutaneous (or plasma) disc decompression (PDD)
  - ◆ Percutaneous intradiscal radiofrequency thermocoagulation (PIRFT)/intradiscal radiofrequency thermomodulation/percutaneous radiofrequency thermomodulation
  - ◆ Radiofrequency annuloplasty (RA)
  - ◆ Targeted disc decompression (TDD)

### **CMM-308.2: Non-indications**

- Based on the lack of conclusive scientific evidence demonstrating the clinical efficacy of thermal intradiscal procedures and the potential to expose patients to serious adverse side effects or complications, the use of thermal intradiscal procedures are considered **experimental, investigational, or unproven**.



**CMM-308.3: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required. Pre- authorization requirements vary by individual payor.

CPT®	Code Description/Definition
<b>22526</b>	Percutaneous intradiscal electrothermal annuloplasty, unilateral or bilateral including fluoroscopic guidance; single level
<b>22527</b>	Percutaneous intradiscal electrothermal annuloplasty, unilateral or bilateral including fluoroscopic guidance; 1 or more additional levels (list separately in addition to code for primary procedure)
<b>62287</b>	Decompression procedure, percutaneous, of nucleus pulposus of intervertebral disc, any method utilizing needle based technique to remove disc material under fluoroscopic imaging or other form of indirect visualization, with discography and/or epidural injection(s) at the treated level(s), when performed, single or multiple levels, lumbar
<b>62292</b>	Injection procedure for chemonucleolysis, including discography, intervertebral disc, single or multiple levels, lumbar
<b>0274T</b>	Percutaneous laminotomy/laminectomy (interlaminar approach) for decompression of neural elements, (with or without ligamentous resection, discectomy, facetectomy and/or foraminotomy), any method, under indirect image guidance (e.g., fluoroscopy, CT), single or multiple levels, unilateral or bilateral; cervical or thoracic
<b>0275T</b>	Percutaneous laminotomy/laminectomy (interlaminar approach) for decompression of neural elements, (with or without ligamentous resection, discectomy, facetectomy and/or foraminotomy), any method, under indirect image guidance (e.g., fluoroscopy, CT), single or multiple levels, unilateral or bilateral; lumbar
<p>This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the individual payor (health insurance company, etc.) and is based on the member/patient/client/beneficiary's policy or benefit entitlement structure as well as any third party payor guidelines and/or claims processing rules. Providers are strongly urged to contact each payor for individual requirements if they have not already done so.</p>	

## CMM-308.4: References

1. Airaksinen O, Brox J, Cedraschi C, et al; On behalf of the COST B13 Working Group on Guidelines for Chronic Low Back Pain. Chapter 4 European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J*. 2006 Mar;15(Supplement 2):s192-s300.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*, 2nd Ed. 2008.
3. American Medical Association. *Current Procedural Terminology – Professional Edition*.
4. Andersson G, Mekhail N, Block J. Treatment of intractable discogenic low back pain. A systematic review of spinal fusion and intradiscal electrothermal therapy (IDET). *Pain Physician*. 2006;9:237-48.
5. Appleby D, Andersson G, Totta M. Meta-analysis of the efficacy and safety of intradiscal electrothermal therapy (IDET). *Pain Med*. 2006;7(4):308-16
6. Barendse G, van Den Berg S, Kessels A et al. Randomized controlled trial of percutaneous intradiscal radiofrequency thermocoagulation for chronic low back pain: lack of effect from a 90 second 70 C lesion. *Spine* 2001;26(3):287-92
7. Bogduk N, Karasek M, Two-year follow-up of a controlled trial of intradiscal electrothermal anuloplasty for chronic low back pain resulting from internal disc disruption, *Spine J*. 2002 Sep-Oct;2(5):343-50.
8. Boswell M, Trescot A, Datta S, et al American Society of Interventional Pain Physicians.
9. Interventional techniques: evidence-based practice guidelines in the management of chronic spinal pain. *Pain Physician*. 2007;10:7-111.
10. Chou L, Lew H, Coelho P, Slipman C. Intradiscal electrothermal anuloplasty. *Am J Phys Med Rehabil*. 2005;84(7):538-49.
11. Chou R. American Pain Society; American College of Physicians. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for a clinical practice guideline. Recommendations on surgery and interventional treatments. American Pain Society 27th Annual Scientific Meeting: Symposium 312. Presented May 8, 2008.
12. Cohen S, Williams S, Kurihara C, et al. Nucleoplasty With or Without Intradiscal Electrothermal Therapy (IDET) as a Treatment for Lumbar Herniated Disc. *J Spinal Disord Tech*. 2005 Feb;18:S119-S124.
13. Davis T, Delamarter R, Sra P, et al. The IDET procedure for chronic discogenic low back pain, *Spine*. 2004 Apr 16;29(7):752-6.
14. Derby R, Baker R, Lee C, Anderson P. Evidence-informed management of chronic low back pain with intradiscal electrothermal therapy. *Spine J*. 2008;8:80-95.
15. Freeman B, Fraser R, Cain C, et al. A randomized, double-blind, controlled trial. Intradiscal electrothermal therapy versus placebo for the treatment of chronic discogenic low back pain. *Spine*. 2005;30(21):2369-77.
16. Freeman B. IDET: a critical appraisal of the evidence. *Eur Spine J*. 2006 Jul 26.
17. Gerszten P, Welch W, McGrath P et al. A prospective outcomes study of patients undergoing intradiscal electrothermy (IDET) for chronic low back pain. *Pain Physician* 2002;5(4):360-364
18. Gibson J, Waddell G. Surgery for degenerative lumbar spondylosis: updated Cochrane Review. *Spine*. 2005 Oct 15;30(20):2312-20.
19. Heary R. Intradiscal electrothermal anuloplasty: the IDET procedure. *J Spinal Disord*. 2001;14(4):353-60.
20. Kapural L, Hayek S, Malak O, et al. Intradiscal thermal anuloplasty versus intradiscal radiofrequency ablation for the treatment of discogenic pain: a prospective matched trial. *Pain Med* 2005;6(6):425-31.
21. Kapural L, Mekhail N, Korunda Z, Basali A. Intradiscal thermal anuloplasty for the treatment of lumbar discogenic pain in patients with multilevel degenerative disc disease. *Anesth Analg*. 2004;99:472-6.
22. Karasek M, Bogduk N. Twelve-month follow-up of a controlled trial of intradiscal thermal anuloplasty for back pain due to internal disc disruption. *Spine* 2000;25(20):2601-2607.
23. Kloth D, Fenton D, Andersson G, Block J. Intradiscal electrothermal therapy (IDET) for the treatment of discogenic low back pain: patient selection and indications for use. *Pain Physician*. 2008;11:659-68.

24. Lutz C, Lutz G, Cooke P. Treatment of chronic lumbar discogenic pain with intradiscal electrothermal therapy: a prospective outcome study. *Arch Phys Med Rehabil* 2003;84: 23-28.
25. Pauza K, Howell S, Dreyfuss P, et al. A randomized, placebo-controlled trial of intradiscal electrothermal therapy (IDET) for the treatment of discogenic low back pain. *Spine J*. 2004;4(1):27-35.
26. Saal J, Saal J. Intradiscal electrothermal therapy for the treatment of chronic discogenic low back pain. *Clin Sports Med*. 2002;21(1):167-87.
27. Saal J, Saal JS Intradiscal electrothermal treatment for chronic discogenic low back pain: a prospective outcome study with minimum 1-year follow-up. *Spine* 2000 Oct 15;25(20):2622-7.
28. Saal J, Saal J. Intradiscal electrothermal treatment for chronic discogenic low back pain: prospective outcome study with a minimum 2-year follow-up, *Spine* 2002 May 1;27(9):966-73; discussion 973-4.
29. Saal J, Saal J. Management of chronic discogenic low back pain with a thermal intradiscal catheter. A preliminary report. *Spine* 2000 Feb 1;25(3):382-8.
30. Stanos S, New Clinical Practice Guidelines on Low Back Pain. *APS Bulletin*, 2007;18(2).
31. Urrútia G, Kovacs F, Nishishinya M, Olabe J. Percutaneous thermocoagulation intradiscal techniques for discogenic low back pain. *Spine*. 2007;32(10):1146-54
32. Webster B, Verma S, Pransky G, Outcomes of workers' compensation claimants with low back pain undergoing intradiscal electrothermal therapy, *Spine*. 2004 Feb 15;29(4):435-41.
33. Wetzel F, McNally T, Phillips F. Intradiscal electrothermal therapy used to manage chronic discogenic low back pain: new directions and interventions. *Spine*. 2002;27(22):2621-6.
34. Wetzel F, McNally T. Treatment of chronic discogenic low back pain with intradiscal electrothermal therapy. *J Am Acad Orthop Surg*. 2003 Jan-Feb;11(1):6-11.
35. Workloss Data Institute. Official Disability Guidelines.

**CMM-310: Manipulation of the Spine Under Anesthesia**

<b>CMM-310.1: Indications</b>	<b>71</b>
<b>CMM-310.2: Non-Indications</b>	<b>71</b>
<b>CMM-310.3: Procedure (CPT®) Codes</b>	<b>71</b>
<b>CMM-310.4 References</b>	<b>72</b>

### **CMM-310.1: Indications**

- The use of manipulation of the spine when the patient is either sedated or under general anesthesia may be considered **medically necessary** as a closed treatment of traumatically induced vertebral fracture or dislocation in an emergent situation to mitigate the potential for neurological compromise when the decision for an open reduction has been considered by a qualified physician.
- Manipulation under anesthesia should be performed in conjunction with an active rehabilitation/therapeutic exercise program.

### **CMM-310.2: Non-Indications**

- In the absence of traumatically induced vertebral fracture or dislocation, based on the lack of evidence of long term efficacy and safety, the use of manipulation of the spine under sedation or general anesthesia is considered **not medically necessary**.
- Manipulations performed in isolation without the patient participating in an active rehabilitation program in conjunction with a home exercise program is considered **not medically necessary**.

### **CMM-310.3: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required. Pre- authorization requirements vary by individual payor.

CPT®	Code Description/Definition
<b>22505</b>	Manipulation of spine requiring anesthesia, any region.

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the individual payor (health insurance company, etc.) and is based on the member/patient/client/beneficiary's policy or benefit entitlement structure as well as any third party payor guidelines and/or claims processing rules. Providers are strongly urged to contact each payor for individual requirements if they have not already done so.

## **CMM-310.4: References**

1. American Chiropractic Association 2006 House of Delegates Meeting – Resolutions; Manipulation Under Anesthesia (MUA).
2. American College of Occupational and Environmental Medicine. Occupational Medicine Practice Guideline, 2nd Ed. 2008.
3. Buchmann J, Wende K, Kundt G, Haessler F. Manual treatment effects to the upper cervical apophysial joints before, during, and after endotracheal anesthesia: a placebo-controlled comparison. *Am J Phys Med Rehabil.* 2005 Apr;84(4):251-7.6.
4. Dagenais S, Mayer J, Wooley J, Haldeman S. Evidence-informed management of chronic low back pain with medicine-assisted manipulation. *Spine J.* 2008 Jan-Feb;8(1):142-9.
5. Dan N, Saccasan P. Serious complications of lumbar spinal manipulation. *Med J Austral.* 1983;2(12):672-3.
6. Greenman, P. Manipulation with the patient under anesthesia. *J Am Osteopath Assoc* 1992 Sept; 92(9): 1159-60, 1167-70.
7. Kivimaki J, Pohjolainen T, Malmivaara A, et al. Manipulation under anesthesia with home exercises versus home exercises alone in the treatment of frozen shoulder: a randomized, controlled trial with 125 patients.[see comment]. *Journal of Shoulder & Elbow Surgery.* 16(6):722-6, 2007 Nov-Dec.
8. Kohlbeck F, Haldeman S. Medication-assisted spinal manipulation. *Spine J.* 2002;2(4):288-302.
9. Lavernia C, Cardona D, Rossi MD, Lee D. Multimodal pain management and arthrofibrosis. *Journal of Arthroplasty.* 23(6 Suppl 1):74-9, 2008 Sep.
10. National Academy of Manipulation Under Anesthesia Physicians National Guidelines.
11. Ongley M, Klein R, Dorman T, et al. A new approach to the treatment of chronic low back pain. *Lancet.* 1987;2(8551):143-6.
12. Palmieri N, Smoyak S. Chronic low back pain: a study of the effects of manipulation under anesthesia. *J ManipulativePhysiol Ther.* 2002;25(e8).
13. West D, Mathews R, Miller M, Kent G, Effective management of spinal pain in one hundred seventy-seven patients evaluated for manipulation under anesthesia, *J Manipulative Physiol Ther* 1999 Jun;22(5):299-308.
14. Workloss Data Institute. Official Disability Guidelines.
15. Xiong X, Bean A, Anthony A, et al. Manipulation for cervical spinal dislocation under general anesthesia: serial review for 4 years. *Spinal Cord.* 36(1):21-4, 1998 Jan.



## **CMM-400: Anesthesia Services for Interventional Pain Procedures**

<b>CMM-400.1: Definitions</b>	<b>74</b>
<b>CMM-400.2: General Guidelines</b>	<b>74</b>
<b>CMM-400.3: Indications</b>	<b>75</b>
<b>CMM-400.4: Non-Indications</b>	<b>76</b>
<b>CMM-400.5: Procedure (CPT®) Codes</b>	<b>77</b>
<b>CMM-400.6: References</b>	<b>77</b>

## **CMM-400.1: Definitions**

- **Conscious sedation** includes:
  - ◆ **Minimal sedation (anxiolysis)** indicates a drug-induced state during which patients respond normally to verbal commands. Although cognitive function and coordination may be impaired, ventilation and cardiovascular functions are unaffected<sup>3</sup>.
  - ◆ **Moderate sedation/analgesia (conscious sedation)** indicates a drug-induced depression of consciousness during which patients respond purposefully to verbal commands, either alone or accompanied by light tactile stimulation. No interventions are required to maintain a patent airway, and spontaneous ventilation is adequate. Cardiovascular function is usually maintained<sup>3</sup>.
  - ◆ **Deep sedation/analgesia** is a drug-induced depression of consciousness during which patients cannot be easily aroused but respond purposefully after repeated or painful stimulation<sup>3</sup>.
- **Monitored anesthesia care (MAC)** includes the administration of sedatives and/or analgesics often used for mild to moderate sedation. An essential component of MAC is the preprocedural anesthesia assessment and understanding of the patient's coexisting medical conditions and management of a patient's actual or anticipated physiological derangements or medical problems that may occur during a diagnostic or therapeutic procedure. The provider of MAC must be prepared and qualified to convert to general anesthesia when necessary. MAC is administered by a certified registered nurse anesthetist (CRNA) or anesthesiologist. Additionally, a provider's ability to intervene to rescue a patient's airway from any sedation-induced compromise is a mandatory professional qualification to provide MAC<sup>7</sup>.
- **General anesthesia** is a drug-induced loss of consciousness during which patients are not arousable, even by painful stimulation. The ability to independently maintain ventilatory function is often impaired. Patients often require assistance in maintaining a patent airway, and positive pressure ventilation may be required because of depressed spontaneous ventilation or drug-induced depression of neuromuscular function. Cardiovascular function may be impaired<sup>3</sup>.

## **CMM-400.2: General Guidelines**

- The determination of medical necessity for the performance of monitored anesthesia care (MAC) is always made on a case-by-case basis.
- The medical necessity of monitored anesthesia care (MAC) is:
  - ◆ Evaluated prior to each procedure and the determination is made independent of any prior medical necessity determinations for monitored anesthesia care (MAC); **and**
  - ◆ Only considered once an interventional pain procedure is approved or if the interventional pain procedure does not require prior authorization.
    - Benefits, coverage policies, and eligibility issues pertaining to each health plan may take precedence over eviCore's guidelines. Providers are urged to obtain written instructions and requirements directly from each payor.

### **CMM-400.3: Indications**

Monitored anesthesia care (MAC) is considered **medically necessary** when **EITHER** of the following are met:

- Monitored anesthesia care (MAC) will be used during **ANY** of the following interventional pain procedures<sup>8</sup>:
  - ◆ Regional sympathetic blocks
  - ◆ Radiofrequency ablation of the medial branch nerves
  - ◆ Discography
  - ◆ Spinal cord stimulator trial and permanent implantation
  - ◆ Vertebral augmentation
  - ◆ Implantation of intrathecal drug delivery systems
- There is a presence of **ANY** of the following:
  - ◆ Attestation that a behavioral health professional has determined that severe anxiety, psychiatric condition(s), or cognitive impairment(s) would decrease patient safety during the procedure<sup>13</sup>
  - ◆ Hyperkinetic movement disorders including **ANY** of the following<sup>12</sup>:
    - Acquired/traumatic/hypoxic brain injury/stroke
    - Athetoid cerebral palsy
    - Basal ganglia disease
    - Dystonia
    - Familial paroxysmal choreoathetosis
    - Hemiballismus
    - Huntington's Chorea
    - Multiple sclerosis
    - Paroxysmal kinesigenic choreathetosis
    - Spasticity related involuntary movements
    - Spinal cord injury
  - ◆ Patients at risk for airway obstruction due to an anatomic variation including **ANY** of the following<sup>11,14</sup>:
    - Dysmorphic facial features
    - History of stridor
    - Jaw abnormalities (e.g., micrognathia)
    - Mallampati score of 4
    - Neck abnormalities (e.g., mass)
    - Oral abnormalities (e.g., macroglossia)
    - Pierre-Robin syndrome
    - Trisomy 21
  - ◆ Significant medical condition that increases the risk for complications including **ANY** of the following<sup>10</sup>:
    - Active hepatitis
    - Cardiac disease including **ANY** of the following:
      - Poorly controlled hypertension
      - Implanted pacemaker/defibrillator
      - Moderate to severe reduction in ejection fraction requiring medical treatment

- End stage renal disease requiring dialysis
- Morbid obesity (BMI  $\geq$  40 kg/m<sup>2</sup>)
- Pulmonary disease including poorly controlled COPD requiring oxygen
- Sleep apnea requiring **BOTH** of the following during sleep:
  - BiPAP support
  - Supplemental oxygen

When the criteria for the performance of monitored anesthetic care (MAC) are met, **ALL** of the following criteria must also be met<sup>6</sup>:

- A preoperative evaluation has been performed by a member of the anesthesia delivery team which includes airway examination and medical assessment.
- Informed consent has been obtained with a discussion of alternative sedation options.
- **BOTH** of the following are present during the delivery of monitored anesthesia care (MAC):
  - ◆ Continual monitoring of ventilatory function with capnography to supplement standard monitoring by observation and pulse oximetry
  - ◆ A qualified medical professional to recognize and treat airway complications.
- Recovery from monitored anesthesia care (MAC) will be managed by skilled nursing personnel with direct supervision by a certified registered nurse anesthetist or anesthesiologist.

#### **CMM-400.4: Non-Indications**

- Monitored anesthesia care (MAC) for manipulation of the spine or for closed procedures on the cervical, thoracic, or lumbar spine is considered **experimental, investigational, or unproven.**

## **CMM-400.5: Procedure (CPT®) Codes**

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required. Pre- authorization requirements vary by individual payor.

<b>CPT®</b>	<b>Code Description/Definition</b>
<b>01991</b>	Anesthesia for Diagnostic or Therapeutic Nerve Blocks and Injections (When Block or Injection is Performed by a Different Physician or Other Qualified Health Care Professional); Other Than the Prone Position
<b>01992</b>	Anesthesia for Diagnostic or Therapeutic Nerve Blocks and Injections (When Block or Injection is Performed by a Different Physician or Other Qualified Health Care Professional); Prone Position
<b>01935</b>	Anesthesia for Percutaneous Image Guided Procedures on the Spine and Spinal Cord; Diagnostic
<b>01936</b>	Anesthesia for Percutaneous Image Guided Procedures on the Spine and Spinal Cord; Therapeutic
<b>CPT®</b>	<b>Code Considered Experimental, Investigational, or Unproven</b>
<b>00640</b>	Anesthesia for Manipulation of the Spine or for Closed Procedures on the Cervical, Thoracic or Lumbar Spine

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the individual payor (health insurance company, etc.) and is based on the member/patient/client/beneficiary's policy or benefit entitlement structure as well as any third party payor guidelines and/or claims processing rules. Providers are strongly urged to contact each payor for individual requirements if they have not already done so.

## **CMM-400.6: References**

1. Abram S, Francis M. Hazards of sedation for interventional pain procedures: the anesthesia patient safety foundation newsletter. 27(2):29-31.
2. American American Society of Anesthesiologists, Anesthesiology 2018; 128:437-79.
3. American Society of Anesthesiologists. Continuum of depth of sedation: definition of general anesthesia and levels of sedation/analgesia. Committee of origin: Quality Management and Departmental Administration. Last amended October 15, 2014.
4. American Society of Anesthesiologists. Position on monitored anesthesia care. Last amended October 16, 2013.
5. American Society of Anesthesiologists. Practice Guidelines for Chronic Pain Management: An Updated Report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. April 2010.
6. American Society of Anesthesiologists. Practice guidelines for moderate procedural sedation and analgesia 2018: a report by the American Society of Anesthesiologists Task Force on Moderate Procedural Sedation and Analgesia, the American Association of Oral and Maxillofacial Surgeons, American College of Radiology, American Dental Association, American Society of Dentist Anesthesiologists, and Society of Interventional Radiology. March 2018.
7. American Society of Anesthesiologists. Standards and Guidelines Distinguishing Monitored Anesthesia Care from Moderate Sedation/Analgesia. October 17 2018.
8. American Society of Anesthesiologists. Statement on anesthetic care during interventional pain procedures for adults. Last amended October 26, 2016.
9. American Society of Anesthesiologists. Statement on regional anesthesia. Last amended October 25, 2017.
10. American Society of Anesthesiologists. Practice guidelines for management of the difficult airway: An updated report by the American Society of Anesthesiologists Task Force on Management of the Difficult Airway. 2013; 118:251-70.

11. Early DS, Lightdale JR, Vargo JJ, et al. Guidelines for sedation and anesthesia in GI endoscopy. *Gastrointestinal Endoscopy*. 2018;98(2):327-337. doi:10.1016/j.gie.2017.07.018.
12. Ene, H. Hyperkinetic movement disorders(including dystonias, choreas). *PM&R Knowledge Now*. 9/20/14
13. Spine Intervention Society FactFinders For Patient Safety. Conscious Sedation. February 2018.
14. Vargo JJ, Delege MH, Feld AD, et al. Multisociety sedation curriculum for gastrointestinal endoscopy. *The American Journal of Gastroenterology*. 2012. doi:10.1038/ajg.2012.112.