

Cigna Medical Coverage Policies – Musculoskeletal Shoulder Surgery – Arthroscopic and Open Procedures

Effective August 15, 2020



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by eviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2020 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

©Copyright 2020 eviCore healthcare

CMM-315: Shoulder Surgery-Arthroscopic and Open Procedures

Definitions

- **Rotator cuff tears** result when there is a disruption of the tendon(s) of the rotator cuff muscles which attach the humerus to the scapula and are important in shoulder movements and maintaining glenohumeral joint stability. The supraspinatus tendon is most commonly involved, but the infraspinatus, teres minor, and subscapularis tendons can also be torn.
 - ◆ Defining whether a rotator cuff tear is acute has relevance to treatment. In evaluating patients, the surgeon should attempt to properly identify patients with acute tears as opposed to patients with pre-existing chronic tears that become symptomatic after an injury event. A discrete traumatic event is more suggestive of acute tear. Physical examination findings including supraspinatus and infraspinatus muscle atrophy, as well as internal and external rotation lag signs, may be indicative of larger and more chronic rotator cuff tears.
 - ◆ Evaluation of rotator cuff muscle quality with CT or MRI is an important consideration. Chronic and larger tears are associated with muscle atrophy and fatty replacement, both of which correlate with inferior functional outcome after rotator cuff repair. It is thought that early repair of acute rotator cuff tears might mitigate the development of chronic tendon and muscle pathology and improve functional outcomes.
 - ◆ Classification of rotator cuff tears (based upon surgical findings):
 - Partial-thickness tears, also called incomplete tears (Ellman):
 - Grade 1: < 3 mm deep (< 25% thickness)
 - Grade 2: 3–6 mm in depth but not exceeding 50% of the tendon thickness
 - Grade 3: > 6 mm deep (> 50% thickness)
 - Full-thickness tears, also called complete tears (Cofield):
 - Small: < 1 cm
 - Medium: 1-3 cm
 - Large: 3-5 cm
 - Massive: > 5 cm
- **Impingement syndrome** commonly results from friction, abrasion, and inflammation of the rotator cuff and the long head of the biceps tendon with the subacromial arch (anterior lip of the acromion, coracoacromial ligament, and acromioclavicular joint) from acute trauma, repetitive use or degenerative changes.
- **Distal clavicle excision** is the removal of the end of the clavicle at the acromioclavicular (AC) joint. The superior AC ligament remains intact so that the joint remains stable.
- **Acromioplasty** is the removal of bone from the acromion and partial resection of the coracoacromial ligament.
- **Subacromial decompression** is the removal of bone or other abnormality to enlarge the space between the rotator cuff musculature and the acromion.

- **Labral tears** result when the glenoid labrum becomes injured or torn. Tears are typically classified by the position of the tear in relation to the glenoid.
 - ◆ **Bankart tear** is a tear in the labrum located in the front, lower (anterior, inferior) part of the glenoid. This type of tear occurs most commonly during a shoulder dislocation and makes the shoulder more prone to recurrent dislocations.
 - ◆ **SLAP tear (Superior Labral, Anterior and Posterior tear)** is a tear in the labrum that covers the top part of the glenoid from the front to back. A SLAP tear occurs at the point where the long head of the biceps tendon attaches. This type of tear occurs most commonly during falls on an outstretched arm.
- **Shoulder dislocation** is defined as the complete loss of the humeral articulation with the glenoid fossa, usually as a result of acute trauma.
- **Shoulder subluxation** is defined as a partial loss of humeral articulation with the glenoid fossa (incomplete or partial dislocation) usually as a result of repetitive trauma to the degree that symptoms are produced.
- **Shoulder instability and/or laxity** is defined as a partial loss of the glenohumeral articulation of which there are two categories:
 - ◆ Post-traumatic shoulder instability includes an individual with a previous injury that has stretched or torn the ligaments of the shoulder.
 - ◆ Atraumatic instability and/or laxity includes an individual with generalized looseness of the joints “double-jointed” or “multi-directional instability” usually representing a type of congenital ligamentous laxity.
- **Adhesive capsulitis** is a condition of the shoulder characterized by stiffness, loss of motion (contracture), and pain due to scarring in and/or around the shoulder joint. Conditions that have been suggested to predispose an individual to adhesive capsulitis include trauma, surgery to the shoulder, inflammatory diseases, diabetes, hyperthyroidism, dyslipidemia. Often called frozen shoulder, adhesive capsulitis is clinically divided into classes:
 - ◆ Primary adhesive capsulitis is characterized by a significant limitation of both active and passive motions on the shoulder; individuals are typically unable to recall a possible cause of the condition (idiopathic adhesive capsulitis).
 - ◆ Secondary adhesive capsulitis is characterized by a trauma or a possible cause prior to the onset of the symptoms, such as fracture of the humerus, rotator cuff repair, shoulder girdle injury/surgery, or prolonged immobilization.
- **Non-surgical management**, with regard to the treatment of shoulder pain, is defined as any provider-directed non-surgical treatment that has been demonstrated in the scientific literature to be efficacious and/or is considered reasonable care in the treatment of shoulder pain. The types of treatment involved can include, but are not limited to: relative rest/activity modification, supervised physiotherapy modalities and supervised therapeutic exercises, oral prescription and non-prescription medications, assistive devices (e.g., sling, splint, brace), and/or injections (i.e., steroid).

General Guidelines

- The determination of medical necessity for the performance of shoulder surgery is always made on a case-by-case basis.
- Refer to **MS-19: Shoulder** for advanced imaging indications for conditions of the shoulder.
- Manipulation of a shoulder joint under general anesthesia is included in all arthroscopic shoulder procedures and is therefore considered incidental to the base procedure requiring medical necessity review.

Indications and Non-Indications

- Shoulder arthroscopic or open procedures may be **considered medically necessary** for individuals when surgery is being performed for fracture, tumor, infection or foreign body that has led to or will likely lead to progressive destruction.

Diagnostic Arthroscopy

- Diagnostic arthroscopy is considered **medically necessary** as a separate procedure when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment) for at least six (6) months in duration
 - ◆ Individual demonstrates **ANY** of the following abnormal, shoulder physical examination findings as compared to the non-involved side:
 - Functionally limited range of motion (active or passive)
 - Measurable loss in strength
 - Positive Neer Impingement Test or Hawkins-Kennedy Impingement Test
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration
 - ◆ Advanced diagnostic imaging study (e.g., MRI, CT) is inconclusive for internal derangement/pathology
 - ◆ Other potential pathological conditions including, but not limited to: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain and advanced glenohumeral osteoarthritis have been excluded.
- Diagnostic arthroscopy is considered **not medically necessary** for all other indications.

Loose Body/Foreign Body Removal

- Loose body or foreign body removal is considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Mechanical symptoms including painful locking, clicking, catching, or popping
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration, except when the loose body or foreign body has caused an acute restriction of shoulder joint range of motion (i.e., locking)
 - ◆ Advanced diagnostic imaging study (e.g., MRI, CT) is conclusive for the presence of a loose body or foreign body within the shoulder joint
 - ◆ Other potential pathological conditions including, but not limited to: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, and advanced glenohumeral osteoarthritis have been excluded
- Loose body or foreign body removal is considered **not medically necessary** for all other indications.

Synovectomy

- Synovectomy (partial or complete) is considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Individual demonstrates functionally limited range of motion (active or passive) on physical examination as compared to the non-involved side
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration
 - ◆ Advanced diagnostic imaging study (e.g., MRI, CT) demonstrates underlying pathology consistent with the individual's reported medical condition (e.g., synovitis, joint effusion) which correlates with the individual's reported symptoms and physical exam findings
 - ◆ Presence of any **ONE** of the following:
 - Inflammatory arthritis (i.e., rheumatoid arthritis, gout, pseudogout, psoriatic arthritis)
 - Pigmented villonodular synovitis (PVNS)
 - Synovial chondromatosis
 - Lyme synovitis
 - Hemophilia
 - Hemochromatosis
 - Non-specific synovitis (including proliferative synovitis, post-operative synovitis as a sequela from a shoulder replacement, etc.)
 - Recurrent hemarthrosis (i.e., secondary to sickle cell anemia, bleeding diathesis, etc.)

- ◆ Other potential pathological conditions including, but not limited to: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, and advanced glenohumeral osteoarthritis have been excluded
- Synovectomy is considered **not medically necessary** for all other indications.

Debridement

- Debridement (limited or extensive) is considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Individual demonstrates the following on physical examination when compared to the non-involved side:
 - **EITHER** of the following:
 - Functionally limited range of motion
 - Measurable loss of strength
 - **ONE OR MORE** of the following positive orthopedic tests/signs:
 - Drop Arm Test
 - Painful Arc Test
 - Jobe or Empty Can Test
 - External Rotation Lag Sign
 - Lift-Off Test
 - Belly-Press Test
 - Cross Body Adduction Test
 - Resisted AC Joint Extension Test
 - Neer Impingement Test
 - Hawkins-Kennedy Impingement Test
 - O'Brien's Test
 - Biceps Load Test
 - Clunk Test
 - Anterior Slide Test
 - Compression Rotation Test
 - Speed's Test
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration
 - ◆ Advanced diagnostic imaging study (e.g., MRI, CT) demonstrates underlying pathology which correlates with the individual's reported symptoms and physical exam findings
 - ◆ Other potential pathological conditions including, but not limited to: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, and advanced glenohumeral osteoarthritis have been excluded
- Debridement is considered **not medically necessary** for all other indications.

Rotator Cuff Repair

- Rotator cuff repair **is considered medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., loss of shoulder function to the extent which interferes with ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Individual demonstrates the following on physical examination when compared to the non-involved side:
 - **EITHER** of the following :
 - Functionally limited range of motion
 - Measurable loss of strength of the rotator cuff musculature
 - **ONE or MORE** of the following positive orthopedic tests/signs:
 - Drop Arm Test
 - Painful Arc Test
 - Jobe or Empty Can Test
 - External Rotation Lag Sign (Dropping Sign)
 - Internal Rotation Lag Sign
 - Lift-Off Test
 - Bear Hug Test
 - Belly-Press Test (Napoleon)
 - Neer Impingement Test
 - Hawkins-Kennedy Impingement Test
 - Hornblower Test (Patte)
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration, except for an individual who suffers a discrete traumatic event that results in an acute full-thickness rotator cuff tear AND associated function-limiting pain
 - Note: The failure of provider-directed non-surgical management for at least three (3) months in duration is required in the presence of these advanced diagnostic imaging (e.g., MRI, CT) findings of fatty infiltration and/or muscle atrophy, regardless of whether a discrete traumatic event occurred.
 - ◆ Advanced diagnostic imaging (e.g., MRI, CT) demonstrates a Grade 2 or 3 partial-thickness rotator cuff tear (Ellman classification) or a full-thickness rotator cuff tear (Cofield classification) that correlates with the individual's reported symptoms and physical exam findings
 - ◆ Other potential pathological conditions including but not limited to fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain and advanced glenohumeral osteoarthritis have been excluded.
- Rotator cuff repair is considered not **medically necessary** for all other indications.

Distal Clavicle Excision/Subacromial Decompression/ Acromioplasty

- Distal clavicle excision is considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., documented loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Individual demonstrates localized tenderness to palpation of the acromioclavicular (AC) joint and **ONE or MORE** of the following positive orthopedic tests on physical examination when compared to the non-involved side:
 - Cross Body Adduction Test
 - Resisted AC Joint Extension Test
 - Neer Impingement Test
 - Hawkins-Kennedy Impingement Test
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration
 - ◆ Plain radiographs demonstrate findings consistent with pathology in the subacromial space and/or at the AC joint
 - ◆ Advanced diagnostic imaging study (e.g., MRI, CT) demonstrates underlying pathology (e.g., AC joint arthritis, impingement, etc.) which correlates with the individual's reported symptoms and physical exam findings
 - Note: Advanced diagnostic imaging is not required for isolated distal clavicle excision when not associated with subacromial decompression/acromioplasty surgery.
 - ◆ Other potential pathological conditions including, but not limited to: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, and advanced glenohumeral osteoarthritis have been excluded
- Subacromial decompression/acromioplasty is considered **medically necessary** as an “add-on procedure” only when performed with other medically necessary primary shoulder surgical procedures **AND ALL** of the above criteria have been met with the exception of localized tenderness to palpation of the acromioclavicular joint
- Distal clavicle excision/subacromial decompression/acromioplasty is considered **not medically necessary** for all other indications.

Labral Repair (e.g., SLAP)/Biceps Tenodesis

- Labral repair (e.g., SLAP)/biceps tenodesis is considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Individual demonstrates **BOTH** of the following on physical examination when compared to the non-involved side:
 - Minimally limited or full shoulder range of motion
 - **ONE or MORE** of the following positive orthopedic tests
 - O'Brien's Test
 - Biceps Load Test
 - Clunk Test
 - Anterior Slide Test
 - Compression Rotation Test
 - Speed's Test
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration
 - ◆ Advanced diagnostic imaging study (e.g., MRI, CT) demonstrates labral (e.g., SLAP) biceps tendon pathology and correlates with the individual's reported symptoms and physical exam findings
 - ◆ Other potential pathological conditions including but not limited to fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain and advanced glenohumeral osteoarthritis have been excluded.
- Labral repair/biceps tenodesis is considered **not medically necessary** for all other indications.

Shoulder Instability and/or Laxity

- Arthroscopic or open surgical procedures for shoulder instability and/or laxity are considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ Documented history of "post-traumatic" or "atraumatic" instability and/or laxity that has resulted in function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment)
 - ◆ Individual demonstrates **ONE OR MORE** of the following positive orthopedic tests on physical examination when compared to the non-involved side:
 - Anterior or Posterior Apprehension Test
 - Sulcus Sign
 - Load and Shift Test
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration that includes shoulder stabilization/strengthening exercises, except when **EITHER** of the following criteria are met in an acute traumatic injury setting:
 - Irreducible shoulder dislocation
 - Anterior shoulder instability in competitive contact or collision athletes

- ◆ Advanced diagnostic imaging study (e.g., MRI, CT) demonstrates labral tear (e.g., Bankart, capsular tear) and correlates with the individual's reported symptoms and physical exam findings
 - ◆ Other potential pathological conditions including, but not limited to: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, and advanced glenohumeral osteoarthritis have been excluded
- Arthroscopic or open surgical procedures for shoulder instability and/or laxity are considered **not medically necessary** for all other indications.

Arthroscopic Capsular Release/Lysis of Adhesions

- Arthroscopic capsular release/lysis of adhesions for an individual with documented chronic refractory adhesive capsulitis/arthrofibrosis which has resulted from disease, injury or surgery is considered **medically necessary** when **ALL** of the following criteria have been met:
- ◆ Function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age appropriate activities of daily living and/or demands of employment) for at least six (6) months in duration
 - ◆ Individual demonstrates functionally limited and painful global loss of active and passive range of motion of at least 50% when compared to the non-involved side
 - ◆ Failure of provider-directed non-surgical management for at least three (3) months in duration, including a combination of anti-inflammatory medication, cortisone injection, and a trial of physical therapy (i.e., active exercise and manual therapy designed to increase joint mobility and range of motion)
 - ◆ Other potential diagnostic conditions including but not limited to fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain and advanced glenohumeral osteoarthritis have been excluded.
- Arthroscopic capsular release/lysis of adhesions is considered **not medically necessary** for all other indications.

Experimental, Investigational, or Unproven

- Based on lack of scientific safety and efficacy, each of the following is considered **experimental, investigational or unproven**:
- ◆ In-office diagnostic arthroscopy (e.g., Mi-Eye™, VisionScope®)
 - ◆ Superior capsular reconstruction using any type of graft (e.g., autograft, allograft, synthetic)
 - ◆ Arthroscopic or open coracoplasty/subcoracoid decompression for the treatment of subcoracoid impingement

Procedure (CPT®) Codes

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
23000	Removal of subdeltoid calcareous deposits, open
23020	Capsular contracture release (e.g. Sever type procedure)
23030	Incision and drainage, shoulder area; deep abscess or hematoma
23031	Incision and drainage, shoulder area; infected bursa
23035	Incision, bone cortex (eg, osteomyelitis or bone abscess), shoulder area
23040	Arthrotomy, glenohumeral joint, including exploration, drainage, or removal of foreign body
23044	Arthrotomy, acromioclavicular, sternoclavicular joint, including exploration, drainage, or removal of foreign body
23065	Biopsy, soft tissue of shoulder area; superficial
23066	Biopsy, soft tissue of shoulder area; deep
23071	Excision, tumor, soft tissue of shoulder area, subcutaneous; 3 cm or greater
23073	Excision, tumor, soft tissue of shoulder area, subfascial (e.g. intramuscular); 5 cm or greater
23075	Excision, tumor, soft tissue of shoulder area, subcutaneous; less than 3 cm
23076	Excision, tumor, soft tissue of shoulder area, subfascial (e.g. intramuscular); less than 5 cm
23077	Radical resection of tumor (e.g. sarcoma), soft tissue of shoulder area; less than 5 cm
23078	Radical resection of tumor (e.g. sarcoma), soft tissue of shoulder area; 5 cm or greater
23100	Arthrotomy, glenohumeral joint, including biopsy
23101	Arthrotomy, acromioclavicular joint or sternoclavicular joint, including biopsy and/or excision of torn cartilage
23105	Arthrotomy; glenohumeral joint, with synovectomy, with or without biopsy
23106	Arthrotomy;sternoclavicular joint, with synovectomy, with or without biopsy
23107	Arthrotomy, glenohumeral joint, with joint exploration, with or without removal of loose or foreign body
23120	Claviculectomy; partial
23125	Claviculectomy; total
23130	Acromioplasty or acromionectomy, partial, with or without coracoacromial ligament release
23140	Excision or curettage of bone cyst or benign tumor of clavicle or scapula
23145	Excision or curettage of bone cyst or benign tumor of clavicle or scapula; with autograft (includes obtaining graft)
23146	Excision or curettage of bone cyst or benign tumor of clavicle or scapula; with allograft
23150	Excision or curettage of bone cyst or benign tumor of proximal humerus
23155	Excision or curettage of bone cyst or benign tumor of proximal humerus; with autograft (includes obtaining graft)
23156	Excision or curettage of bone cyst or benign tumor of proximal humerus; with allograft.
23170	Sequestrectomy (e.g. for osteomyelitis or bone abscess), clavicle
23172	Sequestrectomy (e.g. for osteomyelitis or bone abscess), scapula
23174	Sequestrectomy (e.g. for osteomyelitis or bone abscess), humeral head to surgical neck
23180	Partial excision (craterization, saucerization, or diaphysectomy) bone (e.g. osteomyelitis), clavicle
23182	Partial excision (craterization, saucerization, or diaphysectomy) bone (e.g. osteomyelitis), scapula

23184	Partial excision (craterization, saucerization, or diaphysectomy) bone (e.g. osteomyelitis), proximal humerus
23190	Ostectomy of scapula, partial (eg, superior medial angle)
23195	Resection, humeral head
23200	Radical resection of tumor; clavicle
23210	Radical resection of tumor; scapula
23220	Radical resection of tumor, proximal humerus
23395	Muscle transfer, any type, shoulder or upper arm; single
23397	Muscle transfer, any type, shoulder or upper arm; multiple
23405	Tenotomy, shoulder area; single tendon
23406	Tenotomy, shoulder area; multiple tendons through same incision
23410	Repair of ruptured musculotendinous cuff (e.g. rotator cuff) open; acute
23412	Repair of ruptured musculotendinous cuff (e.g. rotator cuff) open; chronic
23415	Coracoacromial ligament release, with or without acromioplasty
23420	Reconstruction of complete shoulder (rotator) cuff avulsion, chronic (includes acromioplasty)
23430	Tenodesis of long tendon of biceps
23440	Resection or transplantation of long tendon of biceps
23450	Capsulorrhaphy, anterior; Putti-Platt procedure or Magnuson type operation
23455	Capsulorrhaphy, anterior; with labral repair (e.g. Bankart procedure)
23460	Capsulorrhaphy, anterior, any type; with bone block
23462	Capsulorrhaphy, anterior, any type; with coracoid process transfer
23465	Capsulorrhaphy, glenohumeral joint, posterior, with or without bone block
23466	Capsulorrhaphy, glenohumeral joint, any type multi-directional instability
23480	Osteotomy, clavicle, with or without internal fixation
23485	Osteotomy, clavicle, with or without internal fixation; with bone graft for nonunion or malunion (includes obtaining graft and/or necessary fixation)
23490	Prophylactic treatment (nailing, pinning, plating or wiring) with or without methylmethacrylate; clavicle
23491	Prophylactic treatment (nailing, pinning, plating or wiring) with or without methylmethacrylate; proximal humerus
29805	Arthroscopy, shoulder, diagnostic, with or without synovial biopsy (separate procedure)
29806	Arthroscopy, shoulder, surgical; capsulorrhaphy
29807	Arthroscopy, shoulder, surgical; repair of SLAP lesion
29819	Arthroscopy, shoulder, surgical; with removal of loose body or foreign body
29820	Arthroscopy, shoulder, surgical; synovectomy, partial
29821	Arthroscopy, shoulder, surgical; synovectomy, complete
29822	Arthroscopy, shoulder, surgical; debridement, limited
29823	Arthroscopy, shoulder, surgical; debridement, extensive
29824	Arthroscopy, shoulder, surgical; distal claviclectomy including distal articular surface (Mumford procedure)
29825	Arthroscopy, shoulder, surgical; with lysis and resection of adhesions, with or without manipulation
29826	Arthroscopy, shoulder, surgical; decompression of subacromial space with partial acromioplasty, with coracoacromial ligament (i.e. arch) release when performed (List separately in addition to code for primary procedure)
29827	Arthroscopy, shoulder, surgical; with rotator cuff repair
29828	Arthroscopy, shoulder, surgical; biceps tenodesis

This list may not be all inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.

Shoulder Surgery-Arthroscopic and Open Procedures

References

1. AAOS Clinical Practice Guidelines Unit. Optimizing The Management of Rotator Cuff Problems: Guideline and Evidence Report. 2010 Dec 4. V1.1_033011:i-293.
2. Arciero RA, Wheeler JH, Ryan JB, et al. Arthroscopic Bankart Repair Versus Nonoperative Treatment for Acute, Initial Anterior Shoulder Dislocations. *Am J Sports Med.* 1994;22(5):589-594.
3. Audenaert E, Nuffel JV, Schepens A, Verhelst M, Verdonk R. Reconstruction of massive rotator cuff lesions with a synthetic interposition graft: a prospective study of 41 patients. *Knee Surgery, Sports Traumatology, Arthroscopy.* 2005;14(4):360-364. doi:10.1007/s00167-005-0689-7.
4. Boileau P, Baque F, Valerio L et al. Isolated Arthroscopic Biceps Tenotomy or Tenodesis Improves Symptoms in Patients with Massive Irreparable Rotator Cuff Tears. *J Bone Joint Surg Am.* 2007; 89(4):747757.
5. Boorman RS, More KD, Hollinshead RM, et al. What happens to patients when we do not repair their cuff tears? Five-year rotator cuff quality-of-life index outcomes following nonoperative treatment of patients with full-thickness rotator cuff tears. *J Shoulder Elbow Surg.* 2018;27:444-448. doi:10.1016/j.jse.2017.10.009.
6. Budoff J, Nirschl R, Guidi E. Current Concepts Review-Debridement of Partial-Thickness Tears of the Rotator Cuff without Acromioplasty. Long-term follow-up and review of the literature. *J Bone Joint Surg Am.* 1998; 80(5):733-748.
7. Carver TJ, Kraeutler MJ, Smith JR, Bravman JT, Mccarty EC. Nonarthroplasty Surgical Treatment Options for Massive, Irreparable Rotator Cuff Tears. *Orthopaedic Journal of Sports Medicine.* 2018;6(11):232596711880538. doi:10.1177/2325967118805385.
8. Choi L. Overuse injuries. In: DeLee J, et al. *DeLee and Drez's Orthopaedic Sports Medicine.* 3 ed. Philadelphia, Pa.: Saunders Elsevier; 2009.
9. Cofield RH. Subscapular muscle transposition for repair of chronic rotator cuff tears. *Surg Gynecol Obstet.* 1982;154(5):667-672.
10. Denard PJ, Brady PC, Adams CR, et al. Preliminary results of arthroscopic superior capsule reconstruction with dermal allograft. *Arthroscopy.* 2018;34(1): 93-99.
11. DeOrio JK, Cofield RH. Results of a second attempt at surgical repair of a failed initial rotator-cuff repair. *J Bone Joint Surg Am.* 1984;66(4):563-567.
12. Dunn WR, Kuhn JE, Sanders R, et al. 2013 Neer Award: predictors of failure of nonoperative treatment of chronic, symptomatic, full-thickness rotator cuff tears. *J Shoulder Elbow Surg.* 2016;25:1303-1311.
13. Forsythe B, Agarwalla A, Puzitiello RN, Sumner S, Romeo AA, Mascarenhas R. The Timing of Injections Prior to Arthroscopic Rotator Cuff Repair Impacts the Risk of Surgical Site Infection. *The Journal of Bone and Joint Surgery.* 2019;101(8):682-687. doi:10.2106/jbjs.18.00631.
14. Galatz LM, Ball C, Teefey S, et al. The outcome and repair integrity of completely arthroscopically repaired large and massive rotator cuff tears. *J Bone Joint Surg Am.* 2004; 86(2):219-224.
15. Garofalo R, Conti M, Massazza G, Cesari E, Vinci E, Castagna A. Subcoracoid impingement syndrome: a painful shoulder condition related to different pathologic factors. *Musculoskelet Surg.* 2011 Jul;95 Suppl 1:S25-9.
16. Gartsman G, Taverna E. The incidence of glenohumeral joint abnormalities associated with complete, reparable rotator cuff tears. *Arthroscopy.* 1997;12:575-579.
17. Godeneche A, Elia F, Kempf J, et al. Fatty infiltration of stage 1 or higher significantly compromises long-term healing of supraspinatus repairs. *J Shoulder Elbow Surg.* 2017;26:1818-1825. doi: 10.1016/j.jse.2017.03.024.
18. Gupta AK, Hug K, Berkoff DJ, et al. Dermal Tissue Allograft for the Repair of Massive Irreparable Rotator Cuff Tears. *The American Journal of Sports Medicine.* 2012;40(1):141-147. doi:10.1177/0363546511422795.
19. Herbery-Davies J, Teefey SA, Steger-May K, et al. Progression of fatty muscle degeneration in atraumatic rotator cuff tears. *J Bone Joint Surg Am.* 2017;99:832-9. doi:10.2106/JBJS.16.00030.
20. Hippensteel KJ, Brophy R, Smith MV, et al. A comprehensive review of physical examination tests of the cervical spine, scapula, and rotator cuff. *J Am Acad Orthop Surg.* 2019;27: 385-94. doi: 10.5435/JAAOS-D-17-00090.
21. Hirahara AM, Andersen WJ, Panero AJ. Superior capsular reconstruction: clinical outcomes after minimum 2-year follow-up. *Am J Orthop.* 2017: 266-78.

22. Hovelius L, Olofsson A, Sandström B. Nonoperative treatment of primary anterior shoulder dislocation in patients forty years of age and younger. a prospective twenty-five-year follow-up. *J Bone Joint Surg Am.* 2008;90(5):945-952.
23. Hovis W, Dean M, Mallon W, Hawkins R. Posterior instability of the shoulder with secondary impingement in elite golfers. *Am J Sports Med.* 2002;30(6):886-890.
24. Keener JD, Galatz LM, Teefey SA, et al. A Prospective Evaluation of Survivorship of Asymptomatic Degenerative Rotator Cuff Tears. *J Bone Joint Surg Am.* 2015;97:89-98.
25. Keener JD, Patterson BM, Orvets N, Chamberlain AM. Degenerative Rotator Cuff Tears. *Journal of the American Academy of Orthopaedic Surgeons.* 2019;27(5):156-165. doi:10.5435/jaaos-d-17-00480.
26. Khazzam M, Kane SM, Smith MJ. Open Shoulder Stabilization Using bone block technique for treatment of chronic glenohumeral instability associated with glenoid deficiency. *Am J Orthop (Belle Mead NJ).* 2009;38(7):329-335
27. Kim H, Teefey S, Zelig A, et al. Shoulder strength in asymptomatic individuals with intact compared with torn rotator cuffs. *J Bone Joint Surg Am.* 2009; 91(2):289-296.
28. Kim SJ, Choi YR, Jung M, Yoon Y, Chun YM. Concomitant coracoplasty during arthroscopic subscapularis repair does not yield better clinical outcomes and structural integrity. *Knee Surg Sports Traumatol Arthrosc.* 2018 Jan;26(1):56-62.
29. Kragh JF Jr, Doukas WC, Basamania CJ. Primary coracoid impingement syndrome. *Am J Orthop (Belle Mead NJ).* 2004 May;33(5):229-32; discussion 232.
30. Kukkonen J, Joukainen A, Lehtinen J, et al. Treatment of Nontraumatic Rotator Cuff Tears: A Randomized Control Trial with Two Years of Clinical and Imaging Follow-up. *J Bone Joint Surg Am.* 2015;97:1729-37.
31. Lee SJ, Min YK. Can inadequate acromiohumeral distance improvement and poor posterior remnant tissue be the predictive factors of re-tear? Preliminary outcomes of arthroscopic superior capsular reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(7): 2205-13.
32. Leroux, TS, Saltzman BM, Meyer M, et al. The Influence of Evidence-Based Surgical Indications and Techniques on Failure Rates After Arthroscopic Shoulder Stabilization in the Contact or Collision Athlete with Anterior Shoulder Instability. *Am J Sports Med.* 2017;45(5):1218-1225.
33. Lewington MR, Ferguson DP, Smith TD, Burks R, Coady C, Wong IH-B. Graft Utilization in the Bridging Reconstruction of Irreparable Rotator Cuff Tears: A Systematic Review. *The American Journal of Sports Medicine.* 2017;45(13):3149-3157. doi:10.1177/0363546517694355.
34. Lim S, Alramadhan H, Kwak J-M, Hong H, Jeon I-H. Graft tears after arthroscopic superior capsule reconstruction (ASCR): pattern of failure and its correlation with clinical outcome. *Archives of Orthopaedic and Trauma Surgery.* 2018;139(2):231-239. doi:10.1007/s00402-018-3025-7.
35. McKee M, Yoo D. The effect of surgery for rotator cuff disease on general health status. Results of a prospective trial. *J Bone Joint Surg Am.* 2000; 82(7):970-979.
36. Mihata T, Lee TQ, Watanabe C, et al. Clinical results of arthroscopic superior capsule reconstruction for irreparable rotator cuff tears. *Arthroscopy.* 2013;29(3): 459-70.
37. Mishra D, Fanton G. Two-year outcome of arthroscopic Bankart repair and electrothermal-assisted capsulorrhaphy for recurrent traumatic anterior shoulder instability. *Arthroscopy.* 2001;17(8):844- 849.
38. Moosmayer S, Gartner AV, Tariq R. The natural course of nonoperatively treated rotator cuff tears: an 8.8 year follow-up of tear anatomy and clinical outcome in 49 patients. *J Shoulder Elbow Surg.* 2017;26:627-634. doi: 10.1016/j.jse.2016.10.002.
39. Neumann JA, Zgonis MH, Rickert KD, et al. Interposition Dermal Matrix Xenografts: A Successful Alternative to Traditional Treatment of Massive Rotator Cuff Tears. *The American Journal of Sports Medicine.* 2017;45(6):1261-1268. doi:10.1177/0363546516683945.
40. Park M, Jun B, Park C, et al. Biomechanical Analysis of a Knotless Transtendon Interimplant Mattress Repair for Partial-Thickness Articular-Sided Rotator Cuff Tears. *Am J Sports Med.* 2009;37(12):2427-2434.
41. Park JY, Lhee SH, Oh KS, Kim NR, Hwang JT. Is arthroscopic coracoplasty necessary in subcoracoid impingement syndrome? *Arthroscopy.* 2012 Dec;28(12):1766-75.
42. Pennington WT, Bartz BA, Pauli JM, et al. Arthroscopic superior capsular reconstruction with acellular dermal allograft for the treatment of massive irreparable rotator cuff tears, short-term clinical outcomes and the radiographic parameter of superior capsular distance. *Arthroscopy.* 2018;34(6): 1764-73.
43. Petrera A, Dwyer T, Tsuji MRS, et al. Outcomes of Arthroscopic Bankart Repair in Collision Versus

- Noncollision Athletes. *Orthopedics*. 2013;36(5):e621-e626.
44. Ranebo MC, Hallgren HCB, Norlin R, Adolfsson LE. Long-term clinical and radiographic outcome of rotator cuff repair with a synthetic interposition graft: a consecutive case series with 17 to 20 years of follow-up. *Journal of Shoulder and Elbow Surgery*. 2018;27(9):1622-1628. doi:10.1016/j.jse.2018.03.011.
 45. Rees J. The pathogenesis and surgical treatment of tears of the rotator cuff. *J Bone Joint Surg Br*. 2008;90-B(7):827-832.
 46. Rhon DI, Boyles RB, Cleland JA. One-year outcome of subacromial corticosteroid injection compared with manual physical therapy for the management of the unilateral shoulder impingement syndrome: a pragmatic randomized trial. *Ann Intern Med*. 2014;161(3):161-169.
 47. Sachs RA, Lin D, Stone ML, et al. Can the Need for Future Surgery for Acute Traumatic Anterior Shoulder Dislocation Be Predicted? *J Bone Joint Surg Am*. 2007;89:1665-74.
 48. Shen P, Lien S, Shen H, et al. Long-term functional outcomes after repair of rotator cuff tears correlated with atrophy of the supraspinatus muscles on magnetic resonance images. *J Shoulder Elbow Surg*. 2008; 17 (1 Suppl): 1S-7S.
 49. Sochacki KR, Mcculloch PC, Lintner DM, Harris JD. Superior Capsular Reconstruction for Massive Rotator Cuff Tear Leads to Significant Improvement in Range of Motion and Clinical Outcomes: A Systematic Review. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*. 2019;35(4):1269-1277. doi:10.1016/j.arthro.2018.10.129.
 50. Suenaga N, Minami A, Kaneda K. Postoperative subcoracoid impingement syndrome in patients with rotator cuff tear. *J Shoulder Elbow Surg*. 2000 Jul-Aug;9(4):275-8.
 51. Suenaga N, Minami A, Kaneda K. Postoperative subcoracoid impingement syndrome in patients with rotator cuff tear. *J Shoulder Elbow Surg*. 2000 Jul-Aug;9(4):275-8.
 52. Streubel PN, Krych AJ, Simone JP, et al. Anterior Glenohumeral Instability: A Pathology-based Surgical Treatment Strategy. *J Am Acad Orthop Surg*. 2014;22(5):283-294.
 53. Ueda Y, Sugaya H, Takahashi N, et al. Rotator Cuff Lesions in Patients with Stiff Shoulders: A Prospective Analysis of 379 Shoulders. *J Bone Joint Surg Am*. 2015;97:1233-7.
 54. Venouziou AI, Kokkalis ZT, Sotereanos DG. Human dermal allograft interposition for the reconstruction of massive irreparable rotator cuff tears. *Am J Orthop*. 2013;42(2):63-70.
 55. Vitale M, Arons R, Hurwitz S, et al. The Rising Incidence of Acromioplasty. *J. Bone Joint Surg*. 2010;92(9):1842-1850.
 56. Werner BC, Brockmeier SF, Miller MD. Etiology, Diagnosis, and Management of Failed SLAP Repair. *J Am Acad Orthop Surg*. 2014;22(9):554-565.
 57. Whittle S, Buchbinder R. In the Clinic: Rotator Cuff Disease. *Ann Intern Med*. 6 Jan 2015. ITC 1-ITC 15.
 58. Work Loss Data Institute. *Shoulder (acute and chronic)*. Corpus Christi (TX): Work Loss Data Institute; 2008.
 59. Wylie JD, Suter T, Potter MQ, et al. Mental Health Has a Stronger Association with Patient-Reported Shoulder Pain and Function Than Tear Size in Patients with Full-Thickness Rotator Cuff Tears. *J Bone Joint Surg Am*. 2016;98:251-6.