

Cigna Medical Coverage Policies – Radiology Peripheral Vascular Disorders Guidelines

Effective July 30, 2020



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by eviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

These guidelines include procedures eviCore does not review for Cigna. Please refer to the [Cigna CPT code list](#) for the current list of high-tech imaging procedures that eviCore reviews for Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2020 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Peripheral Vascular Disease (PVD) Imaging Guidelines

Abbreviations and Glossary for the PVD Imaging Guidelines	3
PVD-1: General Guidelines	4
PVD-2: Screening for Suspected Peripheral Artery Disease/Aneurysmal Disease	9
PVD-3: Cerebrovascular and Carotid Disease	13
PVD-4: Upper Extremity Peripheral Vascular Disease	19
PVD-5: Pulmonary Artery Hypertension	22
PVD-6: Aortic Disorders, Renal Vascular Disorders and Visceral Artery Aneurysms	24
PVD-7: Lower Extremity Peripheral Vascular Disease	38
PVD-8: Imaging for Hemodialysis Access	45
PVD-9: Arteriovenous Malformations (AVMs)	47
PVD-10: This section intentionally left blank	48
PVD-11: Venous Imaging – General Information	49
PVD-12: Acute Limb Swelling	55
PVD-13: Chronic Limb Swelling Due to Chronic Deep Venous Thrombosis/May Thurner's Syndrome	58
PVD-14: Chronic Limb Swelling Due to Venous Insufficiency/ Venous Stasis Changes/ Varicose Veins	61
PVD-15: Venous Stasis Ulceration	65
PVD-16: IVC Filters	67
PVD-17: Post Iliac Vein Stent/Angioplasty	69

Abbreviations and Glossary for the PVD Imaging Guidelines

(See also: [Cardiac Imaging Guidelines Glossary](#))

AAA	abdominal aortic aneurysm
ABI	Ankle brachial index: a noninvasive, non-imaging test for arterial insufficiency – (see toe-brachial index below). This testing can also be done after exercise if resting results are normal.
Claudication	or Intermittent claudication : usually a painful cramping sensation of the legs with walking or severe leg fatigue
CTA	computed tomography angiography
CTV	computed tomography venography
DLCO	diffusion capacity : defined as the volume of carbon monoxide transferred into the blood per minute per mmHg of carbon monoxide partial pressure
DVT	deep venous thrombosis
ECG	electrocardiogram
ENT	Ears, Nose, Throat
HbA1C	hemoglobin A1C : test used to determine blood sugar control for individuals with diabetes
MRA	magnetic resonance angiography
MRV	magnetic resonance venography
PAD	peripheral artery disease
PAH	pulmonary artery hypertension
PFT	pulmonary function tests
PVD	peripheral vascular disease
SVC	superior vena cava
TIA	transient ischemic attack
TTE	transthoracic echocardiogram
Toe-Brachial Index	useful in individuals with ABI above the normal range due to non-compressible posterior tibial or dorsalis pedis arteries
V/Q Scan	ventilation and perfusion scan

PVD-1: General Guidelines

PVD-1.1: General Information	5
PVD-1.2: Procedure Coding	6
PVD-1.3: General Guidelines – Imaging	8

PVD-1.1: General Information

- A current clinical evaluation (within 60 days), including medical treatments, are required prior to considering advanced imaging, which includes:
 - ◆ Relevant history and physical examination including:
 - Palpation of pulses
 - Evaluation of lower extremities for presence of non-healing wounds or gangrene
 - Associated skin changes such as thickened nails, absence of hair in the feet or calves, cool extremities
 - Evaluation for the presence of arterial bruits
 - Appropriate laboratory studies
 - Non-advanced imaging modalities, such as recent ABIs (within 60 days) after symptoms started or worsened
 - ◆ Other meaningful contact (telephone call, electronic mail or messaging) by an established individual can substitute for a face-to-face clinical evaluation.
- Risk factors for vascular disease include:
 - ◆ Diabetes
 - ◆ Cigarette smoking
 - ◆ Hypertension
 - ◆ Hyperlipidemia
 - ◆ Age >50, with at least one risk factor, are considered “at risk” for vascular disease.
 - ◆ See **PV-17: Impotence/Erectile Dysfunction** in the Pelvis Imaging Guidelines.
- Signs and symptoms of peripheral arterial disease:
 - ◆ Claudication (Cramping pain in the legs, most notably back of the calves but can involve hips or thighs, after walking which is relieved with rest but recurs at a predictable distance)
 - Symptoms that are not consistent with claudication include:
 - Generalized leg pain
 - Nocturnal cramps
 - Pain that is not easily relieved after a few minutes of rest
 - Burning pain in feet
 - ◆ Critical limb ischemia
 - Rest pain: Pain in the foot (not leg) at rest, particularly at night when the leg is elevated. Pain is relieved by dangling the leg off the bed or moving to an upright position
 - Non healing wounds. Wounds present for >2 weeks with little to no evidence of healing
 - ◆ Erectile dysfunction can be associated with vascular disease
- Claudication and critical limb ischemia have different natural histories. Claudication generally follows a benign indolent course. 70% of individuals with claudication will have the same symptoms after five years with no progression. Critical limb ischemia, on the other hand, is associated with a high rate of limb loss (25%) and death (35%) one year after presentation

- Simultaneous venous and arterial systems evaluation are unusual but are occasionally needed.
- Post angioplasty/reconstruction: follow-up imaging is principally guided by symptoms. See
 - ◆ **PVD-6.8: Post Aortic Endovascular/Open Surgery Surveillance Studies**
 - ◆ **PVD-7.3: Post-Procedure Surveillance Studies**

PVD-1.2: Procedure Coding

Non-Invasive Physiologic Studies of Extremity Arteries	CPT®
<ul style="list-style-type: none"> ➤ Limited bilateral noninvasive physiologic studies of upper or lower extremity arteries. ➤ Non-invasive physiologic studies of upper or lower extremity arteries, single level, bilateral (e.g., ankle/brachial indices, Doppler waveform analysis, volume plethysmography, transcutaneous oxygen tension measurement). 	93922
<ul style="list-style-type: none"> ➤ Complete bilateral noninvasive physiologic studies of upper or lower extremity arteries, 3 or more levels. ➤ Non-invasive physiologic studies of upper or lower extremity arteries, multiple levels or with provocative functional maneuvers, complete bilateral study (e.g., segmental blood pressure measurements, segmental Doppler waveform analysis, segmental volume plethysmography, segmental transcutaneous oxygen tension measurements, measurements with postural provocative tests, measurements with reactive hyperemia). 	93923

- CPT® 93922 and CPT® 93923 can be requested and reported only once for the upper extremities and once for the lower extremities.
- CPT® 93922 and CPT® 93923 should not be ordered on the same request nor billed together for the same date of service.
- CPT® 93924 and CPT® 93922 and/or CPT® 93923 should not be ordered on the same request and should not be billed together for the same date of service.
- ABI studies performed with handheld dopplers, where there is no hard copy output for evaluation of bidirectional blood flow, are not reportable by these codes.

Non-Invasive Physiologic Studies of Extremity Arteries	CPT®
Non-invasive physiologic studies of lower extremity arteries, at rest and following treadmill stress testing, complete bilateral study.	93924

Arterial Duplex – Upper and Lower Extremities	CPT®
Duplex scan of lower extremity arteries or arterial bypass grafts; complete bilateral.	93925
<ul style="list-style-type: none"> ➤ A complete duplex scan of the lower extremity arteries includes examination of the full length of the common femoral, superficial femoral and popliteal arteries. ➤ The iliac, deep femoral, and tibioperoneal arteries may also be examined. 	
Duplex scan of lower extremity arteries or arterial bypass grafts; unilateral or limited study.	93926
<ul style="list-style-type: none"> ➤ The limited study is reported when only one extremity is examined or when less than a full examination is performed (e.g. only one or two vessels or follow-up). 	
Duplex scan of upper extremity arteries or arterial bypass grafts; complete bilateral.	93930

Arterial Duplex – Upper and Lower Extremities		CPT®
<ul style="list-style-type: none"> ➤ A complete duplex of the upper extremity arteries includes examination of the subclavian, axillary, and brachial arteries. ➤ The radial and ulnar arteries may also be included. 		
Duplex scan of upper extremity arteries or arterial bypass grafts; unilateral or limited study.		93931
<ul style="list-style-type: none"> ➤ The limited study is reported when only one extremity is examined or when less than a full examination is performed (e.g. only one or two vessels or follow-up). 		
Cerebrovascular Artery Studies		CPT®
Duplex scan of extracranial arteries; complete bilateral study.		93880
Duplex scan of extracranial arteries; unilateral or limited study.		93882
<ul style="list-style-type: none"> ➤ This study is often referred to as a “carotid ultrasound” or “carotid duplex”. ➤ Typically, it includes evaluation of the common, internal, and external carotid arteries. 		
Transcranial Doppler Studies		CPT®
Transcranial Doppler study of the intracranial arteries; complete study		93886
Transcranial Doppler study of the intracranial arteries; limited study		93888
Transcranial Doppler vasoreactivity study		93890
Transcranial Doppler study of the intracranial arteries; emboli detection without intravenous microbubble injection		93892
Transcranial Doppler study of the intracranial arteries; emboli detection with intravenous microbubble injection		93893
Venous Studies - Extremities		CPT®
Non-invasive physiologic studies of extremity veins, complete bilateral study (e.g. Doppler waveform analysis with responses to compression and other maneuvers, phleborheography, impedance plethysmography). This study is rarely performed.		93965
Duplex scan of extremity veins, including responses to compression and other maneuvers; complete bilateral study.		93970
Duplex scan of extremity veins, including responses to compression and other maneuvers; unilateral or limited study.		93971
<ul style="list-style-type: none"> ➤ These codes are used to report studies of lower <i>or</i> upper extremity veins. ➤ A complete bilateral study of the lower extremity veins includes examination of the common femoral, proximal deep femoral, great saphenous and popliteal veins. Calf veins may also be included. ➤ A complete bilateral study of upper extremity veins includes examination of the subclavian, jugular, axillary, brachial, basilica, and cephalic veins. Forearm veins may also be included. 		
Visceral Vascular Studies		CPT®
Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; complete study		93975
Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; limited study		93976
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study		93978
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; unilateral or limited study		93979

Duplex for Hemodialysis Access	CPT®
Duplex scan of hemodialysis access (including arterial inflow, body of access and venous outflow).	93990

PVD-1.3: General Guidelines – Imaging

- ABI should be measured first:
 - ◆ If normal, then further vascular studies are generally not indicated.
 - ◆ If clinical suspicion for PAD remains high with normal ABI's, exercise ABI's (CPT® 93924) can be performed on a treadmill to elicit ischemia
 - ◆ The TBI (toe-brachial index) is used to establish the diagnosis of PAD in the setting of non-compressible arteries (ABI >1.40) and may also be used to assess perfusion in individuals with suspected CLI (rest pain and/or non-healing wound)
- Imaging Studies:
 - ◆ Carotid studies (MRA Neck or CTA Neck) capture the area from the top of the aortic arch (includes the origin of the innominate artery, common carotid artery, and subclavian artery, which gives off the vertebral artery) to the base of the skull.
 - ◆ CTA or MRA Abdomen (CPT® 74175 or CPT® 74185) images from the diaphragm to the umbilicus or iliac crest.
 - ◆ CTA or MRA Chest (CPT® 71275 or CPT® 71555) images from the base of the neck to the dome of the liver.
 - ◆ Runoff studies (CPT® 75635 for CTA or CPT® 74185, CPT® 73725, and CPT® 73725 for MRA) image from the umbilicus to the feet.
 - CTA Abdomen and lower extremities should be reported as CPT® 75635, rather than using the individual CPT® codes for the abdomen, pelvis, and legs
 - MRA Abdomen, MRA Pelvis and MRA Lower extremities should be reported as CPT® 74185, CPT® 73725, and CPT® 73725. The CPT® code for MRA Pelvis (CPT® 72198) should not be included in this circumstance.
 - ◆ If a prior imaging study (Ultrasound, MRA, CTA, Catheter angiogram, etc.) has been completed for a condition, a follow-up, additional, or repeat study for the same condition is generally not indicated unless there has been a change in the individual's condition, previous imaging showed an indeterminate finding, or eviCore healthcare guidelines support routine follow-up imaging.

References

1. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the management of patients with lower extremity peripheral artery disease. *J Am Coll Cardiol.* 2017 Mar 69;(11):1467-1508.
2. Perlstein TS and Creager MA. The ankle-brachial index as a biomarker of cardiovascular risk: it's not just about the legs. *Circulation.* 2009 Nov 29;120(21):2033-2035.

PVD-2: Screening for Suspected Peripheral Artery Disease/Aneurysmal Disease

PVD-2.1: Asymptomatic Screening	10
PVD-2.2: Screening for Vascular Related Genetic Connective Tissue Disorders (Familial Aneurysm Syndromes/Spontaneous Coronary Artery Dissection (SCAD)/Ehlers-Danlos/Marfan/Loeys-Dietz)	10
PVD-2.3: Screening for TAA in Individuals with Bicuspid Aortic Valves	11

PVD-2.1: Asymptomatic Screening

- Routine screening of asymptomatic individuals for PAD is not advised. Those with CVD risk factors should be placed on best medical management and should be questioned on symptoms of PAD at annual physicals
- Resting ABI's (CPT® 93922) may be appropriate in individuals with abnormal pulse exams.
- Currently, there is no evidence to demonstrate that screening all individuals with PAD for asymptomatic atherosclerosis in other arterial beds improves clinical outcome

PVD-2.2: Screening for Vascular Related Genetic Connective Tissue Disorders (Familial Aneurysm Syndromes/Spontaneous Coronary Artery Dissection (SCAD)/Ehlers-Danlos/Marfan/Loeys-Dietz)

- Screening for Familial Syndromes in individuals with a positive family history (1st degree relative with dissection/TAA) but no known genetic syndrome/mutation, otherwise known as Suspected Familial Aneurysm syndrome
 - ◆ ECHO (CPT® 93306, CPT® 93307, or CPT® 93308) and chest x-ray for all First-degree relatives (parents, siblings, children) of individuals with TAA and/or dissection
 - Any imaging listed can be performed if these studies identify a TAA or are equivocal or do not visualize the ascending aorta adequately
 - Studies can be repeated at 2 year intervals if negative
- Follow-Up per TAA Follow-Up guidelines
 - ◆ For individuals with documented Marfan/Loeys-Dietz/Ehlers-Danlos type IV on initial diagnosis full vascular imaging should be performed from head to pelvis with CTA Head, carotid duplex, CTA Chest or CT Chest with contrast, and abdominal duplex. If there are no identified aneurysms or dissections, repeat imaging can be obtained at 2 year intervals
 - ◆ Imaging should be every 6 months once an aneurysm has been identified until a decision has been made to repair
 - Intracranial aneurysm – CTA or MRA Head
 - Aneurysm of a cervical artery – Carotid duplex or CTA Neck if unable to fully visualize with carotid duplex
 - Thoracic aorta – CTA Chest or CT Chest with or without contrast
 - Abdominal aneurysm – Abdominal duplex
 - Visceral aneurysm – These can be difficult to visualize on duplex. If not visible on duplex, can obtain a CTA Abdomen

PVD-2.3: Screening for TAA in Individuals with Bicuspid Aortic Valves

- Screening in individuals with bicuspid aortic valve:
 - ◆ Screening, any requested imaging from the “Table of Thoracic Aorta Imaging Options” in **PVD-6.2: Thoracic Aortic Aneurysm (TAA)** and/or ECHO (CPT® 93306, CPT® 93307, or CPT® 93308)
 - Additional imaging such as Cardiac MRI, Cardiac CT, or CCTA is NOT generally indicated
 - There is no evidence-based data to support screening relatives of individuals with bicuspid aortic valve
 - ◆ Follow-up per TAA Follow-Up guidelines See **PVD-6.2: Thoracic Aortic Aneurysm (TAA)**
- If no dilatation of the aortic root or ascending thoracic aorta is found, there is no evidence-based data to support continued surveillance imaging

References

1. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: Executive Summary. *Journal of the American College of Cardiology*. 2017;69(11):1465-1508. doi:10.1016/j.jacc.2016.11.008.
2. Perlstein TS, Creager MA. The Ankle-Brachial Index as a Biomarker of Cardiovascular Risk. *Circulation*. 2009;120(21):2033-2035. doi:10.1161/circulationaha.109.907238.
3. Saydah SH, Poor Control of Risk Factors for Vascular Disease among Adults with Previously Diagnosed Diabetes. *JAMA*. <https://jamanetwork.com/journals/jama/fullarticle/198035>. Published January 21, 2004.
4. Hennion DR, Siano KA. Diagnosis and Treatment of Peripheral Arterial Disease. *American Family Physician*. <https://www.aafp.org/afp/2013/0901/p306.html#afp20130901p306-c1>. Published September 1, 2013.
5. Hirsh AT. Peripheral Arterial Disease Detection, Awareness, and Treatment in Primary Care. *Jama*. 2001;286(11):1317. doi:10.1001/jama.286.11.1317.
6. Mohler ER, Gornik HL, Gerhard-Herman M, Misra S, Olin JW, Zierler RE. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 Appropriate Use Criteria for Peripheral Vascular Ultrasound and Physiological Testing Part I: Arterial Ultrasound and Physiological Testing. *Journal of the American College of Cardiology*. 2012;60(3):242-276. doi:10.1016/j.jacc.2012.02.009.
7. US Preventive Services Task Force, Curry SJ, Krist AH, et al. Screening for Peripheral Artery Disease and Cardiovascular Disease Risk Assessment With the Ankle-Brachial Index:US Preventive Services Task Force Recommendation Statement. *Jama*. 2018;320(2):177. doi:10.1001/jama.2018.8357.
8. MS Conte, FB Pomposelli, DG Clair, et al. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: Management of asymptomatic disease and claudication. *Journal of Vascular Surgery* 2015. Vol 6:1S-41S.
9. MacCarrick G, III JHB, Bowdin S, et al. Loeys–Dietz syndrome: a primer for diagnosis and management. *Nature News*. <https://www.nature.com/articles/gim201411>. Published February 27, 2014.
10. Olin JW, Gornik HL, Bacharach JM, et al. Fibromuscular Dysplasia: State of the Science and Critical Unanswered Questions. *Circulation*. 2014;129(9):1048-1078. doi:10.1161/01.cir.0000442577.96802.8c.
11. Persu A, Niepen PVD, Touzé E, et al. Revisiting Fibromuscular Dysplasia. *Hypertension*. 2016;68(4):832-839. doi:10.1161/hypertensionaha.116.07543.
12. Hayes SN, Kim ES, Saw J, et al. Spontaneous Coronary Artery Dissection: Current State of the Science: A Scientific Statement From the American Heart Association. *Circulation*. 2018;137(19). doi:10.1161/cir.0000000000000564.

13. Hiratzka LF, Bakris GL, Beckman JA, et al. 2010
ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with thoracic aortic disease. *J Am Coll Cardiol* 2010; 55: e27-e129.

PVD-3: Cerebrovascular and Carotid Disease

PVD-3.1: Initial Imaging	14
PVD-3.2: Surveillance Imaging with NO History of Carotid Surgery or Intervention	16
PVD-3.3: Surveillance Imaging WITH History of Carotid Surgery or Intervention	17

PVD-3.1: Initial Imaging

- Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral), prior to considering advanced imaging, should generally be used to evaluate possible carotid artery disease when ANY of the following apply:
 - ◆ Hemispheric neurologic symptoms including stroke, TIA, or amaurosis fugax
 - ◆ Known or suspected retinal arterial emboli or Hollenhorst plaque
 - ◆ Suspected carotid dissection
 - ◆ Pulsatile neck masses
 - ◆ Carotid or cervical bruit
 - ◆ Abnormal findings on physical exam of the carotid arteries (e.g. aneurysm or absent carotid pulses)
 - ◆ Preoperative evaluation of individuals with evidence of severe diffuse atherosclerosis, scheduled for major cardiovascular surgical procedures
 - ◆ Preoperative evaluation of individuals prior to elective coronary artery bypass graft (CABG) surgery in individuals older than 65 years of age and in those with peripheral artery disease, history of cigarette smoking, history of stroke or TIA, or carotid bruit
 - ◆ Suspected Subclavian Steal Syndrome
 - See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging Guidelines
 - ◆ Blunt neck trauma
 - ◆ Neurologic complaints after chiropractic neck manipulation
 - ◆ Vasculitis potentially involving carotid arteries, i.e. Takayasu's arteritis and fibromuscular dysplasia (FMD)
- Carotid ultrasound screening in asymptomatic individuals due only to risk factors is not indicated
- New signs and symptoms consistent with carotid artery disease (e.g. TIA, amaurosis fugax, change in nature of a carotid bruit) are an indication to re-image the cervical vessels (regardless of when the previous carotid imaging was performed) using ANY of the following:
 - ◆ Duplex ultrasound (CPT® 93880 bilateral study or CPT® 93882 unilateral study),
 - ◆ MRA Neck with contrast (CPT® 70548) or without and with contrast (CPT® 70549),
 - ◆ CTA Neck (CPT® 70498)
- For Typical Symptoms of TIA/Stroke or Carotid Dissection:
 - ◆ See **HD-21: Stroke/TIA** in the Head Imaging Guidelines
- For Suspected Vertebrobasilar Pathology:
 - ◆ Symptoms include:
 - Vertigo associated with nausea and vomiting
 - Diplopia
 - Loss of vision in one or both eyes
 - Dysarthria
 - Bifacial numbness
 - Bilateral extremity weakness and/or numbness
 - Acute changes in mental status

- Loss of consciousness
 - Ataxia
- ◆ Mechanisms of injury for concern of arterial dissection including, but not exclusive to:
 - Chiropractic manipulation of neck
 - Whiplash injury
 - Fibromuscular dysplasia
 - Stroke in the young (age ≤50)
- ◆ Initial Imaging
 - Carotid duplex-Note: Carotid duplex provides limited information on vertebral disease
 - If clinical suspicion is high
 - CTA Neck/MRA Neck can be considered medically necessary.
 - MRA/CTA Neck and Head for evaluation of posterior circulation disease, both are required to visualize the entire vertebral-basilar system. See **HD-1.5: General Guidelines – CT and MR Angiography** in the Head Imaging Guidelines
 - See **HD-21: Stroke/TIA** in the Head Imaging Guidelines
- ◆ Surveillance imaging post-stenting or known vertebrobasilar disease interval determined by Vascular Specialist, Neurologist, or Neurosurgeon for ANY of the following:
 - Asymptomatic
 - Unchanged symptoms
 - New or worsening symptoms
- After Intracranial Hemorrhage:
 - ◆ Initial Imaging See **HD-13.1: Head Trauma** in the Head Imaging Guidelines
 - ◆ Surveillance Imaging
 - Interval determined by neurosurgeon or neurologist
- For Suspected Subclavian Steal Syndrome:
 - ◆ Initial imaging should be a carotid duplex
 - If initial duplex demonstrates high grade stenosis or occlusion of the subclavian artery, advanced imaging is NOT indicated unless the individual is symptomatic with arm claudication or signs of hypo-perfusion of the vertebral artery with recurrent dizziness
 - ◆ Surveillance of subclavian arterial disease is NOT indicated if there has not been any intervention such as a carotid-subclavian bypass or subclavian stent
 - ◆ Advanced imaging See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging

PVD-3.2: Surveillance Imaging with NO History of Carotid Surgery or Intervention

- Surveillance imaging once a year for individuals with fibromuscular dysplasia of the extracranial carotid arteries.
- Reporting standards for carotid stenosis varies widely. The most commonly used criteria, however, is noted in the chart below published by the Society of Radiology in 2003

Degree of Stenosis (%)	Primary Parameters		Additional Parameters	
	ICA PSV (cm/sec)	Plaque Estimate (%)*	ICA/CCA PSV Ratio	ICA EDV (cm/sec)
Normal	<125	None	<2.0	<40
<50	<125	<50	<2.0	<40
50–69	125–230	≥50	2.0–4.0	40–100
≥70 but less than near occlusion	>230	≥50	>4.0	>100
Near occlusion	High, low, or undetectable	Visible	Variable	Variable
Total occlusion	Undetectable	Visible, no detectable lumen	Not applicable	Not applicable

- If normal study, no routine follow-up imaging is indicated
- If <50% carotid stenosis
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) can be performed every two years
- Between 50% and 70% carotid stenosis
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) can be performed annually
 - ◆ A repeat duplex (CPT® 93880 bilateral or CPT® 93882 unilateral) may be performed in three to six months until stability is reached when ONE of the following occurs:
 - Change in the character of the bruit
 - Duplex demonstrates rapid progression, including:
 - Doubling of peak systolic velocities
 - Increase of the ICA/CCA ratio
 - Heavy calcification
 - Thrombus
 - Ulcerated plaque
 - Echolucent plaque
- Carotid stenosis ≥70% or ICA/CCA ratio >4
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) or MRA Neck with contrast (CPT® 70548) or CTA Neck (CPT® 70498) at the following intervals:
 - Every 6 months until ONE of the following occurs:

- Intervention is performed
- Decision is made to not intervene
- MRA Neck with contrast (CPT® 70548) or CTA Neck (CPT® 70498) if duplex Ultrasound shows $\geq 70\%$ occlusion/stenosis of the internal carotid artery OR the ICA/CCA ratio is >4.0 even with a lower percentage of stenosis
 - ◆ If carotid stent is planned
 - MRA Head (CPT® 70544, or CPT® 70545, or CPT® 70546) or CTA Head (CPT® 70496) can be added

PVD-3.3: Surveillance Imaging WITH History of Carotid Surgery or Intervention

- Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) post carotid surgery or intervention at the following intervals:
 - ◆ 1 month after procedure
 - ◆ Every 6 months for 2 years after procedure
 - ◆ Then annually
- If $\geq 70\%$ residual carotid stenosis is seen at 1 month after procedure
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) at the following intervals:
 - Every 3-6 months for one year
 - Then annually

Background and Supporting Information

- Carotid intima-media thickness using duplex ultrasound imaging (Category III code 0126T) is not recommended in clinical practice for risk assessment for a first ASCVD event. Although outcomes data are lacking, Texas has adopted this method in Texas Heart Attack Preventive Screening Bill (HR 1290).
- Texas Heart Attack Preventive Screening Law (HR 1290) mandates that insurers in Texas cover either a calcium scoring study (CPT® 75571 or HCPCS S8092) or a carotid intima-media thickness study (ultrasound—Category III code 0126T) every five years for certain populations. To qualify, the following must apply:
 - ◆ Must be a Texas resident
 - ◆ Must be a member of a fully-insured Texas health plan
 - ◆ Must be a man age 45 to 75 or a woman age 55 to 75
 - ◆ Must have either diabetes or a Framingham cardiac risk score of intermediate or higher
 - ◆ Must not have had a calcium scoring study or a carotid intima-media thickness study within the past 5 years
- MRA Neck (CPT® 70548) or CTA Neck (CPT® 70498) may be indicated if ultrasound is technically difficult or confirmation of the degree of stenosis on ultrasound is needed because an interventional procedure is being considered.

References

1. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS Guideline on the Management of Patients With Extracranial Carotid and Vertebral Artery Disease. *Journal of the American College of Cardiology*. 2011;57(8). doi:10.1016/j.jacc.2010.11.006.
2. Gerhard-Herman M, Gardin JM, Jaff M, Mohler E, Roman M, Naqvi TZ. Guidelines for Noninvasive Vascular Laboratory Testing: A Report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology. *Journal of the American Society of Echocardiography*. 2006;19(8):955-972. doi:10.1016/j.echo.2006.04.019.
3. Reutern G-MV, Goertler M-W, Bornstein NM, et al. Grading Carotid Stenosis Using Ultrasonic Methods. *Stroke*. 2012;43(3):916-921. doi:10.1161/strokeaha.111.636084.
4. ACR–AIUM–SPR–SRU. AIUM Practice Parameter for the Performance of an Ultrasound Examination of the Extracranial Cerebrovascular System. *Journal of Ultrasound in Medicine*. 2016;35(9):1-11. doi:10.7863/ultra.35.9.1.
5. Moresoli P, Habib B, Reynier P, et al. Carotid Stenting versus Endarterectomy for Asymptomatic Carotid Artery Stenosis. *Stroke*. 2017;48(8):2150-2157. doi:10.1161/strokeaha.117.016824.
6. Rogers RK, Bishu K. Optimal Treatment of Extracranial Carotid Artery Disease: Carotid Endarterectomy, Carotid Stenting, or Optimal Medical Therapy. *Current Cardiology Reports*. 2015;17(10). doi: 10.1007/s11886-015-0636-2.
7. Shakarchi JA, Lowry D, Nath J, et al. Duplex ultrasound surveillance after carotid artery endarterectomy. *Journal of Vascular Surgery*. 2016;63(6):1647-1650. doi:10.1016/j.jvs.2016.01.054.
8. Aburahma AF, Srivastava M, Aburahma Z, et al. The value and economic analysis of routine postoperative carotid duplex ultrasound surveillance after carotid endarterectomy. *Journal of Vascular Surgery*. 2015;62(2):378-384. doi:10.1016/j.jvs.2015.03.023.
9. Aboyans V, Ricco J-B, Bartelink M-LEL, et al. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS). *European Heart Journal*. 2017;39(9):763-816. doi:10.1093/eurheartj/ehx095.
10. Naylor A, Ricco J-B, Borst GD, et al. Editor's Choice – Management of Atherosclerotic Carotid and Vertebral Artery Disease: 2017 Clinical Practice Guidelines of the European Society for Vascular Surgery (ESVS). *European Journal of Vascular and Endovascular Surgery*. 2018;55(1):3-81. doi:10.1016/j.ejvs.2017.06.021.
11. Mohler ER, Gornik HL, Gerhard-Herman M, Misra S, Olin JW, Zierler RE. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 Appropriate Use Criteria for Peripheral Vascular Ultrasound and Physiological Testing Part I: Arterial Ultrasound and Physiological Testing. *Journal of the American College of Cardiology*. 2012;60(3):242-276. doi:10.1016/j.jacc.2012.02.009.
12. Goff DC, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk. *Circulation*. 2014;129(25 suppl 2):s49-s73. doi:10.1161/01.cir.0000437741.48606.98.
13. Ricotta JJ, Aburahma A, Ascher E, et al. Updated Society for Vascular Surgery guidelines for management of extracranial carotid disease. *Journal of Vascular Surgery*. 2011;54(3):1-31. doi:10.1016/j.jvs.2011.07.031.
14. Ballotta E, Giau GD, Meneghetti G, et al. Progression of atherosclerosis in asymptomatic carotid arteries after contralateral endarterectomy: A 10-year prospective study. *Journal of Vascular Surgery*. 2007;45(3):516-522. doi:10.1016/j.jvs.2006.11.011.
15. Derdeyn CP, Zipfel GJ, Albuquerque FC, et al. Management of Brain Arteriovenous Malformations: A Scientific Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. *Stroke*. 2017;48(8). doi:10.1161/str.000000000000134.

PVD-4: Upper Extremity Peripheral Vascular Disease

PVD-4.1: Upper Extremity PVD – Imaging

20

PVD-4.1: Upper Extremity PVD – Imaging

- For signs and symptoms of arterial insufficiency, appropriate studies include:
 - ◆ Arterial ultrasound upper extremities (CPT® 93930 or CPT® 93931), or
 - ◆ CTA Upper extremity (CPT® 73206) or MRA Upper extremity (CPT® 73225), and/or
 - ◆ CTA Chest (CPT® 71275) or MRA Chest (CPT® 71555).
- For suspected Fibromuscular Dysplasia of the brachial artery, appropriate studies include:
 - ◆ MRA Upper extremity (CPT® 73225).
 - ◆ CTA Upper extremity (CPT® 73206).
 - ◆ Arterial Ultrasound (CPT® 93930 bilateral study or CPT® 93931 unilateral study).
- Arterial Duplex (CPT® 93931) following upper extremity arterial revascularization at:
 - ◆ Baseline (within one month)
 - ◆ 6 months
 - ◆ Then annually if stable
 - ◆ Anytime for new or worsening symptoms
- For symptoms of venous insufficiency including but not limited to unilateral pain and swelling of the upper extremity
 - ◆ Venous duplex upper extremities (CPT® 93970 or CPT® 93971) should be performed initially.
 - ◆ If duplex ultrasound is nondiagnostic:
 - MRV Upper extremity (CPT® 73225) and/or MRV Chest (CPT® 71555), or
 - CTV Upper extremity (CPT® 73206) and/or CTV Chest (CPT® 71275).
 - CTV Upper extremity (CPT® 73206) or MRV Upper extremity (CPT® 73225), and/or CTV Chest (CPT® 71275) or MRV Chest (CPT® 71555) if there is a history of exertion with the limb such as with weight lifting or in the presence of central venous access (port, PICC line, to name a few) with a negative venous duplex. See **CH-31.1: Thoracic Outlet Syndrome** in the Chest Imaging Guidelines
- For Superior Vena Cava Syndrome (upper extremity and facial symptoms):
 - ◆ CT Chest with contrast (CPT® 71260).
 - ◆ MRV (CPT® 71555) or CTV (CPT® 71275) Chest may be considered when stenting of the SVC is being considered.

Background and Supporting Information

- Signs and symptoms of arterial insufficiency include but are not limited to:
 - ◆ Arm or hand claudication, cramping or fatigue of the unilateral extremity with use or with raising limb overhead that is relieved with rest and is reproducible. See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging Guidelines
 - ◆ Systolic blood pressure differential between arms of <15mmHg. See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging Guidelines
 - ◆ Bluish discoloration of the hand or fingers
 - ◆ Unilateral cold painful pulseless hand

- ◆ Non healing wound (>2 weeks with no healing or evidence of healing) or frank gangrene

References

1. Desjardins B, Rybicki FJ, Kim HS, et al. ACR Appropriateness Criteria® Suspected Upper Extremity Deep Vein Thrombosis. *Journal of the American College of Radiology*. 2012;9(9):613-619. doi:10.1016/j.jacr.2012.05.021.
2. Yoshimuta T, Akutsu K, Okajima T, et al. "String of Beads" Appearance of Bilateral Brachial Artery in Fibromuscular Dysplasia. *Circulation*. 2008;117(19):2542-2543. doi:10.1161/circulationaha.107.747089.
3. Birch A, Um D, Laselle B. Ultrasound Detection of Superior Vena Cava Thrombus. *Western Journal of Emergency Medicine*. 2014;15(6):715-718. doi:10.5811/westjem.2014.6.14006.

PVD-5: Pulmonary Artery Hypertension

PVD-5.1: Pulmonary Artery Hypertension – Imaging

23

PVD-5.1: Pulmonary Artery Hypertension – Imaging

- Pulmonary artery hypertension (PAH) comprises a spectrum of diseases which will need direct evaluation, including ECG (right ventricular hypertrophy with/without strain, right atrial dilatation); chest x-ray; arterial blood gas, PFT's or V/Q scan. Imaging is based on suspected etiology.
- Transthoracic echocardiogram (TTE) (CPT® 93306) should be performed initially and may be accompanied by:
 - ◆ Pulmonary venous hypertension - Stress echocardiogram (CPT® 93350 or CPT® 93351) or left and/or right heart catheterization.
 - ◆ Pulmonary hypertension associated with hypoxemia - High resolution CT Chest (CPT® 71250) to rule out restrictive lung disorders such as idiopathic pulmonary fibrosis.
- Acute or chronic pulmonary embolism – CTA Chest (CPT® 71275)
- See specific subsections
 - ◆ **CD-2.2: Transthoracic Echocardiogram (TTE) - Indications** in the Cardiac Imaging Guidelines
 - ◆ **CD-7.4: Right Heart Catheterization (RHC)** in the Cardiac Imaging Guidelines
 - ◆ **CD-11.3.12: Severe Pulmonary artery hypertension (PHT) and Eisenmenger syndrome** in the Cardiac Imaging Guidelines
 - ◆ **PEDCD-2.3: Congenital Heart Disease Modality Considerations** in the Pediatric Cardiac Imaging Guidelines
 - ◆ **PEDCD-7: Pediatric Pulmonary Hypertension-General** in the Pediatric Cardiac Imaging Guidelines
 - ◆ **CH-25: Pulmonary Embolism (PE)** in the Chest Imaging Guidelines

References

1. Barbosa EJM, Gupta NK, Torigian DA, et al. Current role of imaging in the diagnosis and management of pulmonary hypertension. *AJR Am J Roentgenol.* 2012; 198 (6): 1320-1331.
2. Galiè A, Torbicki A, Barst R, et al. Guidelines on diagnosis and treatment of pulmonary arterial hypertension: the task force on diagnosis and treatment of pulmonary arterial hypertension of the European Society of Cardiology. *Eur Heart J.* 2004 Dec 1; 25 (24): 2243-2278.
3. Peña E, Dennie C, Veinot J, et al. Pulmonary hypertension: how the radiologist can help. *RadioGraphics.* 2012; 32 (1):9-32.

PVD-6: Aortic Disorders, Renal Vascular Disorders and Visceral Artery Aneurysms

PVD-6.1: Aortic Disorders General Information	25
PVD-6.2: Thoracic Aortic Aneurysm (TAA)	26
PVD-6.3: Abdominal Aortic Aneurysm (AAA)	28
PVD-6.4: Iliac Artery Aneurysm (IAA)	29
PVD-6.5: Visceral Artery Aneurysm	30
PVD-6.6: Renovascular Hypertension/Renal Artery Stenosis	31
PVD-6.7: Aortic Dissection and Other Aortic Conditions	33
PVD-6.8: Post Aortic Endovascular/Open Surgery Surveillance Studies	35

PVD-6.1: Aortic Disorders General Information

Duplex ultrasound for visceral vascular studies	CPT®
➤ Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; complete study.	93975
➤ Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; limited study.	93976
➤ Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study.	93978
➤ Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; unilateral or limited study.	93979

- In clinical practice, CT, CTA, MRA are usually preferred to evaluate for stenosis of these vessels rather than ultrasound (Exception: Duplex ultrasound is appropriate to rule out testicular or ovarian torsion or to evaluate an abdominal bruit or a pulsatile abdominal mass).
- Mesenteric Ischemia
 - ◆ See **AB-6: Mesenteric/Colonic Ischemia** in the Abdomen Imaging Guidelines.

References

1. Hirsch AT, Haskal ZJ, Hertzler NR, et al. ACC/AHA Practice Guidelines: ACC/AHA 2005 Practice guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery,* Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines. *Circulation*. 2006; 113: e463-e654. <http://circ.ahajournals.org/content/113/11/e463>.
2. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. This article is copublished in the *Journal of the American College of Cardiology*, *Catheterization and Cardiovascular Interventions*, the *Journal of Vascular Surgery*, and *Vascular Medicine*. *Circulation*. 2011 Nov 1; 124 (18): 2020-2045. <http://circ.ahajournals.org/content/124/18/2020>.
3. Mohler III ER, Gornik HL, Gerhard-Herman M, et al. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 Appropriate use criteria for peripheral vascular ultrasound and physiological testing part I: arterial ultrasound and physiological testing. *J Am Coll Cardiol* 2012;60:242–76. [https://www.jvascsurg.org/article/S0741-5214\(12\)01174-3/fulltext](https://www.jvascsurg.org/article/S0741-5214(12)01174-3/fulltext).

PVD-6.2: Thoracic Aortic Aneurysm (TAA)

- Advanced imaging with a CT or MR is preferred imaging for this diagnosis. Given the diversity of studies, pathology and provider preference, approved thoracic imaging for this indication can be ONE of the following studies listed in the table below:

Table of Thoracic Aorta Imaging Options	CPT®
CT Chest, and/or Abdomen, and/or Pelvis	71260
	74177
	74160
	72193
CTA Chest, and/or Abdomen, and/or Pelvis	71275
	74175
	72191
	74174
MRA Chest, and/or Abdomen, and/or Pelvis	71555
	74185
	72198

- For suspected TAA, any requested imaging from the “Table of Thoracic Aorta Imaging Options” above:
 - ◆ Abnormalities identified on chest x-ray (abnormality including widened mediastinum, suspicious calcifications) or other imaging studies (fluoroscopy, MRI Spine, etc.) abnormality.^{1,2,3,4,5}
- For known TAA accompanied with chest pain or back pain and suspicion of rupture, any requested imaging from the “Table of Thoracic Aorta Imaging Options” above.^{1,2,3,4,5}
- For planning for pre–thoracic endovascular repair (TEVAR) of thoracic aorta disease.⁹
 - ◆ CTA Chest, and/or Abdomen, and/or Pelvis (CPT® 71275, CPT® 74175, CPT® 72191, CPT® 74174); or
 - ◆ MRA Chest, and/or Abdomen, and/or Pelvis (CPT® 71555, CPT® 74185, CPT® 72198)
- For follow-up of ascending aortic aneurysms
 - ◆ Operative treatment is reasonable for asymptomatic individuals when the diameter of the arch exceeds 5.5 cm
 - ◆ **For individuals with isolated aortic arch aneurysms <4.0 cm in diameter**
 - Repeat imaging annually
 - ◆ For individuals with isolated aortic arch aneurysms >4.0 cm
 - Repeat imaging 6 months
- For follow-up of descending aortic aneurysms, any requested imaging from the “Table of Thoracic Aorta Imaging Options” above for the following:^{4,5,7,9}
 - ◆ “Medically” treated/observation
 - 3.5 to 4.4 cm TAA can be followed annually
 - ≥4.5 cm TAA can be followed every 6 months
 - ≥3.0 cm TAA when there is concern for growth can have a one-time 3 month interval advanced imaging

PVD-6.3: Abdominal Aortic Aneurysm (AAA)

- Ultrasound Abdominal aorta (CPT® 76706) is the preferred initial imaging study to screen and retroperitoneal ultrasound (CPT® 76775) to survey for AAA or to evaluate a pulsatile abdominal mass.
- **Obese Individual (BMI ≥35):** CT Abdomen and Pelvis with contrast (CPT® 74177) or without contrast (CPT® 74176) can be substituted for US using the same timeline as a non-obese individual. Ultrasound abdominal aorta should ideally first be attempted to see if the image quality is adequate
- Screening
 - ◆ One-time screening recommendations for AAA (Ultrasound CPT® 76706)
 - Men and women 65 to 75 years of age with a history of tobacco use
 - Men and women older than 75 years with a history of tobacco use and in otherwise good health who have not previously received a screening ultrasound examination
 - All first-degree relatives of individuals who present with an AAA and are between 65 and 75, or in those older than 75 in good health
 - ◆ Ultrasound (CPT® 76706) for AAA screening is reasonable if there is a documented thoracic aortic aneurysm; however, there is insufficient evidence to support the use of advanced imaging to screen for a thoracic aortic aneurysm in individuals with known abdominal aortic aneurysm.
- Surveillance recommendations for AAA (CPT® 76775)
 - ◆ >2.5 cm but <3.0 cm: 10 years
 - ◆ 3.0 cm to 3.9 cm: 3 year intervals
 - ◆ 4.0 cm to 4.9 cm: every 12 months
 - ◆ 5.0 cm to 5.4 cm: every 6 months
 - ◆ >5.4 cm or aortic diameter has increased in size by 0.7 cm in six months, or at least 1 cm in a year may undergo more frequent monitoring and should be evaluated by a Vascular Specialist.
- Additional Imaging
 - ◆ CT Abdomen and Pelvis with contrast (CPT® 74177), or CT Abdomen and Pelvis without and with contrast (CPT® 74178), or CTA Abdomen and Pelvis (CPT® 74174), or CTA Abdomen (CPT® 74175), or CTA Pelvis (CPT® 72191)
 - Individuals suspected to have AAA presenting with recent-onset abdominal or back pain, particularly in the presence of a pulsatile epigastric mass or significant risk factors for AAA
 - Pre-operative imaging for AAA repair

PVD-6.5: Visceral Artery Aneurysm

- Treatment is generally indicated for aneurysm >2cm
- Workup for suspected visceral artery aneurysm (spleen, kidney, liver or intestines) if calcifications seen on plain film imaging can include:
 - ◆ Ultrasound (CPT® 76700 or CPT® 76705), **or**
 - ◆ CTA Abdomen (CPT® 74175), **or**
 - ◆ CT Abdomen with contrast (CPT® 74160).
- Ultrasound (CPT® 76700 or CPT® 76705) **or** CTA Abdomen (CPT® 74175) **or** CT Abdomen with contrast (CPT® 74160) for further monitoring based on the intervals below or as determined by a vascular specialist:
 - ◆ Splenic artery aneurysms:
 - <20mm can be imaged every three years
 - If >25mm, they should be referred for treatment, either stent, excision or splenectomy
 - ◆ For all other visceral artery aneurysms:
 - Initial evaluation with six-month follow-up for one year
 - Further follow-up annually if no significant enlargement is seen
- CTA Abdomen (CPT® 74175), MRA Abdomen (CPT® 74185), or CT Abdomen (CPT® 74160) are indicated following stent placement at:
 - ◆ 1 month
 - ◆ 6 months
 - ◆ 12 months
 - ◆ Then every year

Background and Supporting Information

- Visceral Artery Aneurysms are defined by an increase of more than 50% of the original arterial diameter.
- Vascular specialty consultation is beneficial in order to determine the time frame to intervention.
- Splenic artery aneurysms, the most common (60%), tend to exhibit very slow rates of growth, while the other visceral artery aneurysms are more unpredictable in their rate of growth with a greater tendency to rupture

References

1. Zierler RE, Jordan WD, Lal BK, et al. The Society for Vascular Surgery practice guidelines on follow-up after vascular surgery arterial procedures. *Journal of Vascular Surgery*. 2018;68(1):256-284. doi:10.1016/j.jvs.2018.04.018.
2. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *Journal of Vascular Surgery*. 2018;67(1). doi:10.1016/j.jvs.2017.10.044.
3. Corey MR, Ergul EA, Cambria RP. The natural history of splanchnic aneurysms and outcome after operative intervention. *J Vasc Surg*. 2016 April 63 (4):949-57.
4. Erben Y, Brownstein AJ, Rajaei S. Natural history of and management of splanchnic artery aneurysms in a single tertiary referral center. *J Vasc Surg* 2018 Oct; 68(4): 1079-1087

PVD-6.6: Renovascular Hypertension/Renal Artery Stenosis

- MRA Abdomen without or with contrast (CPT® 74185) or CTA Abdomen with contrast (CPT® 74175) if:
 - ◆ The individual is adherent to full doses of three blood pressure medications (including a diuretic) yet has still not achieved goal.
 - ◆ Sudden and persistent worsening of previously controlled hypertension
 - ◆ Onset of hypertension younger than 30 years of age.
 - ◆ Malignant hypertension with coexistent evidence of acute end-organ damage (acute renal failure, new visual or neurological disturbance and/or advanced retinopathy) or flash pulmonary edema.
 - ◆ Women who develop hypertension ($\geq 140/90$) within the first 20 weeks of pregnancy, if hypertension persists > 12 weeks post-partum.
 - ◆ New or worsening renal function/increasing creatinine (especially after the administration of an ACE inhibitor or with angiotensin receptor blocking agent).
 - ◆ Unexplained atrophic kidney or discrepancy in size between kidneys of greater than 1.5cm.
- Gadolinium agents may be contraindicated in individuals with severe renal disease or on dialysis due to the risk of developing nephrogenic systemic sclerosis
- US kidney retroperitoneal (CPT® 76775) and/or Doppler (CPT® 93975 or CPT® 93976) if expertise is available
- Screening carotid duplex (CPT® 93880) is reasonable to assess for carotid involvement in individuals with documented or highly suspicious renal artery stenosis due to fibromuscular dysplasia (mostly women between 15 and 50 years of age). CTA Abdomen (CPT® 74175) or MRA Abdomen (CPT® 74185) to screen for renovascular fibromuscular dysplasia in hypertensive individuals with documented cervicocephalic fibromuscular dysplasia. The assessment of other vascular beds should be considered if supported by suggestive symptoms or medical history.

Background and Supporting Information

- Renal artery revascularization has NOT been shown to be more effective than medical therapy in most situations and should not be pursued except in extreme cases, or if there is concern for Takayasu arteritis or fibromuscular dysplasia

PVD-6.7: Aortic Dissection and Other Aortic Conditions

- CTA or MRA entire aorta (including arch branches) and extending through the femoral arteries for suspected aortic dissection.^{1,2,3,4,5}
- ANY of the following studies can be used if acute dissection is suspected:
 - ◆ CT Chest (CPT® 71260 or CPT® 71270) and/or CT Abdomen (CPT® 74160 or CPT® 74170) and/or CT Pelvis (CPT® 72193 or CPT® 72194) or
 - If CT Abdomen and Pelvis with or with and without is requested, codes: (CPT® 74177 or CPT® 74178) are appropriate.
 - ◆ CTA Chest (CPT® 71275) and/or CTA Abdomen and/or Pelvis (CPT® 74175 or CPT® 72191 or CPT® 74174), or
 - ◆ MRA Chest and/or Abdomen and/or Pelvis (CPT® 71555 and/or CPT® 74185 and/or CPT® 72198)
- Chronic Aortic Dissections
 - ◆ 1/3 of individuals with chronic type B dissections that were not treated via open or endovascular repair will go on to develop aneurysmal disease requiring subsequent intervention. Advanced imaging of the affected segment of the aorta can be performed as follows with any of the studies noted above:
 - In individuals with a persistent false lumen or initial aortic diameter of >4cm:
 - Every 6 months for two years until stability has been reached
 - Then annually
 - In individuals with initial aortic diameter of <4cm and/or a thrombosed false lumen:
 - Annually
 - Any time if the individual is symptomatic with chest pain, back pain or has any evidence of end organ ischemia: renal dysfunction, mesenteric ischemia or acute limb ischemia
 - In individuals with Marfan syndrome/Loeys-Dietz/Ehlers-Danlos
 - ◆ As aneurysmal expansion within a dissection can occur rapidly, post-dissection imaging in these individuals is indicated as follows:
 - 1 month
 - 3 months
 - 6 months
 - 12 months
 - Yearly thereafter
 - ◆ Depending on the location of the dissection the following may be approved:
 - CTA or MRA Head
 - Carotid duplex or CTA Neck or MRA Neck
 - CTA or MRA Chest
 - CTA or MRA Abdomen and Pelvis or CTA or MRA Abdomen or CTA or MRA Pelvis

PVD-6.8: Post Aortic Endovascular/Open Surgery Surveillance Studies

- Aortic root/ascending aortic aneurysm repair post-operative echocardiography (TEE/TTE) can be obtained:
 - ◆ Every three months for the first year
 - ◆ Every six months year 2
 - ◆ Annually thereafter

PVD-6.8.1: Post-operative surveillance after TEVAR for any indication

Imaging for post-operative abdominal EVAR	CPT®
CT Chest, and/or Abdomen, and/or Pelvis	71260 74177 74160 72193
CTA Chest, and/or Abdomen, and/or Pelvis	71275 74175 72191 74174
MRA Chest, and/or Abdomen, and/or Pelvis	71555 74185 72198

- ANY of the above studies listed in the table can be performed at one month, six months, twelve months and then annually
- Abdomen and Pelvis imaging is indicated if TEVAR performed for a dissection that extends into the abdomen or pelvis

PVD-6.8.2: Post-operative surveillance after abdominal EVAR (endovascular aneurysm repair)

Imaging for post-operative abdominal EVAR	CPT®
CT Abdomen and/or Pelvis with contrast	74160 72193 74177
CT Abdomen and/or Pelvis without and with contrast	74170 72194 74178
CTA Abdomen and/or Pelvis	74175 72191 74174
MRA Abdomen and/or Pelvis	74185 72198

- CT as per above coding as requested and color duplex ultrasound (CPT® 93975 or CPT® 93976) one month after EVAR

PVD-7: Lower Extremity Peripheral Vascular Disease

PVD-7.1: Claudication	39
PVD-7.2: Popliteal Artery Entrapment Syndrome	40
PVD-7.3: Post-Procedure Surveillance Studies	40
PVD-7.4: Lower Extremity Artery Aneurysms	43

PVD-7.1: Claudication

- Resting ABI for initial evaluation for suspected PAD. This can be accomplished at the bedside as part of the physical examination or requested as CPT® 93922 (limited Doppler ultrasound) or CPT® 93923 (multi-level complete Doppler ultrasound).
 - ◆ CPT® 93923 may be performed once.
 - ◆ Follow-up studies may be performed with CPT® 93922.
 - ◆ Post-exercise ABI (CPT® 93924) can be performed if the resting ABI is >0.89 and PAD is still highly suspected clinically.
- History and physical suggestive of PAD include:
 - ◆ History
 - Claudication - reproducible calf or thigh cramping with exertion that is relieved completely with rest
 - Critical limb ischemia
 - Rest pain suggestive of ischemia - pain in the ball of foot when the leg is in an elevated position particularly at night
 - Distal non-healing wound or punched out ulcer with sharply demarcated edges present for >2 weeks with no evidence of healing, i.e. presence of granulation tissue
 - ◆ Physical Examination
 - Abnormal lower extremity pulse examination
 - Vascular bruit
 - Non-healing lower extremity wound
 - Lower extremity gangrene
 - Other suggestive lower extremity physical findings (e.g., elevation pallor/dependent rubor)
 - Atrophic nails, hair loss, shiny skin
- If resting ABI (CPT® 93922) is normal (0.9 to 1.3) and disease is still suspected:
 - ◆ Differentiate from “pseudoclaudication” See **SP-9: Lumbar Spinal Stenosis** in the Spine Imaging Guidelines
 - ◆ Re-measure ABI after exercise (CPT® 93924)¹
 - ◆ A toe-brachial index may be used as further screening in individuals with ABI’s greater than 1.3
 - ◆ Advanced imaging is necessary only if there is consideration for invasive therapy^{2,3,4,5} not to confirm diagnosis
- Duplex ultrasound (CPT® 93925 bilateral study or CPT® 93926 unilateral study) and Doppler studies are adjuncts to abnormal ABI that may be used to identify location and extent of disease once there has been a decision for revascularization:^{6,7}
- MRA Aorta and Pelvic vessels, and Lower extremities (CPT® 74185, CPT® 73725 and CPT® 73725), **or** CTA with run off (CPT® 75635) to further evaluate the lower extremity arteries for the purpose of preoperative planning for ANY of the following:^{2,8}
 - ◆ Intermittent claudication (i.e. non-limb threatening ischemia) and either:
 - Failed 3 months conservative medical therapy (physician supervised walking/exercise program plus medical therapy), **or**

- Functional disability (e.g. exercise impairment sufficient to threaten the individual's employment or to require significant alterations in the individual's lifestyle)
- ◆ Potentially limb-threatening vascular disease evidenced by:
 - Skin breakdown
 - Non-healing ischemic ulcers
 - Resting leg pain
 - Gangrene
- ◆ Blue Toe Syndrome:
 - Emboli from aortic plaque or mural thrombus
 - Hyperviscosity syndrome
 - Hypercoagulable states
 - Vasculitis
- ◆ Note: MRA Pelvis should not be requested/billed with CPT® 74185, CPT® 73725 and CPT® 73725.

Background and Supporting Information

Claudication symptoms usually remain stable (70% to 80% of individuals) and do not worsen or improve at rapid rates. Repeat studies to assess the efficacy of medical therapy are not indicated unless there is a negative change in clinical status for the purpose of preoperative planning such as worsening claudication or progression to critical limb ischemia.

PVD-7.2: Popliteal Artery Entrapment Syndrome

- Diagnosis of popliteal artery stenosis or occlusion due to compression by adjacent muscle and tendons seen in young men (ages 20 to 40).¹⁰
 - ◆ Ultrasound (CPT® 93926 unilateral study), CTA Lower extremity (CPT® 73706), or MRA Lower extremity (CPT® 73725).
 - ◆ CT or MRI Lower Extremity (contrast as requested) if requested by the operating surgeon.

PVD-7.3: Post-Procedure Surveillance Studies

- Scheduled Interval
 - ◆ ABI (CPT® 93922) following any revascularization procedure.
 - ABI (CPT® 93922) or Duplex ultrasound (CPT® 93926 unilateral study) at each routine follow up is appropriate generally after a history/physical has been performed.
 - Further imaging studies such as CTA or MRA are indicated for worsening symptoms, an abnormal duplex or a significant reduction (>0.15) in the ABI

PVD-7.4: Lower Extremity Artery Aneurysms

- For Iliac artery aneurysm See **PVD-6.3: Iliac Artery Aneurysm (IAA)**
- Femoral artery aneurysm
 - ◆ Initial imaging
 - Ultrasound (CPT® 93925 bilateral study or CPT® 93926 unilateral study).
 - ◆ Surveillance imaging
 - Symptomatic true femoral aneurysms smaller than 2.5 cm in diameter
 - Ultrasound (CPT® 93926 unilateral study) annually
 - Symptomatic true femoral aneurysms larger than 2.5 cm
 - Ultrasound (CPT® 93926 unilateral study) every 6 months
 - ◆ Other imaging
 - CTA Lower extremity or MRA Lower extremity without or with contrast when:
 - Preoperative study for individuals with no plans for invasive angiography.
 - Technically limited or abnormal ultrasound results.
- Popliteal artery aneurysm
 - ◆ Initial imaging
 - Ultrasound (CPT® 93925 bilateral study or CPT® 93926 unilateral study) and Ultrasound to assess for a contralateral popliteal aneurysm and abdominal aortic aneurysm (CPT® 76770 or CPT® 76775).
 - ◆ Surveillance imaging
 - Ultrasound (CPT® 93926 unilateral study) annually.
 - Post interventional functional testing (ABI) (CPT® 93922) may be useful as clinically indicated.
 - ◆ Other imaging
 - CTA or MRA for:
 - Preoperative study for individuals with no plans for invasive angiography.
 - Technically limited or abnormal ultrasound results.

PVD-8: Imaging for Hemodialysis Access

PVD-8.1: Preoperative Arterial Evaluation and Venous Mapping Prior to AV Fistula Creation

46

PVD-8.1: Preoperative Arterial Evaluation and Venous Mapping Prior to AV Fistula Creation

- For vessel mapping prior to AV fistula creation CPT® 93985 or 93986
- In some instances, MRA Upper Extremity (CPT® 73225) may be needed if duplex imaging is equivocal
- Arterial evaluation to assess arterial suitability (size, degree of stenosis and calcification) prior to AV fistula creation may be appropriate
 - ◆ CPT® 93930 or CPT® 93931 can be used to report upper extremity arterial evaluation
 - ◆ Venous mapping to assess venous suitability prior to AV fistula creation may be appropriate
 - CPT® 93970 or CPT® 93971 can be used to report venous mapping
- Indications for Duplex ultrasound (CPT® 93990) of hemodialysis access include but are not limited to:
 - ◆ Individuals with decreased flow rates during hemodialysis.
 - ◆ Development of arm swelling or discomfort after access placement surgery or a hemodialysis session.
 - ◆ Prolonged immaturity of a surgically created AV fistula.
 - ◆ Suspected pseudoaneurysm.
 - ◆ Suspected AV fistula or graft stenosis.
 - ◆ Known or suspected fluid collection adjacent to an AV fistula or graft.
 - ◆ Though it is generally not needed, one Duplex ultrasound (CPT® 93990) can be performed after a surgically created AV fistula for assessment.

References

1. Sidawy AN, Spergel LM, Besarab A, et al. The Society for Vascular Surgery: Clinical practice guidelines for the surgical placement and maintenance of arteriovenous hemodialysis access. *Journal of Vascular Surgery*. 2008;48(5). doi:10.1016/j.jvs.2008.08.042.

PVD-9: Arteriovenous Malformations (AVMs)

See **PEDPVD-2.5: Arteriovenous Malformations (AVMs) and Fistulas** in the Pediatric Peripheral Vascular Disease Imaging Guidelines

PVD-10: This section intentionally left blank

PVD-11: Venous Imaging – General Information

PVD-11.1: Abbreviations and Glossary	50
PVD-11.2: Background Information	51
PVD-11.3: Procedure Coding	53
PVD-11.4: General Guidelines – Imaging	54

PVD-11.1: Abbreviations and Glossary

Abbreviation	Definition
CTV	Computed Tomography Venography
DVT	Deep Venous Thrombosis
EVA	Endovenous ablation – a minimally invasive procedure using heat to obliterate the saphenous vein for the treatment of venous reflux
IVC	Inferior vena cava
May-Thurner's	Syndrome of compression of the left iliac vein via an overlying right common iliac artery. The pulsations of the artery into the vein against the 5 th lumbar vertebra can predispose to DVT
MRV	Magnetic Resonance Venography
Phlebectomy	Removal of a vein usually through a small incision
Post thrombotic syndrome	Constellation of symptoms including chronic edema and pain that develops after a DVT
Sclerotherapy	Injection of an irritant into a vein to obliterate it
SEPS	Sub-fascial endoscopic perforator surgery
SVT	Superficial venous thrombosis
VVI	Venous Valvular Insufficiency – a study utilizing ultrasound to assess for the presence of reflux within the superficial and deep veins of the lower extremity.

PVD-11.2: Background Information

- A current clinical evaluation (within 60 days), including medical treatments, are required prior to considering advanced imaging, which includes:
 - ◆ Relevant history and physical examination including:
 - The affected limb(s), the extent of the edema (calf and/or thigh), pitting or non-pitting. With regard to venous insufficiency, presence or absence of hyperpigmentation or other skin changes, ulcerations if applicable, size of varicosities if present as well as distribution
 - Arterial examination to rule out phlegmasia alba/cerulea dolens which is comprised arterial flow secondary to extensive DVT if applicable
 - Appropriate laboratory studies, for example d-dimer, if applicable
 - Non-advanced imaging modalities, such as a venous duplex or venous valvular insufficiency study (VVI) after symptoms started or worsened
 - ◆ Other meaningful contact (telephone call, electronic mail or messaging) by an established individual can substitute for a face-to-face clinical evaluation.
- Venous disease can be classified into three categories:
 - ◆ Venous-occlusive disease
 - ◆ Venous insufficiency
 - ◆ Venous malformations

Veno-occlusive disease

- Types of thrombotic disease:
 - ◆ Superficial venous thrombosis
 - ◆ Deep venous thrombosis
 - ◆ Iliac vein obstruction, unilateral or bilateral
 - ◆ May-Thurner's syndrome
- Signs/Symptoms of veno-occlusive disease is generally sudden onset of pain and edema in the limb.
- Risk factors include age >40, obesity, pregnancy, prolonged immobility, post-surgery, and malignancy among others.
- Procedures related to veno-occlusive disease include:
 - ◆ Thrombolysis
 - ◆ Thrombectomy
 - ◆ Post Iliac vein angioplasty/stenting

Venous insufficiency

- Types of venous insufficiency:
 - ◆ Superficial and deep venous reflux
 - ◆ Varicose veins
 - ◆ Reticular and spider veins
- Signs/symptoms of venous insufficiency include:
 - ◆ Chronic swelling in the leg that is relieved with elevation
 - ◆ Chronic swelling in the leg that is worse in the evenings

PVD-11.3: Procedure Coding

Venous Studies – Extremities	CPT®
CTV Abdomen and Pelvis involves obtaining images from the diaphragm to just below the inguinal ligament after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.	74174
CTV Pelvis involves obtaining images from the top of the pelvic brim to the upper thighs or just below the inguinal ligament. The venogram portion is performed by obtaining images after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.	72191
MRV Abdomen and Pelvis involves taking images from the diaphragm to just below the inguinal ligament after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.	74185 and 72198
MRV Pelvis involves obtaining images from the top of the pelvic brim to the upper thighs or just below the inguinal ligament. The venogram portion is performed by obtaining images after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.	72198
Non-invasive physiologic studies of extremity veins, complete bilateral study (e.g. Doppler waveform analysis with responses to compression and other maneuvers, phleborheography, impedance plethysmography). This study is rarely performed.	93965
Duplex scan of extremity veins, including responses to compression and other maneuvers; complete bilateral study.	93970
Duplex scan of extremity veins, including responses to compression and other maneuvers; unilateral or limited study.	93971
<ul style="list-style-type: none"> ➤ These codes are used to report studies of lower or upper extremity veins. ➤ A complete bilateral study of the lower extremity veins includes examination of the external iliac veins, common femoral, proximal deep femoral, great saphenous and popliteal veins. Calf veins may also be included. ➤ A complete bilateral study of upper extremity veins includes examination of the subclavian, jugular, axillary, brachial, basilic, and cephalic veins. Forearm veins may also be included. 	
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study	93978
Duplex scan of extremity veins, including responses to compression and other maneuvers; unilateral or limited study.	93979

PVD-11.4: General Guidelines – Imaging

- Venous duplex (CPT® 93970, CPT® 93971) of the limb is the initial imaging of choice
 - ◆ Follow-up duplex imaging (CPT® 93970, CPT® 93971) is not generally indicated to document resolution and should only be obtained for new signs/symptoms or for concerns of propagation of thrombus when the treatment plan would change (Insertion of IVC filter, change of anticoagulation, etc.)
- Imaging studies
 - ◆ Venous duplex (CPT® 93970, CPT® 93971) should visualize the veins, with demonstration of the presence or absence of compressibility and venous flow.
 - ◆ Venous valvular insufficiency studies (CPT® 93970, CPT® 93971) visualize the veins of the lower extremity, assess for reflux (reversal of venous antegrade flow after valve closure) and measure its duration.
 - ◆ CTV or MRV Abdomen and Pelvis with contrast involves taking images from the diaphragm to just below the inguinal ligament after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.

References

1. Ageno W et al. Characteristics and Management of Patients with Venous Thromboembolism: The GARFIELD-VTE Registry. *Thromb Haemost* 2019. 119(2):319-327.
2. Jones WS, Vemulapalli S, Parikh KS et al. Treatment Strategies for Patients with Lower Extremity Chronic Venous Disease (LECVD. Agency for Healthcare Research and Quality (US); 2017 Apr.
3. Gloviczki, P., Comerota, AJ, Dalsing, MD et al. The care of patients with varicose veins and associated chronic venous diseases: Clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum. *Journal of Vascular Surgery*. 2011. 53(16S):1-47.
4. Wolpert, L, Rahmani O, Stein B, et al. Magnetic Resonance Venography in the Diagnosis and Management of May-Thurner syndrome. *Vasc Endovascular Surg*. 2002 Jan-Feb;36(1):51-7.

PVD-12: Acute Limb Swelling

PVD-12.1: Superficial Venous Thrombosis (SVT)	56
PVD-12.2: Acute Deep Venous Thrombosis (DVT)	56
PVD-12.3: Follow-up Imaging of Known DVT	57
PVD-12.4: Follow-up Imaging after Venous Surgery	57

PVD-12.1: Superficial Venous Thrombosis (SVT)

- The diagnosis of superficial venous thrombosis is generally made on the basis of physical examination.
 - ◆ Duplex ultrasound (CPT® 93970, CPT® 93971) is the initial imaging if the diagnosis is equivocal
 - ◆ Follow-up duplex ultrasound (CPT® 93970, CPT® 93971) is indicated only if thrombus in the superficial systems is encroaching onto the deep venous system (saphenofemoral or saphenopopliteal junction)

Background and Supporting Information

- Superficial venous thrombosis (SVT) refers to acute or chronic thrombosis of the superficial veins in both the upper (cephalic and basilic veins) and lower extremities (greater [great] saphenous vein, lesser [small] saphenous vein, gastrocnemius and soleal veins). Treatment: Elevation and warm compresses until pain and swelling subsides.

PVD-12.2: Acute Deep Venous Thrombosis (DVT)

- Duplex ultrasound (CPT® 93970 bilateral study or CPT® 93971 unilateral study) is the initial imaging study for any suspected DVT
 - ◆ Deep venous thrombosis can present as:
 - Symptomatic
 - Swelling
 - Pain
 - Warmth
 - Erythema
 - Pain with dorsiflexion of the foot (Homan's Sign)
 - Or with progression, such as phlegmasia cerulea dolens
 - Risk factors for DVT include age >40, obesity, malignancy, prolonged immobilization, hypercoagulability as well as those outlined in **CH-25: Pulmonary Embolism (PE)** in Chest Imaging Guidelines.
- CTA/CTV Abdomen and Pelvis with contrast can be performed to rule out IVC thrombus secondary to the filter when there is acute bilateral lower extremity swelling in an individual with a history of an IVC filter in place.
- When there is concern for proximal DVT (iliofemoral):
 - ◆ Focused abdominal duplex can generally visualize the iliac veins and IVC to determine the absence or presence of iliac vein thrombus in an individual. If the results are equivocal or indeterminate:
 - CTV or MRV Abdomen and Pelvis with contrast can be performed.
- For proximal DVT's (iliac vein DVT's or in cases of phlegmasia (extensive DVT compromising arterial inflow), thrombectomy (rarely performed) or thrombolysis can be performed.

- If the cause of the DVT is found to be due to May-Thurner, iliac vein angioplasty followed by stenting of the left iliac vein is generally performed. See **PVD-13.3: May-Thurner Syndrome**

Background and Supporting Information

- Deep venous thrombosis is characterized by thrombosis of a deep vein in either the upper (brachial, axillary, subclavian veins) or the lower extremity (peroneal, posterior tibial, popliteal, femoral or iliac veins).

PVD-12.3: Follow-up Imaging of Known DVT

- Duplex ultrasound (CPT® 93970 bilateral study or CPT® 93971 unilateral study) can be repeated in order to rule out proximal extension of a calf vein DVT in those individuals who cannot be anticoagulated, most commonly after recent surgery. Time interval for follow-up study includes:
 - ◆ One week after the initial diagnosis.
 - ◆ Serial imaging (up to 3 studies) over the first three weeks if calf DVT is not treated.
- Imaging during or to terminate long-term anticoagulation therapy to determine venous recanalization is not supported by evidence. Repeat imaging to make decisions on whether or not to continue or terminate anticoagulation is not indicated.

PVD-12.4: Follow-up Imaging after Venous Surgery

- Venous duplex (CPT® 93971 unilateral study) can be obtained of the treated limb to rule out a DVT within seven days of endovenous ablation.
- Follow-up routine imaging is not indicated after other venous procedures including:
 - ◆ Saphenous vein ligation and stripping
 - ◆ Phlebectomy
 - ◆ Sclerotherapy

References

1. Lim W, Gal GL, Bates SM, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Advances*. 2018;2(22):3226-3256. doi:10.1182/bloodadvances.2018024828.
2. American College of Phlebology. Treatment of Superficial Venous Disease of the Lower Leg Guidelines. American Vein & Lymphatic Society. <http://www.phlebology.org/member-resources/publications/treatment-superficial-venous-disease-lower-leg-guidelines>. Published 2016.
3. American College of Phlebology. Duplex Ultrasound Imaging of Lower Extremity Veins in Chronic Venous Disease. American Vein & Lymphatic Society. <http://www.phlebology.org/member-resources/publications/duplex-ultrasound-imaging-lower-extremity-veins-chronic-venous-disease>. Published 2012.
4. Healy DA, Kimura S, Power D, et al. A Systematic Review and Meta-analysis of Thrombotic Events Following Endovenous Thermal Ablation of the Great Saphenous Vein. *European Journal of Vascular and Endovascular Surgery*. 2018;56(3):410-424. doi:10.1016/j.ejvs.2018.05.008.
5. Jacobs CE, Pinzon MM, Orozco J, et al. Deep Venous Thrombosis after Saphenous Endovenous Radiofrequency Ablation: Is it Predictable? *Annals of Vascular Surgery*. 2014;28(3):679-685. doi:10.1016/j.avsg.2013.08.012.

PVD-13: Chronic Limb Swelling Due to Chronic Deep Venous Thrombosis/May Thurner's Syndrome

PVD-13.1: Chronic Deep Venous Thrombosis – General Information	59
PVD-13.2: Post Thrombotic Syndrom	59
PVD-13.3: May-Thurner's Syndrome – General Information	60

PVD-13.1: Chronic Deep Venous Thrombosis – General Information

- In individuals **with** a history of proximal (iliofemoral) DVT who have subsequently developed post thrombotic syndrome, imaging can be performed to evaluate for iliac venous obstruction which can result from incompletely lysed thrombus.
 - ◆ Initial imaging should be Duplex (CPT® 93970 bilateral study or CPT® 93971 unilateral study) followed by either a CT or MR Venogram Abdomen and Pelvis or CT or MR Venogram Pelvis or Venography for treatment planning purposes.
 - ◆ Selected individuals may be a candidate for iliac vein angioplasty/stenting.

Background and Supporting Information

- Chronic deep venous thrombosis is defined as an acute DVT that is greater than 14 days old.
- Individual with incompletely lysed or residual DVT can develop post-thrombotic syndrome that can be characterized as chronic edema, venous stasis changes, pain and in advanced cases venous stasis ulceration.
 - ◆ Incompletely lysed DVT can cause luminal narrowing of the vein restricting venous outflow leading to stenosis or occlusion and /or can lead to valve dysfunction resulting in reflux of venous blood retrograde towards gravity. Both pathologies ultimately lead to chronic edema which can cause chronic pain and venous stasis disease.
 - Imaging for post-thrombotic syndrome is not indicated unless there are signs and symptoms suggestive of a new acute DVT or for preoperative planning for iliac vein/stenting for suspected iliac vein stenosis or occlusion.
 - The mainstay of treatment for chronic deep venous thrombosis is compression stockings.

PVD-13.2: Post Thrombotic Syndrome

- Imaging for post-thrombotic syndrome is indicated when:
 - ◆ There are signs and symptoms suggestive of a new acute DVT (NOT for chronic swelling that has not changed in severity or character)
 - ◆ For preoperative planning for iliac vein/stenting in the setting of known iliac venous obstruction in those with a history of a proximal (iliofemoral) DVT.
- See **PVD-13.3: May-Thurner's syndrome – General Information**
- See **PVD-3.1: Chronic Deep Venous Thrombosis – General Information**
- See **PVD-15: Venous Stasis Ulceration**

PVD-13.3: May-Thurner's Syndrome – General Information

- Duplex (CPT® 93970 bilateral study or CPT® 93971 unilateral study) will confirm the presence of a left common iliac vein DVT but diagnosis is made with advanced imaging such as CTV or MRV Abdomen and Pelvis, Venography or Peri-Procedural Intravascular ultrasound demonstrating compression of the vein.
- Treatment is with iliac vein angioplasty/stenting for both acute and chronic left-sided DVT
- Prophylactic treatment of May-Thurner's syndrome in the absence of acute or chronic DVT **OR** chronic left lower extremity edema and its sequelae such as varicose veins or venous stasis ulcers is **NOT** considered medically necessary.

Background and Supporting Information

- In approximately 25% of people, the right iliac artery overlies the left iliac vein over the fifth lumbar vertebra and its pulsations can compress the vein increasing the risk of DVT in the left extremity

References

1. Partsch H, Flour M, Smith PC, et al. Indications for compression therapy in venous and lymphatic disease. Consensus based on experimental data and scientific evidence. *Int Angiol.* 2008; 27:193-219.
2. Nayak L, Vedantham S. Multifaceted Management of the Postthrombotic Syndrome. *Seminars in Interventional Radiology.* 2012; 29:1.
3. Christof T, Kaltenmeier, Erben Y, et al. Systematic review of May-Thurner syndrome with emphasis on gender differences. *Journal of Vascular Surgery Venous and Lymphatic Disorders.* 2018;6(3):399-407.
4. Rossi FH, Kambara AM, Izukawa NM, et al. Randomized double-blinded study comparing medical treatment versus iliac vein stenting in chronic venous disease. *Journal of Vascular Surgery: Venous and Lymphatic Disorders.* 2018;6(2):183-191.
5. Sermsathanasawadi N, Pruekprasert K, Pitaksantayothin W et al. Prevalence, risk factors, and evaluation of ilio caval obstruction in advanced chronic venous insufficiency. *J Vasc Surg Venous Lymphat Disord.* 2019 May;7(3):441-447.
6. Vedantham S, Kahn SR, Goldhaber SZ, Comerota AJ, et al. Endovascular therapy for advanced post-thrombotic syndrome: Proceedings from a multidisciplinary consensus panel. *Vasc Med.* 2016 Aug;21(4):400-7.

PVD-17.1: Post Iliac Vein Stenting/Angioplasty

- Iliac venous stents can be placed after thrombolysis for DVT associated with May-Thurner's syndrome, DVT associated with extrinsic compression or for post thrombotic iliac obstruction.
 - ◆ Surveillance of iliac venous stents with an arterial duplex (CPT® 93975) can be obtained
 - ◆ For worsening signs or symptoms, including increased edema when stent malfunction is suspected
 - ◆ Postoperatively within the first month, at six months, twelve months and then annually
- Advanced imaging CTV or MRV Abdomen and Pelvis can be obtained for an abnormal or indeterminate duplex

References

1. Washington State Health Care Authority. Selected Treatments for Varicose Veins, A Health Technology Assessment. Prepared by Hayes, Inc. Final Report: April 2017.
2. Proebstle TM, Alm BJ, Göckeritz, et al. Five-year results from the prospective European multicenter cohort study on radiofrequency segmental thermal ablation for incompetent great saphenous veins. *BJS*. 2015; 102: 212-218.
3. Nesbitt C, Bedenis R, Bhattacharya V, et al. Endovenous ablation (radiofrequency and laser) and foam sclerotherapy versus open surgery for great saphenous vein varices (Review). *Cochrane Database of Systematic Reviews*. 2014; Issue 7. Art. No.: CD005624.
4. Weiss RA, Weiss MA, Eimpunth S, et al. Comparative Outcomes of Different Endovenous Thermal Ablation Systems on Great and Small Saphenous Vein Insufficiency: Long-Term Results. *Lasers in Surgery and Medicine*. 2015; 47: 156-160.
5. Brittenden J, Cotton SC, Elders A, et al. Clinical effectiveness and cost-effectiveness of foam sclerotherapy, endovenous laser ablation and surgery for varicose veins: results from the Comparison of LAser, Surgery and foam Sclerotherapy (CLASS) randomised controlled trial. *Health Technol Assess*. 2015; 19(27): 1-342.
6. Biemans AAM, Kockaert M, Akkersdijk GP, et al. Comparing endovenous laser ablation, foam sclerotherapy, and conventional surgery for great saphenous varicose veins. *J Vasc Surg*. 2013; 58: 727-734.
7. van der Velden SK, Biemans AA, De Maeseneer MG, et al. Five-year results of a randomized clinical trial of conventional surgery, endovenous laser ablation and ultrasound-guided foam sclerotherapy in patients with great saphenous varicose veins. *Br J Surg*. 2015; 102(10): 1184-1194.
8. Rasmussen L, Lawaetz M, Serup J, et al. Randomized clinical trial comparing endovenous laser ablation, radiofrequency ablation, foam sclerotherapy and surgical stripping for great saphenous varicose veins with 3-year follow-up. *J Vasc Surg: Venous and Lym Dis*. 2013; 1:349-356.
9. Brittenden J, Cotton SC, Elders A, et al. A randomized trial comparing treatments for varicose veins. *N Engl J Med*. 2014; 371(13): 1218-1227. doi:10.1056/NEJMoa1400781.
10. Venermo M, Saarinen J, Eskelinen E, et al. Randomized clinical trial comparing surgery, endovenous laser ablation and ultrasound-guided foam sclerotherapy for the treatment of great saphenous varicose veins. *BJS*. 2016; 103:1438-1444.