



CLINICAL GUIDELINES

Neck Imaging Policy

Version 20.0.2018
Effective May 17, 2018



eviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Imaging requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2016 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Neck Imaging Guidelines

Abbreviations for Neck Imaging Guidelines	3
Neck-1: General	4
Neck-2: Cerebrovascular and Carotid Disease	6
Neck-3: Dysphagia	7
Neck-4: Esophagus	10
Neck-5: Cervical Lymphadenopathy	12
Neck-6: Neck Masses	14
Neck-7: Malignancies Involving the Neck	17
Neck-8: Recurrent Laryngeal Palsy	18
Neck-9: Thyroid and Parathyroid	19
Neck-10: Trachea and Bronchus	26
Neck-11: Neck Pain	28
Neck-12: Salivary Gland Disorders	31

Abbreviations For Neck Imaging Guidelines

ALS	amyotrophic lateral sclerosis
CT	computed tomography
ENT	Ear, Nose, Throat
FNA	fine needle aspiration
GERD	gastroesophageal reflux disease
GI	gastrointestinal
HIV	human immunodeficiency virus
MRI	magnetic resonance imaging

Neck-1: General

- A current clinical evaluation (within 60 days), which includes a relevant history and physical examination and appropriate laboratory studies and non-advanced imaging modalities, such as plain x-ray or ultrasound, are required prior to considering advanced imaging. Other meaningful contact (telephone call, electronic mail or messaging) by an established patient can substitute for a face-to-face clinical evaluation
- Advanced imaging of the neck covers the following areas:
 - ◆ Skull base (thus a separate CPT® code for head imaging in order to visualize the skull base is not necessary).
 - ◆ Nasopharynx
 - ◆ Upper oral cavity to the head of the clavicle
 - ◆ Parotid glands and the supraclavicular region
- Ultrasound of the soft tissues of the neck including thyroid, parathyroid, parotid and other salivary glands, lymph nodes, cysts, etc. is coded as CPT® 76536. This can be helpful in more ill-defined masses or fullness and differentiating adenopathy from mass or cyst, to define further advanced imaging.
- Neck CT
 - ◆ A Neck CT is usually obtained with contrast only (CPT® 70491).
 - Little significant information is added by performing a Neck CT without and with contrast (CPT® 70492), and there is the risk of added radiation exposure, especially to the thyroid.
 - CT Neck without contrast (CPT® 70490) can be difficult to interpret due to difficulty identifying the blood vessels
 - Exception: Contrast is not generally used when evaluating the trachea with CT. Evaluate salivary duct stones in the appropriate clinical circumstance where intravenous contrast may obscure high attenuation stones
 - Contrast enhanced CT is helpful in the assessment of cervical adenopathy and preoperative planning in the setting of thyroid carcinomas
 - Contrast is recommended as an adjunct to US for patients with clinical suspicion for advanced disease, including invasive primary tumor, or clinically apparent multiple or bulky lymph node involvement
 - Contrast may cause intense and prolonged enhancement of the thyroid gland which interferes with radioactive iodine nuclear medicine studies.
 - Use of IV contrast is an important adjunct because it helps to delineate the anatomic relationship between the primary tumor and metastatic disease. Iodine is generally cleared within four to eight weeks in most patients, so concern about iodine burden from IV contrast causing a clinically significant delay in subsequent whole-body scans (WBSs) or radioactive iodine (RAI) treatment after the imaging followed by surgery is generally unfounded. The benefit gained from improved anatomic imaging generally outweighs any potential risk of a several week delay in RAI imaging or

therapy. Where there is concern, a urinary iodine to creatinine ratio can be measured.

➤ Neck MRI

- ◆ MR Neck is used less frequently than Neck CT.
- ◆ MRI Neck without and with contrast (CPT® 70543) is appropriate if CT suggests the need for further imaging or if ultrasound or CT suggests any of the following:
 - Neurogenic tumor (schwannoma, neurofibroma, glomus tumor, etc.)
 - Vascular malformations
 - Deep neck masses
 - Angiofibromas

References

1. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid*. 2016 Jan;26(1):1-133. Accessed November 16, 2017. <http://online.liebertpub.com/doi/10.1089/thy.2015.0020>.

Neck-2: Cerebrovascular and Carotid Disease

- See these related topics in the Head Imaging Guidelines:
 - ◆ **HD-1.5: General Guidelines – CT and MR Angiography: (CTA and MRA)**
 - ◆ **HD-12: Aneurysm and AVM**
 - ◆ **HD-21: Stroke/TIA**
 - ◆ **HD-22: Cerebral Vasculitis**
 - ◆ **HD-23: Dizziness, Vertigo and Syncope**
 - ◆ **HD-31: Tinnitus**
 - ◆ **HD-32: Eye Disorder**
- See **PVD-3: Cerebrovascular and Carotid Disease** in Peripheral Vascular Disease Imaging Guidelines.

Neck-3: Dysphagia

Neck-3.1: Dysphagia

8

Neck-3.1: Dysphagia

- Mass suspected, either intrinsic or extrinsic, to the esophagus.
 - ◆ Esophagram (Barium Swallow) evaluation is considered the initial study in the evaluation of dysphagia. These results can then lead to further evaluation with:
 - Endoscopy, and/or¹
 - CT Neck with contrast (CPT® 70491), and/or CT Chest with contrast (CPT® 71260), and/or CT Abdomen with contrast (CPT® 74160) (if requested).²
- Dysmotility suspected²
 - ◆ Esophagram and Motility study
- Vascular Ring suspected¹
 - ◆ CT angiography Chest with contrast can be used in the evaluation of suspected vascular ring.
 - ◆ MRI Chest without contrast, or MRI Chest without and with contrast (CPT® 71550 or CPT® 71552), can be performed if vascular ring is suspected.
- Globus Sensation
 - ◆ Findings typical of globus sensation (lump in the throat) need no advanced imaging and have a benign natural history.^{2,3,4}
 - ◆ If the diagnosis is unclear or the clinician cannot adequately visualize the pharynx, after examination and laryngoscopy, the following imaging can be considered:
 - Esophagram
 - Endoscopy and/or
 - X-ray pharynx dynamic and static imaging.^{2,3,4}
 - ◆ Dysphagia, weight loss, odynophagia, throat pain and hoarseness
 - Neck CT with contrast (CPT® 70491)⁴
 - Current or previous upper aerodigestive or esophageal malignancy, or lymphoma
 - Previous neck, esophageal, or gastric surgery
 - Palpable neck abnormality (see: **NECK-6: Neck Masses**)

Practice Notes

- A detailed history of the dysphagia symptoms is important to distinguish neurogenic, pharyngeal and esophageal disorders.
- Dysphagia (difficulty swallowing) can be caused by a wide range of benign and malignant causes that affects the body's ability to move food or liquid from the mouth to the pharynx and into the esophagus.
- A short duration (weeks to months) of rapidly progressive esophageal dysphagia with associated weight loss is highly suggestive of esophageal cancer.
- A study including 148 neck CTs and 104 barium esophagrams done for the evaluation of globus sensation, found no malignancies.^{5, 6}

➤ (See **ONC-9: Esophageal Cancer** in the Oncology Imaging Guidelines).

References

1. Pasha SF, Acosta RD, Chandrasekhara V, et al. The role of endoscopy in the evaluation and management of dysphagia. *Gastrointest Endoscopy*. 2014 Feb;79(2):191-201. Accessed November 16, 2017. [http://www.giejournal.org/article/S0016-5107\(13\)02209-8/fulltext](http://www.giejournal.org/article/S0016-5107(13)02209-8/fulltext).
2. Carucci L, Lalani T, Rosen M, et al. ACR Appropriateness Criteria® Dysphagia. *American College of Radiology (ACR)*. Date of origin: 1998. Last review date: 2013. Accessed November 16, 2017. <https://acsearch.acr.org/docs/69471/Narrative/>.
3. Alhilali L, Seo S-h, Branstetter BF, et al. Yield of neck CT and barium esophagram in patients with globus sensation. *AJNR Am J Neuroradiol*. 2014 Feb;35(2):386-389. Accessed November 16, 2017. <http://www.ajnr.org/content/35/2/386>.
4. Lee BE and Kim GH. Globus pharyngeus: a review of its etiology, diagnosis and treatment. *World J of Gastroenterol*. 2012 May;18(20):2462-2471. Accessed November 16, 2017. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3360444/>.
5. Wippold II F, Cornelius RS, Berger KL, et al. ACR Appropriateness Criteria® Neck mass/adenopathy. *American College of Radiology (ACR)*. Date of origin: 2009. Last review date: 2012. Accessed on November 16, 2017. <https://acsearch.acr.org/docs/69504/Narrative>.
6. Kalva SP, Rybicki FJ, Dill KE, et al. ACR Appropriateness Criteria® Nontraumatic aortic disease. *American College of Radiology (ACR)*. Date of origin: 2013. Accessed on November 16, 2017. <https://acsearch.acr.org/docs/3082597/Narrative>.

Neck-4: Esophagus

Neck-4.1: Imaging

11

Neck-4.1: Imaging

- CT Neck, Chest and/or Abdomen all with contrast (CPT® 70491, CPT® 71260, and/or CPT® 74160) can be performed to evaluate any of the following:
 - ◆ GERD, sliding or paraesophageal hiatal hernias: preoperative planning, (CT Chest and/or Abdomen)¹
 - ◆ Hiatal hernia surgery: for GI Specialist or surgeon treatment/pre-operative planning or signs/symptoms of a potential complication, (CT Chest and Abdomen)
 - ◆ Mallory Weiss tear: suspected after endoscopy, (CT Chest and Abdomen)
 - ◆ Esophageal cancer: biopsy proven
 - ◆ See: **ONC-9:Esophageal Cancer** in the Oncology Imaging Guidelines
 - ◆ Esophageal perforation: suspected (CT Neck and/or Chest and/or Abdomen)
 - ◆ Esophageal diverticulum: Depending on location, any of the CT studies above can be used
- CT Neck and/or Chest or MRI Neck and/or Chest (CPT® 70543 and/or CPT® 71552) **and** endoscopic ultrasound (CPT® 76975) can be used for leiomyoma, depending on the location.
- Suspected foreign body obstructing the esophagus should be evaluated with x-ray.
 - ◆ If x-ray is negative, use contrast study such as esophagram.
 - ◆ A location appropriate CT can be used for further evaluation.
- Any type of esophageal stricture (radiation, peptic, lye, neoplastic, postoperative, drug-induced, Crohn's disease, Schatzki's ring, esophageal web) should be evaluated with esophagram (barium swallow) and endoscopy prior to CT.
 - ◆ If esophagram findings are negative, use CT of appropriate location.¹
- Esophageal motility study (CPT® 78258) can be considered for any of the following:
 - ◆ Dysphagia associated with chest pain and difficulty swallowing both solids and liquids
 - ◆ Gastroesophageal reflux
- Gastroesophageal Reflux Study (CPT® 78262) can be considered for any of the following:
 - ◆ Chronic heartburn
 - ◆ Dysphagia
 - ◆ Family history of Barrett's esophagus or esophageal carcinoma
- Gastric Mucosa Imaging (CPT® 78261)
 - ◆ To evaluate Barrett's esophagus when there is dyspepsia or esophagitis.
- Globus (Pharyngeus, "Hystericus") sensation, Lump in throat (see **Neck-3.1: Dysphagia**)

Practice Notes

- A variety of mechanical and motility lesions occur in the esophagus. Dysphagia is difficulty swallowing; odynophagia is painful swallowing.

Neck-5: Cervical Lymphadenopathy

Neck-5.1: Imaging

13

Neck-5.1: Imaging

- Ultrasound (CPT® 76536) can be considered for any of the following:^{1,2}
 - ◆ Inflammatory, infective, or reactive adenopathy but has failed a 2 week trial of treatment or observation (including antibiotics if appropriate).
 - ◆ To further evaluate an ill-defined mass
 - ◆ High suspicion of malignancy
- Neck CT with contrast (CPT® 70491) can be considered if: ²
 - ◆ Carcinoma found in a lymph node or in an organ known not to be primary (see: **ONC-31.7: Carcinoma of Unknown Primary Site**)
 - ◆ Ultrasound is indeterminate or suspicious for malignancy.

Practice Notes

- Chest x-ray is helpful to identify primary lung disease, involvement of mediastinal lymph nodes or other metastases.
- Inflammatory neck adenopathy is often associated with upper respiratory infection, pharyngitis, dental infection. Occasionally, it is associated with sarcoidosis, toxoplasmosis and HIV.
- Most common causes of neoplastic adenopathy are metastasis from head and neck tumors and lymphoma.

References

1. Ferrer R. Lymphadenopathy: differential diagnosis and evaluation. *Am Fam Physician*. 1998 Oct;58(6):1313-1320. Accessed November 16, 2017. <http://www.aafp.org/afp/1998/1015/p1313.html>.
2. Wippold II F, Cornelius RS, Berger KL, et al. ACR Appropriateness Criteria® Neck mass/adenopathy. *American College of Radiology (ACR)*. Date of origin: 2009. Last review date: 2012. Accessed on November 16, 2017. <https://acsearch.acr.org/docs/69504/Narrative>.

Neck-6: Neck Masses

Neck-6.1: Imaging

15

- See **Pediatric Neck Imaging Guidelines**, if under age 18.

Neck-6.1: Imaging

- Ultrasound (CPT® 76536) is the initial study for:¹
 - ◆ Anterior neck masses
 - ◆ Lateral or posterior neck masses that are tender and have been observed for 2 weeks under physician care and reassessed (generally an acute, infections, or inflammatory mass).
 - ◆ Otherwise ill-defined masses, fullness or asymmetry
- Neck CT with contrast (CPT® 70491) is supported for:¹
 - ◆ Lateral or posterior neck masses that are non-tender and discrete in the adult (> age 18)
 - ◆ History of malignancy that would be primary or metastatic to the neck
 - ◆ Suspected peritonsillar, retropharyngeal or other head and neck abscesses
 - ◆ If sarcoidosis is suspected the Neck CT with contrast (CPT® 70491) should be followed by biopsy.
 - ◆ Preoperative evaluations of any neck mass
- Neck MRI without and with contrast (CPT® 70543) if:¹
 - ◆ CT suggests the need for further imaging.
 - ◆ Ultrasound or CT suggests neurogenic tumor (schwannoma, neurofibroma, glomus tumor, etc.), vascular malformations, deep neck masses and angiofibromas.
- Uncomplicated Pharyngitis or Tonsillitis should undergo conservative therapy including antibiotics, if appropriate. Advanced imaging is not indicated.²

Practice Notes

- Although CT is considered the preferred initial modality in neck mass in adults, the use of US is steadily increasing and should be considered when malignancy is not obvious.
- Most lateral neck masses are enlarged lymph nodes.
- Malignancy is a greater possibility in adults that are heavy drinkers and smokers.
- ENT evaluation can be helpful in determining the need for advanced imaging.
- Although CT and MRI can have characteristic appearances for certain entities, biopsy and histological diagnosis are the only way to obtain a definitive diagnosis.

References

1. Wippold II F, Cornelius RS, Berger KL, et al. ACR Appropriateness Criteria® Neck mass/adenopathy. *Am Coll Radiol (ACR)*. Date or origin: 2009. Last review date: 2012. Accessed on November 16, 2017. <https://acsearch.acr.org/docs/69504/Narrative>.
2. Shulman ST, Bisno AL, Clegg HW, et al.
3. Clinical practice guideline for the diagnosis and management of group A streptococcal pharyngitis: 2012 update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2010 Nov;55(10):e86-e102. Accessed November 16, 2017. <https://academic.oup.com/cid/article/55/10/e86/321183>.

Neck-7: Malignancies Involving the Neck

- See the following in the Oncology Imaging Guidelines:
 - ◆ **ONC-3: Squamous Cell Carcinomas - Head and Neck**
 - ◆ **ONC-4: Salivary Gland Cancers**
 - ◆ **ONC-6: Thyroid Cancer**
 - ◆ **ONC-9: Esophageal Cancer**
 - ◆ **ONC-27: Non-Hodgkin Lymphoma**
 - ◆ **ONC-28: Hodgkin Lymphoma**

Neck-8: Recurrent Laryngeal Palsy

- See: **HD-7: Recurrent Laryngeal Palsy** in the Head Imaging Guidelines

Neck-9: Thyroid and Parathyroid

Neck-9.1: Thyroid Nodule	20
Neck-9.2: Hyperthyroidism	22
Neck-9.3: Parathyroid Imaging	22

Neck-9.1: Thyroid Nodule

- Ultrasound (US) of the Neck (CPT® 76536) is the appropriate initial study for anterior neck masses, including goiter and thyroid^{2,6} mass(es) (see: **Neck-6.1: Imaging**). US is also appropriate for evaluation of thyroid nodules in the following clinical scenarios:
 - ◆ Normal or High serum thyrotropin (TSH)^{1,3,6}
 - ◆ Low TSH and nuclear scan shows non-functioning nodule.^{1,6,8}
 - ◆ Incidentally found on CT, MRI, or PET (focal activity).^{2,3,6}
 - ◆ Nodules ≤ 1 cm with very low suspicion US pattern including spongiform pattern and pure cysts do not require repeat US.⁶
 - ◆ For more suspicious or larger nodules, if Fine Needle Aspiration (FNA) is not performed or was not diagnostic for malignancy, US can be repeated:
 - If US features are highly suspicious: repeat US every 6 months for up to 24 months.
 - If US features are of low to intermediate suspicion: repeat US at 12 and 24 months.
 - If nodule is stable after 24 months, follow-up ultrasound exams (CPT® 76536) can be performed every 3 to 5 years for interval surveillance.¹²
- FNA should be considered for thyroid nodules using the American Thyroid Association criteria listed below. Note that FNA procedures do not require prior authorization.⁶

Sonographic Pattern	US features	Estimated risk % of Malignancy	FNA size cutoff (largest dimension)
High Suspicion	Solid hypoechoic nodule or solid hypoechoic component of a partially cystic nodule with one or more of the following features: Irregular margins (infiltrative, microlobulated), microcalcifications, taller than wide shape, rim calcifications with small extrusive soft tissue component, evidence of ETE	>70-90	Recommend FNA at ≥ 1cm
Intermediate Suspicion	Hypoechoic solid nodule with smooth margins without microcalcifications, ETE, or taller than wide shape	10-20	Recommend FNA at ≥ 1cm
Low Suspicion	Isoechoic or hypoechoic solid nodule, or partially cystic nodule with eccentric solid areas, without microcalcifications, irregular margin or ETE, or taller than wide shape	5-10	Recommend FNA at ≥ 1.5cm
Very Low Suspicion	Spongiform or partially cystic nodule without any of the sonographic features described in low, intermediate, or high suspicion patterns	<3	Consider FNA at ≥ 2cm Observation without FNA is also a reasonable option
Benign	Purely cystic nodules (no solid component)	<1	No biopsy

(Source: [2015 American Thyroid Management Guideline for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer](#))

See also **TI-RADS**¹²

- Nuclear medicine thyroid scan (CPT® 78013 or CPT® 78014) is considered for any of the following:
 - ◆ Non-diagnostic or indeterminate FNA of thyroid nodule, e.g. follicular lesion of undetermined significance to see if hot (functioning) nodule that may be benign vs cold nodule for treatment planning.³
 - ◆ Substernal goiter with any one of the following:
 - Dyspnea (including exertional)
 - Wheezing or stridor
 - Cough
 - Dysphagia
 - ◆ Evaluate eligibility for radioiodine therapy³
 - ◆ Select nodules to biopsy in multinodular goiter even if TSH not low^{1,6}
- Nuclear medicine thyroid scan (CPT® 78013 or CPT® 78014) is considered if low serum thyrotropin (TSH) and any of the following:
 - ◆ Single or multiple thyroid nodules^{3,6}
 - ◆ Suspicion of ectopic thyroid tissue³
 - ◆ Presence of thyroid nodule in the setting of Grave's disease (to rule out cold nodule).³
- CT Neck with contrast (CPT® 70491) or CT Neck without contrast (CPT® 70490), or MRI Neck without and with contrast (CPT® 70543). MRI and CT **are not** indicated for routine thyroid nodule evaluation and should only be considered for:
 - ◆ Evaluation of size/substernal extension of a nodular goiter³
 - ◆ Airway compression³
 - ◆ Presence of pathologic lymph nodes in cervical regions not visualized on ultrasound³
 - ◆ In selected cases for nodules with aggressive features for more accurate pre-operative staging^{3,6}
 - ◆ Clinically suspected advanced disease, including invasive primary tumor^{3,6}
 - ◆ Inconclusive US with suspected thyroid cancer⁹
 - ◆ Preoperative planning for any thyroid disease
- A thyroid nodule detected for the first time during pregnancy should be managed in the same way as in non-pregnant patients, except for avoiding the use of radioactive agents for diagnostic and therapeutic purposes³

Practice Notes

- The basis of thyroid nodule management is the use of ultrasonography (US), thyrotropin (TSH, formerly thyroid-stimulating hormone) assay, and FNA biopsy, together with clinical findings.
- Patient Features Suggesting Increased Risk for Thyroid Malignancy.
 - ◆ History of head and neck irradiation
 - ◆ Family history of medullary thyroid carcinoma, multiple endocrine neoplasia type 2, or papillary thyroid carcinoma
 - ◆ Age < 14 or > 70 years

- ◆ Male sex
- ◆ Growth of the nodule
- ◆ Firm or hard nodule consistency
- ◆ Cervical adenopathy
- ◆ Fixed nodule
- ◆ Persistent dysphonia, dysphagia, or dyspnea
- Iodinated CT contrast may interfere with diagnostic nuclear medicine thyroid scans (scintigraphy) and radioiodine treatment.
- There is insufficient evidence supporting the use of PET to distinguish indeterminate thyroid nodules that are benign from those that are malignant.
- ¹⁸FDG-PET imaging is not routinely recommended for the evaluation of thyroid nodules with indeterminate cytology. Routine preoperative ¹⁸FDG-PET scanning is not recommended.

Neck-9.2: Hyperthyroidism

- Hyperthyroidism suspected^{4,7}
 - ◆ Thyroid Uptake Study (CPT® 78012 or CPT® 78014) if one of the following:
 - TSH below normal range and elevated free T4 and/or free T3, OR
 - Subclinical hyperthyroidism with TSH < 0.1 mU/L and normal free T4 and free T3.

Neck-9.3: Parathyroid Imaging

- Primary Hyperparathyroidism suspected
 - ◆ Parathyroid Planar Imaging (CPT® 78070), Parathyroid Planar Imaging with SPECT (CPT® 78071), Parathyroid Planar Imaging with SPECT and CT (CPT® 78072) or Ultrasound (CPT® 76536) if either:
 - Elevated serum calcium and elevated serum parathyroid hormone level.
 - Serum calcium 1 mg/dL more over lab normal value
 - ◆ CT or MRI Neck without and with contrast (CPT® 70492 or CPT® 70543):
 - Very high calcium (≥ 13) suggesting parathyroid carcinoma
 - Preoperative localization including 4D Neck CT without and with contrast (CPT® 70492 or 77293).⁷
 - Recurrent or persistent hyperparathyroidism following neck exploration (MRI preferred).
 - ◆ Chest CT with contrast may be indicated in rare circumstances in the evaluation of ectopic mediastinal parathyroid adenomas.⁶

Practice Notes

- A thyroid nodule is distinct either on palpation or radiologically (incidentaloma). Nonpalpable nodules have the same risk of cancer as palpable. Nodules > 1 cm are evaluated, while smaller nodules are generally evaluated if suspicious, associated with adenopathy or a history of radiation or cancer exists.

- Ultrasound is not used to screen: 1) the general population, 2) patients with normal thyroid on palpation with a low risk of thyroid cancer, 3) patients with hyperthyroidism, 4) patients with hypothyroidism or 5) patients with thyroiditis. Conversely, US can be considered in patients who have no symptoms but are high risk as a result of: history of head and neck irradiation, total body irradiation for bone marrow transplant, exposure to fallout from radiation during childhood or adolescence, family history, thyroid cancer syndromes such as MEN2, medullary or papillary thyroid cancer, Cowden's disease, familial adenomatous polyposis, Carney complex, Werner syndrome/progeria.
- Radionuclide thyroid scan can be considered to evaluate nodules when hyperthyroidism is present, for surveillance of thyroid cancer, or to detect non-palpable nodules. This scan is not useful for other nodules since hyper functioning nodules rarely harbor malignancy. Thyroid nodules > 4 cm may be considered for thyroid lobectomy due to a high incidence of both false negative FNA biopsies and malignancy (26%).
- FNA may be repeated after an initial non-diagnostic cytology result, because repeat FNA with US guidance will yield a diagnostic cytology specimen in 75% of solid nodules and 50% of cystic nodules. However, up to 7% of nodules continue to yield non-diagnostic cytology results despite repeated biopsies and may be malignant at the time of surgery.
- Thyroid nodules may be stratified as to risk of thyroid cancer based on sonographic findings of microcalcification, hypervascularity on Doppler ultrasound, solid or cystic nature of mass and margins of mass.
- Incidental focal FDG-PET uptake often corresponds to a clinically relevant thyroid nodule and ultrasound is recommended; incidentally noted diffuse thyroid FDG-PET uptake most often corresponds to inflammatory uptake, however, ultrasound should be done to ensure that there is no evidence of clinically relevant nodularity.
- Elastography provides information about nodule stiffness that is complementary to gray scale ultrasound findings in nodules with indeterminate cytology or ultrasound findings. It should not be used as a substitute for gray scale ultrasound.
- Use of ultrasound contrast medium is not recommended for the diagnostic evaluation of thyroid nodules and its current use is restricted to definition of size and limits of necrotic zones after minimally invasive nodule ablation techniques.

References

Thyroid

1. Cooper DS, Doherty GM, Haugen BR et al. Revised American Thyroid Association management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid*. 2009 November;19(11):1167-1214. Accessed November 16, 2017. <https://www.ncbi.nlm.nih.gov/pubmed/19860577>.
2. Hoang JK, Langer JE, Middleton WD, et al. Managing incidental thyroid nodules detected on imaging: white paper of the ACR incidental thyroid findings committee. *J Am Coll Radiol*. 2015 Feb;12(2):143-150. Accessed November 16, 2017 [http://www.jacr.org/article/S1546-1440\(14\)00627-9/fulltext](http://www.jacr.org/article/S1546-1440(14)00627-9/fulltext).
3. Gharib H, Papini E, Garber JR, et al. American Association Of Clinical Endocrinologists, American College Of Endocrinology, And Associazione Medici Endocrinologi medical guidelines for clinical practice for the diagnosis and management of thyroid nodules--2016 update. *Endocr Pract*. 2016 May;22(Suppl 1):1-60. Accessed November 16, 2017. <http://journals.aace.com/doi/10.4158/EP161208.GL>.
4. Burch H, Cooper D, Garber J, et al. Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists. *Endocr Pract*. 2011 May-Jun;17(3):456-520. Accessed November 16, 2017. <http://journals.aace.com/doi/pdf/10.4158/EP.17.3.456>.
5. Guidelines and Protocols Advisory Committee, approved by the British Columbia Medical Association, and adopted by the Medical Services Commission. Thyroid function tests in the diagnoses and monitoring of adults. Effective January 1, 2010. Accessed November 16, 2017. <https://www2.gov.bc.ca/gov/content/health/practitioner-professional-resources/bc-guidelines/thyroid-testing>.
6. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid*. 2016 Jan;26(1):1-133. Accessed November 16, 2017. <http://online.liebertpub.com/doi/10.1089/thy.2015.0020>.
7. Donangelo I and Suh SY. Subclinical hyperthyroidism: when to consider treatment. *Am Fam Physician*. 2017 Jun;95(11):710-716. Accessed November 16, 2017 <http://www.aafp.org/afp/2017/0601/p710.html>.
8. National Comprehensive Cancer Network (NCCN) Guidelines Version 2.2017 – May 17, 2017. Thyroid Carcinoma. Accessed November 16, 2017. https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines™) for Thyroid Carcinoma 2.2017. ©2017 National Comprehensive Cancer Network, Inc. All rights reserved. The NCCN Guidelines™ and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org.
9. Goiter. *American Thyroid Association*. Accessed November 16, 2017. <http://www.thyroid.org/goiter/>.
10. Mohebbati A and Shaha A. Imaging techniques in parathyroid surgery for primary hyperparathyroidism. *Am J Otolaryngol*. 2012 Jul-Aug;33(4):457-468. Accessed November 16, 2017. <https://dx.doi.org/10.1016%2Fj.amjoto.2011.10.010>.
11. Khan AA, Hanley DA, Rizzoli R, et al. Primary hyperparathyroidism: review and recommendations on evaluation, diagnosis, and management. A Canadian and international consensus. *Osteoporos Int*. Accessed November 16, 2017. <https://link.springer.com/article/10.1007%2Fs00198-016-3716-2>.
12. Grant EG, Tessler FN, Hoang JK, et al. Thyroid ultrasound reporting lexicon: white paper of the ACR thyroid imaging, reporting and data system (TIRADS) committee. *J Am Coll Radiology* 2015 Dec;12(12) Part A:1272-1279. Accessed November 16, 2017. [http://www.jacr.org/article/S1546-1440\(15\)00684-5/abstract](http://www.jacr.org/article/S1546-1440(15)00684-5/abstract).

Parathyroid

1. Carty SE, Worsey MJ, Virji MA, et al. Concise parathyroidectomy: the impact of preoperative SPECT ^{99m}Tc sestamibi scanning and intraoperative quick parathormone assay. *Surgery*. 1997 Dec;122(6):1107-1116. Accessed November 16, 2017. [http://www.surgjournal.com/article/S0039-6060\(97\)90215-4/pdf](http://www.surgjournal.com/article/S0039-6060(97)90215-4/pdf).
2. Arici C, Cheah WK, Ituarte P, et al. Can localization studies be used to direct focused parathyroid operations? *Surgery*. 2001 Jun;129(6):720-729. Accessed November 16, 2017. [http://www.surgjournal.com/article/S0039-6060\(01\)41394-8/abstract](http://www.surgjournal.com/article/S0039-6060(01)41394-8/abstract).
3. Udelsman R. Six hundred fifty-six consecutive explorations for primary hyperparathyroidism. *Ann Surg*. 2002 May;235(5):665-672. Accessed November 16, 2017. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1422492/#_ffn_sectitle.
4. Eslamy HK and Ziessman HA. Parathyroid scintigraphy in patients with primary hyperparathyroidism: ^{99m}Tc sestamibi SPECT and SPECT/CT. *RadioGraphics*. 2008 Sep-Oct;28(5):1461-1477. Accessed November 16, 2017. <http://pubs.rsna.org/doi/pdf/10.1148/rq.285075055>.
5. Lubitz CC, Stephen AE, Hodin RA, et al. Preoperative localization strategies for primary hyperparathyroidism: an economic analysis. *Ann Surg Oncol*. 2012 Dec;19(13):4202-4209. Accessed November 16, 2017. <https://link.springer.com/article/10.1245%2Fs10434-012-2512-2>.
6. Mortenson MM, Evans DB, Lee JE, et al. Parathyroid exploration in the reoperative neck: improved preoperative localization with 4D-computed tomography. *J Am Coll Surg*. 2008 May;206(5):888-895. Accessed November 16, 2017. [http://www.journalacs.org/article/S1072-7515\(08\)00032-X/pdf](http://www.journalacs.org/article/S1072-7515(08)00032-X/pdf).
7. Kukar M, Platz TA, Schaffner TJ, et al. The use of modified four-dimensional computed tomography in patients with primary hyperparathyroidism: an argument for the abandonment of routine sestamibi single-positron emission computed tomography (SPECT). *Ann Surg Oncol*. 2015 Jan;22(1):139-145. Accessed November 16, 2017. <https://link.springer.com/article/10.1245/s10434-014-3940-y>.

Neck-10: Trachea and Bronchus

Neck-10.1: Imaging

27

Neck-10.1: Imaging

- Plain x-rays of the neck and chest and bronchoscopy are the initial imaging studies for evaluating patients with suspected tracheal and visualized bronchial pathology. Bronchoscopy can further evaluate the distal (endo) bronchial tree.
 - ◆ Suspected tracheal disease can be identified by inspiratory stridor and a characteristic flow-volume loop of PFTs.¹
- Neck CT with contrast (CPT® 70491) or without contrast (CPT® 70490) and/or Chest CT with contrast (CPT® 71260) or without contrast (CPT® 71250) can be performed to further evaluate abnormalities, which include tracheal or bronchial tumor, foreign bodies, or persistent segmental or lobar lung collapse seen on other imaging studies based on the physician's preference.^{1,2}
- Expiratory HRCT (CPT® 71250) is indicated in patients with obstructive physiology tracheomalacia and can also be useful in the evaluation of interstitial lung disease.¹
- Trachea or bronchial "inspissation" without an abnormality described above, is not a risk for malignancy.³

References

1. Dyer DS, Mohammed T-LH, Kirsch J, et al. ACR Appropriateness Criteria® Chronic dyspnea: suspected pulmonary origin. *Am Coll Radiol (ACR)*. Date of origin: 1995. Last review date: 2012. Accessed November 16, 2017. <https://acsearch.acr.org/docs/69448/Narrative/>.
2. Obusez EC, Jamjoom L, Kirsch J, et al. Computed tomography correlation of airway disease with bronchoscopy: part I--nonneoplastic large airway diseases. *Curr Probl Diagn Radiol*. 2014 Sep-Oct;43(5):268-277. Accessed November 16, 2017. [http://www.cpdjournal.com/article/S0363-0188\(14\)00038-3/fulltext](http://www.cpdjournal.com/article/S0363-0188(14)00038-3/fulltext).
3. Gould MK, Donington J, Lynch WR, et al. Evaluation of individuals with pulmonary nodules: when is it lung cancer? Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2013 May;143(5):e93S-e120S. Accessed November 16, 2017. [http://journal.chestnet.org/article/S0012-3692\(13\)60291-3/fulltext](http://journal.chestnet.org/article/S0012-3692(13)60291-3/fulltext).

Neck-11: Neck Pain

Neck-11.1: Neck Pain (Cervical)	29
Neck-11.2: Torticollis and Dystonia	29

Neck-11.1: Neck Pain (Cervical)

- Neck pain is usually related to a specific process including pharyngitis, radiculopathy, adenopathy, mass, carotid dissection and torticollis, and therefore found elsewhere in these guidelines.¹
- For the evaluation of neck pain or other symptoms which may involve the cervical spine, including myelopathy and cervical radiculopathy ¹ see: **Spine Imaging Guidelines**

Neck-11.2: Torticollis and Dystonia

Newborn Infant:

- Ultrasound of the Neck is the initial study to determine if congenital muscular torticollis is present
 - ◆ Positive→No further imaging is needed since the diagnosis is defined
 - ◆ Negative→CT Neck with contrast (CPT® 70491) or MRI Neck without and with contrast (CPT® 70543) to potentially identify other etiologies

Older Child (beyond infancy) or Adult

- For trauma, CT Neck with contrast (CPT® 70491) and/or CT Cervical Spine without contrast (CPT® 72125) is the initial study to identify fracture or mal-alignment
- For no trauma, CT Neck with contrast (CPT® 70491), and/or MRI Cervical Spine without contrast (CPT® 72141), or CT Cervical Spine without contrast (CPT® 72125) is the initial study to locate a soft tissue or neurological cause
 - ◆ Positive→Further advanced imaging is not required if CT Neck or CT Cervical Spine has identified local cause
 - ◆ Negative→MRI Brain without and with contrast (CPT® 70553) to exclude CNS cause

Practice Notes

- Torticollis or cervical dystonia is an abnormal twisting of the neck with head rotated or twisted. Its causes are many and may be congenital or acquired and caused by trauma, infection/inflammation, neoplasm and those less defined and idiopathic. It occurs more frequently in children and on the right side (75%).
- Retropharyngeal space abscess could be associated with torticollis because child would not move neck freely.

References

1. Newman JS, Weissman BN, Angevine PD, et al. ACR Appropriateness Criteria® Chronic neck pain. *Am Coll Radiol (ACR)*. Date of origin: 1998. Last review date: 2013. Accessed November 16, 2017. <https://acsearch.acr.org/docs/69426/Narrative/>.
2. Anderson PA and Montesano PX. Morphology and treatment of occipital condyle fractures. *Spine*. 1988 Jul;13(7):731-736. Accessed November 16, 2017. http://journals.lww.com/spinejournal/Abstract/1988/07000/Morphology_and_Treatment_of_Occipital_Condyle.4.aspx.
3. Ballock RT and Song KM. The prevalence of nonmuscular causes of torticollis in children. *J Pediatr Orthop*. 1996 Jul-Aug;16(4):500-504. Accessed November 16, 2017. http://journals.lww.com/pedorthopaedics/Abstract/1996/07000/The_Prevalence_of_Nonmuscular_Causes_of.16.aspx.
4. Castillo M, Albernaz VS, Mukherji SK, et al. Imaging of Bezold's abscess. *AJR Am J Roentgenol*. 1998 Dec;171(6):1491-1495. Accessed November 16, 2017. <http://www.ajronline.org/doi/pdf/10.2214/ajr.171.6.9843276>.
5. Federico F, Lucivero V, Simone IL, et al. Proton MR spectroscopy in idiopathic spasmodic torticollis. *Neuroradiology*. 2001 Jul;43(7):532-536. Accessed November 16, 2017. <https://link.springer.com/article/10.1007/s002340000532>.
6. Fielding JW and Hawkins RJ. Atlanto-axial rotatory fixation. (fixed rotatory subluxation of the atlanto-axial joint). *J Bone Joint Surg Am*. 1977 Jan;59(1):37-44. Accessed November 16, 2017. http://journals.lww.com/jbjsjournal/Abstract/1977/59010/Atlanto_axial_rotatory_fixation_Fixed_rotatory.5.aspx.
7. Kraus R, Han BK, Babcock DS, et al. Sonography of neck masses in children. *AJR Am J Roentgenol*. 1986 Mar;146(3):609-613. Accessed November 16, 2017. <http://www.ajronline.org/doi/pdf/10.2214/ajr.146.3.609>.
8. Roche CJ, O'Malley M, Dorgan J, et al. A pictorial review of atlantoaxial rotatory fixation: key points for the radiologist. *Clinical Radiology*. 2001 Dec;56(12):947-958. Accessed November 16, 2017. [http://www.clinicalradiologyonline.net/article/S0009-9260\(01\)90679-8/fulltext](http://www.clinicalradiologyonline.net/article/S0009-9260(01)90679-8/fulltext).
9. Tracy MR, Dormans JP, Kusumi K. Klippel-Feil syndrome: clinical features and current understanding of etiology. *Clin Orthop Relat Res*. 2004 Jul;424:183-190. Accessed November 16, 2017. <https://www.ncbi.nlm.nih.gov/pubmed/15241163>.

Neck-12: Salivary Gland Disorders

- Xerostomia (Dry Mouth)
 - ◆ Salivary Gland Nuclear Imaging (one of CPT® 78230, CPT® 78231, or CPT® 78232) can be considered for any one of the following:
 - Dry mouth and either:
 - Sjögren's syndrome
 - Sialadenitis
 - History of head or neck radiation therapy
 - History of cerebral palsy
 - Parotid mass to allow preoperative diagnosis of Warthin's tumor
- Salivary Gland Stones:¹
 - ◆ For suspected salivary duct or gland stone, CT of the Neck without contrast (CPT® 70490) or CT of the Neck without and with contrast (CPT® 70492) or CT of the Maxillofacial area without and with contrast (usually CPT® 70488) or MRI Neck without and with contrast (CPT® 70543).
 - ◆ Sialography (contrast dye injection) under fluoroscopy, may be performed to rule out a stone, with post sialography CT (CPT® 70486), **or** post sialography MRI (CPT® 70540).
- Parotid Mass
 - ◆ Any one of the following can be approved:²
 - MRI Orbits/Face/Neck without and with contrast (CPT® 70543)
 - CT Neck with contrast (CPT® 70491)
 - CT Neck without contrast (CPT® 70490)
 - In addition to one of the above:
 - Salivary Gland Nuclear Imaging (CPT® 78230, CPT® 78231, or CPT® 78232) is indicated If salivary gland stone is suspected, CT of the maxillofacial area without and with contrast (usually CPT® 70488) or Neck MRI without and with contrast (CPT® 70543) can be considered in place of Neck CT.

References

1. Wilson KF, Meier JD, and Ward PD. Salivary gland disorders. *Am Fam Physician*. 2014 Jun;89(11):882-888. Accessed November 16, 2017. <http://www.aafp.org/afp/2014/0601/p882.html>.
2. Wippold II F, Cornelius RS, Berger KL, et al. ACR Appropriateness Criteria® Neck mass/adenopathy. *American College of Radiology (ACR)*. Date of origin: 2009. Last review date: 2012. Accessed on November 16, 2017. <https://acsearch.acr.org/docs/69504/Narrative>.