

Acthar® Gel (repository corticotropin injection)



When requesting Acthar® Gel (repository corticotropin injection), the individual requiring treatment must be diagnosed with infantile spasms and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-approved Indication

Acthar Gel (repository corticotropin injection) is indicated as monotherapy for the treatment of infantile spasms (West syndrome) in infants and children under 2 years of age.

(Note: All other FDA-approved indications will be reviewed by medical directors.)

Coverage Guidelines

Infantile Spasms

The individual must meet **BOTH** of the following criteria for approval:

- Is less than 2 years of age; AND
- Acthar is prescribed by a physician who has consulted with or specializes in neurology.

Approval duration: 1 month

Dosing Recommendation

The recommended dose is 150 U/m² divided into twice daily intramuscular injections of 75 U/m². After 2 weeks of treatment, dosing should be gradually tapered and discontinued over a 2-week period.

References

1. H.P. Acthar® Gel injection for subcutaneous and intramuscular use [prescribing information]. Bedminster, NJ: Mallinckrodt; October 2021.
2. Go CY, Mackay MT, Weiss SK, et al. Evidence-based guideline update: medical treatment of infantile spasms: Report of the guideline development subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2012;78:1974-1980.
3. Pellock JM, Hrachovy R, Shinnar S, et al. Infantile spasms: a US consensus report. *Epilepsia*. 2010;51(10):2175-2189.
4. National Multiple Sclerosis Society. Expert Opinion Paper. National Clinical Advisory Board of the National Multiple Sclerosis Society. Treatment Recommendations for Physicians. Recommendations Regarding Corticosteroids in the Management of Multiple Sclerosis. Available at: http://www.nationalmssociety.org/NationalMSSociety/media/MSNationalFiles/Brochures/ExpOp_Steroids.pdf. Accessed on November 30, 2021.
5. Thompson AJ, Kennard C, Swash M, et al. Relative efficacy of intravenous methylprednisolone and ACTH in the treatment of acute relapses in MS. *Neurology*. 1989;39:969-971.
6. Kidney Disease: Improving Global Outcomes (KDIGO) glomerulonephritis Work Group. KDIGO Clinical Practice Guidelines for Glomerulonephritis. *Kidney Int. Suppl.* 2012;2:139-274. Available at <https://kdigo.org/wp-content/uploads/2017/02/KDIGO-2012-GN-Guideline-English.pdf>. Accessed on November 30, 2021.

7. Berkovich R, Fernandez M, Subhani D. Monthly pulse adrenocorticotrophic hormone (ACTH) or methylprednisolone therapy for long-term treatment of multiple sclerosis as an add-on therapy to beta-interferons: interim results from a pilot study [poster P576]. Presented at: European Committee for Treatment and Research in Multiple Sclerosis (ECTRIMS) (with ACTRIMS [Americas Committee for Treatment and Research in Multiple Sclerosis]). Amsterdam, The Netherlands, October 19-22, 2011.
8. Tumlin JA, Galphin CM, Rovin BH. Advanced diabetic nephropathy with nephrotic range proteinuria: a pilot study of the long-term efficacy of subcutaneous ACTH gel on proteinuria, progression of CKD, and urinary levels of VEGF and MCP-1. *J Diabetes Res*. 2013;2013:489869.
9. Chen Y, Schieppati A, Chen X, et al. Immunosuppressive treatment for idiopathic membranous nephropathy in adults with nephrotic syndrome. *Cochrane Database Syst Rev*. 2014 Oct 16;10:DC004293.
10. Bomback AS, Tumlin JA, Baranski J, et al. Treatment of nephrotic syndrome with adrenocorticotrophic hormone (ACTH) gel. *Drug Des Devel Ther*. 2011;5:147-153.
11. Bomback AS, Canetta PA, Beck LH, et al. Treatment of resistant glomerular diseases with adrenocorticotrophic hormone gel: a prospective trial. *Am J Nephrol*. 2012;36:58-67.
12. Hladunewich MA, Dattran D, Beck LH, et al. A pilot study to determine the dose and effectiveness of adrenocorticotrophin hormone (H.P. Acthar® Gel) in nephrotic syndrome due to idiopathic membranous nephropathy. *Nephrol Dial Transplant*. 2014;29(8):1570-1577.
13. Berg AL, Arnadottir M. ACTH-induced improvement in the nephrotic syndrome in patients with a variety of diagnoses. *Nephrol Dial Transplant*. 2004;19:1305-1307.
14. Ponticelli C, Passerini P, Salvadori M, et al. A randomized pilot trial comparing methylprednisolone plus a cytotoxic agent versus synthetic adrenocorticotrophic hormone in idiopathic membranous nephropathy. *Am J Kidney Dis*. 2006;47(2):233-240.
15. Lorusso P, Bottai A, Mangione E, et al. Low-dose synthetic adrenocorticotrophic hormone-analog therapy for nephrotic patients: results from a single-center pilot study. *Int J Nephrol Renovasc Dis*. 2015;8:7-12.
16. Madan A, Mijovic-Das S, Stankovic A, et al. Acthar gel in the treatment of nephrotic syndrome: a multicenter retrospective case series. *BMC Nephrology*. 2016;17:37.
17. Wang CS, Travers C, McCracken C, et al. Adrenocorticotrophic hormone for childhood nephrotic syndrome. The ATLANTIS randomized trial. *Clin J Am Soc Nephrol*. 2018;13:1859-1865.
18. Hartung DM, Johnston K, Deodhar A, et al. Repository corticotropin versus glucocorticoid for nephrotic syndrome: when will we see the evidence? *Am J Kidney Dis*. 2019;74(2):256-262.
19. Fiechtner JJ, Montroy T. Treatment of moderately to severely active systemic lupus erythematosus with adrenocorticotrophic hormone: a single-site, open-label trial. *Lupus*. 2014;23:905-912.
20. Tumulin J, Galphin C, Santos R, Rovin B. Safety and efficacy of combination ACTHar gel and tacrolimus in treatment-resistant focal segmental glomerulosclerosis and membranous glomerulopathy. *Kidney Int Rep*. 2017;2:924-932.
21. Li X, Golubovsky J, Hui-Yuen J, et al. Adrenocorticotrophic hormone gel in the treatment of systemic lupus erythematosus: a retrospective study of patients. *F1000Res*. 2015 Oct 23;4:1103 [eCollection 2015].
22. Filippone EJ, Dopson SJ, Rivers DM, et al. Adrenocorticotrophic hormone analog use for podocytopathies. *Int Med Case Rep J*. 2016;9:125-133.
23. Furie R, Mitran M, Zhao E, et al. Efficacy and tolerability of repository corticotropin injection in patients with persistently active SLE: results of a phase 4, randomized, controlled pilot study. *Lupus Sci Med*. 2016;3:e000180.
24. Hogan J, Bomback AS, Mehta K, et al. Treatment of idiopathic FSGS with adrenocorticotrophic hormone gel. *Clin J Am Soc Nephrol*. 2013;8:2072-2081.
25. Hahn BH, McMahon MA, Wilkinson A, et al. American College of Rheumatology guidelines for screening, treatment, and management of lupus nephritis. *Arthritis Care Res (Hoboken)*. 2012;64(6):797-808.
26. Levine T. Treating refractory dermatomyositis or polymyositis with adrenocorticotrophic hormone gel: a retrospective case series. *Drug Des Devel Ther*. 2012;6:133-139.
27. Ernste FC, Reed AM. Idiopathic inflammatory myopathies: current trends in pathogenesis, clinical features, and up-to-date treatment recommendations. *Mayo Clin Proc*. 2013;88(1):83-105.