

Acthar® Gel (repository corticotropin injection)



When requesting Acthar® Gel (repository corticotropin injection), the individual requiring treatment must be diagnosed with infantile spasms and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-approved Indication

Acthar Gel (repository corticotropin injection) is indicated as monotherapy for the treatment of infantile spasms (West syndrome) in infants and children under 2 years of age.

(Note: All other FDA-approved indications will be reviewed by medical directors.)

Coverage Guidelines

Infantile Spasms

The individual must meet **BOTH** of the following criteria for approval:

- Is less than 2 years of age; AND
- Acthar is prescribed by a physician who has consulted with or specializes in neurology.

Approval duration: 1 month

Dosing Recommendation

The recommended dose is 150 U/m² divided into twice daily intramuscular injections of 75 U/m². After 2 weeks of treatment, dosing should be gradually tapered and discontinued over a 2-week period.

References

1. Acthar® Gel injection for subcutaneous and intramuscular use [prescribing information]. Bedminster, NJ: Mallinckrodt; October 2021.
2. Tran KA, Harrod C, Bourdette DN, et al. Characterization of the clinical evidence supporting repository corticotropin injection for FDA-approved indications. A scoping review. *JAMA Intern Med.* 2022;182(2):206-217.
3. Go CY, Mackay MT, Weiss SK, et al. Evidence-based guideline update: medical treatment of infantile spasms: Report of the guideline development subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology.* 2012;78:1974-1980.
4. Pellock JM, Hrachovy R, Shinnar S, et al. Infantile spasms: a US consensus report. *Epilepsia.* 2010;51(10):2175-2189.
5. KDIGO 2021 clinical practice guidelines for the management of glomerular diseases. *Kidney Int.* 2021;100:S1-S276. Available at: <https://www.kidney-international.org/action/showPdf?pii=S0085-2538%2821%2900562-7>. Accessed on April 8, 2022.
6. National Multiple Sclerosis Society. Expert Opinion Paper. National Clinical Advisory Board of the National Multiple Sclerosis Society. Treatment Recommendations for Physicians. Recommendations Regarding Corticosteroids in the Management of Multiple Sclerosis. Available at: http://www.nationalmssociety.org/NationalMSSociety/media/MSNationalFiles/Brochures/ExpOp_Steroids.pdf. Accessed on April 8, 2022.

7. Thompson AJ, Kennard C, Swash M, et al. Relative efficacy of intravenous methylprednisolone and ACTH in the treatment of acute relapses in MS. *Neurology*. 1989;39:969-971.
8. American College of Rheumatology. Clinical Practice Guidelines. Available at: [Clinical Practice Guidelines \(rheumatology.org\)](https://www.rheumatology.org). Accessed on April 15, 2022.
9. Fitzgerald JD, Dalbeth N, Mikuls T, et al. 2020 American College of Rheumatology guideline for the management of gout. *Arthritis Care Res (Hoboken)*. 2020;72(6):744-760.
10. Baughman RP, Valeyre D, Korsten P, et al. ERS clinical practice guidelines on treatment of sarcoidosis. *Eur Respir J*. 2021;58(6):2004079.
11. Ward MM, Deodhar A, Gensler LS, et al. 2019 update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis Rheumatol*. 2019;71(10):1599-1613.
12. Ward MM, Deodhar A, Akl EA, et al. American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network 2015 recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Care Res*. 2016;68:151-166.
13. Oldroyd AGS, Lilleker JB, Amin T, et al. British Society for Rheumatology guideline on management of paediatric, adolescent and adult patients with idiopathic inflammatory myopathy. *Rheumatology*. 2022 March 31. [Online ahead of print].
14. American Diabetes Association Professional Practice Committee. Chronic Kidney Disease and Risk Management: Standards of Medical Care in Diabetes-2022. *Diabetes Care*. 2022;45(1):S175-S184.
15. Onel KB, Horton DB, Lovell DJ, et al. 2021 American College of Rheumatology Guideline for the treatment of juvenile idiopathic arthritis: therapeutic approaches for oligoarthritis, temporomandibular joint arthritis, and systemic juvenile idiopathic arthritis. *Arthritis Rheumatol*. 2022;74(4):553-569.
16. Fanouriakis A, Kostopoulou M, Cheema K, et al. 2019 Update of the Joint European League Against Rheumatism and European Renal Association-European Dialysis and Transplant Association (EULAR/ERA-EDT) recommendations for the management of lupus nephritis. *Ann Rheum Dis*. 2020;79:713-732.
17. Crane AB, Sharon Y, Chu DS. Use of adrenocorticotrophic hormone in ophthalmology. *J Ocul Pharmacol Ther*. 2020;36(9):661-667.
18. Wirta D, McLaurin E, Ousler G, et al. Repository corticotropin injection (Acthar® gel) for refractory severe noninfectious keratitis: efficacy and safety from a phase 4, multicenter, open-label study. *Ophthalmol Ther*. 2021;10:1077-1092.
19. Anesi SD, Chang PY, Maleki A, et al. Treatment of noninfectious retinal vasculitis using subcutaneous repository corticotropin injection. *J Ophthalmic Vis Res*. 2021;16:219-233.
20. Singh JA, Guyatt G, Ordie A, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol*. 2019;71(1):5-32.
21. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Care Res (Hoboken)*. 2021;73(7):924-939.

Acthar® Gel (repository corticotropin injection)