

Soliris® (eculizumab)

When requesting Soliris® (eculizumab), the individual requiring treatment must be diagnosed with an FDA-approved indication and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-approved Indications

Soliris is a complement inhibitor indicated for:

- The treatment of patients with paroxysmal nocturnal hemoglobinuria (PNH) to reduce hemolysis.
- The treatment of patients with atypical hemolytic uremic syndrome (aHUS) to inhibit complement-mediated thrombotic microangiopathy.
- The treatment of adult patients with generalized myasthenia gravis (gMG) who are anti-acetylcholine receptor (AChR) antibody positive.
- The treatment of neuromyelitis optica spectrum disorder (NMOSD) in adults who are anti-aquaporin-4 (AQP4) antibody positive.

Coverage Guidelines

Atypical Hemolytic Uremic Syndrome (aHUS)

The individual must meet **all** of the following criteria for authorization:

- The individual does not have Shiga toxin *Escherichia coli* related hemolytic uremic syndrome; AND
- Soliris is prescribed by or in consultation with a hematologist and/or a nephrologist.

Approval duration: 12 months

Paroxysmal Nocturnal Hemoglobinuria (PNH)

The individual must meet **all** of the following criteria for initial authorization:

- The diagnosis of PNH has been confirmed by peripheral blood flow cytometry results showing the absence or deficiency of glycosylphosphatidylinositol (GPI)-anchored proteins on at least 2 cell lineages;
- The individual is 18 years of age or older; AND
- Soliris is prescribed by or in consultation with a hematologist.

For re-authorization:

- The individual is continuing to receive benefits from Soliris therapy (e.g., stabilization of hemoglobin levels, decreased transfusion requirement/transfusion independence, reduction in hemolysis).

Approval duration (initial): 6 months

Approval duration (renewal): 12 months

Generalized Myasthenia Gravis (gMG)

The individual must meet **all** of the following criteria for initial authorization:

- Has a confirmed diagnosis of anti-acetylcholine receptor (AChR) antibody positive generalized myasthenia gravis;
- Is 18 years of age or older;
- Is currently receiving pyridostigmine unless failed, contraindicated or intolerant to pyridostigmine;
- Is currently receiving 2 different immunosuppressive therapies for over 1 year unless failed, contraindicated, or intolerant to 2 different immunosuppressive therapies (e.g., azathioprine, cyclosporine, mycophenolate mofetil, methotrexate, tacrolimus, cyclophosphamide);
- Has evidence of unresolved symptoms of generalized myasthenia gravis resulting in the discontinuation of physical activity (e.g., difficulty swallowing, difficulty breathing, double vision, impairment of mobility); AND
- Soliris is prescribed by or in consultation with a neurologist.

For re-authorization:

- The individual is continuing to receive benefits from Soliris therapy (e.g., reductions in exacerbations of myasthenia gravis; improvement in speech, swallowing, mobility, and respiratory functions).

Approval duration (initial): 6 months

Approval duration (renewal): 12 months

Neuromyelitis Optica Spectrum Disorder (NMOSD)

The individual must meet **all** of the following criteria for authorization:

- Diagnosis has been confirmed by blood serum test for anti-aquaporin-4 antibody positive NMOSD;
- Is 18 years of age or older;
- Has previously tried one of the following therapies: azathioprine, corticosteroid, mycophenolate mofetil, or rituximab; AND
- Soliris is prescribed by or in consultation with a neurologist.

Approval duration: 12 months

Dosing Recommendations

Soliris is only administered as an intravenous infusion.

Paroxysmal nocturnal hemoglobinuria (PHN)

- 600 mg weekly for the first 4 weeks, followed by
- 900 mg for the fifth dose 1 week later, then
- 900 mg every 2 weeks thereafter.

Atypical hemolytic uremia syndrome (aHUS) (for patients 18 years of age and older)

- 900 mg weekly for the first 4 weeks, followed by
- 1200 mg for the fifth dose 1 week later, then
- 1200 mg every 2 weeks thereafter.

Atypical hemolytic uremia syndrome (aHUS) (for patients less than 18 years of age)

Body weight	Induction	Maintenance
40 kg and over	900 mg weekly x 4 doses	1200 mg at week 5; then 1200 mg every 2 weeks
30 kg to less than 40 kg	600 mg weekly x 2 doses	900 mg at week 3; then 900 mg every 2 weeks
20 kg to less than 30 kg	600 mg weekly x 2 doses	600 mg at week 3; then 600 mg every 2 weeks
10 kg to less than 20 kg	600 mg weekly x 1 dose	300 mg at week 2; then 300 mg every 2 weeks
5 kg to less than 10 kg	300 mg weekly x 1 dose	300 mg at week 2; then 300 mg every 3 weeks

Generalized myasthenia gravis (gMG)

- 900 mg weekly for the first 4 weeks, followed by
- 1200 mg for the fifth dose 1 week later, then
- 1200 mg every 2 weeks thereafter.

Neuromyelitis optica spectrum disorder (NMOSD)

- 900mg weekly for first 4 weeks, followed by
- 1200mg for the fifth dose 1 week later, then
- 1200mg every 2 weeks thereafter.

References

1. Soliris® injection [prescribing information]. New Haven, CT: Alexion Pharmaceuticals, Inc.; June 2019.
2. Campistol JM, Arias M, Ariceta G, et al. An update for atypical haemolytic uraemic syndrome: diagnosis and treatment. A consensus document. *Nefrologia*. 2015;35:421–447.
3. Genetics Home Reference. Atypical hemolytic-uremic syndrome. National Institutes of Health (NIH). Available at: <https://ghr.nlm.nih.gov/condition/atypical-hemolytic-uremic-syndrome#sourcesforpage>. Accessed on May 23, 2019.
4. National Institute of Neurological Disorders and Stroke (NINDS). Myasthenia Gravis Fact Sheet. National Institutes of Health (NIH) Publication No. 17-768. Publication last updated: May 14, 2019. Available at: <https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Fact-Sheets/Myasthenia-Gravis-Fact-Sheet>. Accessed on May 23, 2019.
5. Sanders DB, Wolfe GI, Benatar M, et al. International consensus guidance for management of myasthenia gravis. *Neurology*. 2016;87:419–425.
6. Brodsky RA. Paroxysmal nocturnal hemoglobinuria. *Blood*. 2014;124(18):2804–2811.
7. Taylor CM, Machin S, Wigmore SJ, et al. Clinical Practice Guidelines for the management of atypical Haemolytic Uraemic Syndrome in the United Kingdom. *Br J Haematol*. 2010;148(1):37-47.
8. Loirat C, Fakhouri F, Ariceta G, et al. An international consensus approach to the management of atypical hemolytic uremic syndrome in children. *Pediatr Nephrol*. 2016;31(1):15-39.
9. National Organization for Rare Disorders. Neuromyelitis Optica Spectrum Disorder. Available at: <https://rarediseases.org/rare-diseases/neuromyelitis-optica/>. Accessed July 25, 2019.

10. National Institute of Health, U.S. National Library of Medicine. Genetics Home Reference. Neuromyelitis optica. Available at: <https://ghr.nlm.nih.gov/condition/neuromyelitis-optica#genes>. Accessed July 25, 2019.
11. Eculizumab (Soliris®). In: DRUGDEX® System Micromedex. Greenwood Village, CO: Thompson Reuters (Healthcare) Inc. [Updated; November 06, 2019; Accessed December 04, 2019].