

Spinraza® (nusinersen)

When requesting Spinraza® (nusinersen), the individual requiring treatment must be diagnosed with an FDA-approved indication and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-approved Indication

Spinraza is indicated for the treatment of spinal muscular atrophy (SMA).

Coverage Guidelines

Treatment of spinal muscular atrophy (SMA)

An individual must meet **all** of the following criteria for initial approval:

- Has had a genetic test confirming the diagnosis of SMA with bi-allelic mutations in the *survival motor neuron 1 (SMN1)* gene reported as at least one of the following: homozygous deletion, homozygous mutation, or compound heterozygous mutation;
- Has 2 or 3 survival motor neuron 2 (SMN2) gene copies OR has 4 or more SMN2 gene copies with symptoms consistent with Types 1, 2, or 3 SMA;
- Has not received Zolgensma in the past;
- The following laboratory tests will be evaluated prior to the administration of Spinraza: prothrombin time and/or activated partial thromboplastin time, platelet count, and quantitative spot urine protein test; AND
- Spinraza is prescribed by or in consultation with a physician who specializes in the management of spinal muscular atrophy and/or neuromuscular disorders.

For reauthorization, an individual must meet all of the above **and** the following criteria:

- Has responded to Spinraza therapy (e.g., improvement, achievement, and/or maintenance in motor milestones, reduced need for respiratory support, prevention of permanent assisted ventilation).

Approval duration (initial): 3 months

Approval duration (renewal): 4 months (one dose)

Dosing Recommendation

Spinraza is administered intrathecally.

The recommended dose is 12 mg (5mL) per administration. Initiate Spinraza treatment with 4 loading doses. The first three loading doses should be administered at 14-day

intervals. The 4th loading dose should be administered 30 days after the 3rd dose. A maintenance dose of 12mg should be administered once every 4 months thereafter.

References

1. Spinraza® injection for intrathecal use [prescribing information]. Cambridge, MA: Biogen; June 2019.
2. Darras B, Markowitz J, Monani U, De Vivo D. Chapter 8-Spinal Muscular Atrophies. In: Vivo BT, ed. *Neuromuscular Disorders of Infancy, Childhood, and Adolescence (Second Edition)*. San Diego: Academic Press; 2015;117-145.
3. Carre A, Empey C. Review of spinal muscular atrophy (SMA) for prenatal and pediatric genetic counselors. *J Genet Couns*. 2016;25:32-43.
4. Nash LA, Burns JK, Chardon JW, et al. Spinal muscular atrophy: more than a disease of motor neurons. *Curr Mol Med*. 2016;16(9):779-792.
5. Darras BT. Spinal muscular atrophies. *Pediatr Clin N Am*. 2015;62:743-766.
6. Kolb SJ, Kissel JT. Spinal muscular atrophy. *Neurol Clin*. 2015;33(4):831-846.
7. Chiriboga CA. Nusinersen for the treatment of spinal muscular atrophy. *Expert Rev Neurother*. 2017;17(10):955-962.
8. Finkel RS, Mercuri E, Darras BT, et al, for the ENDEAR Group. Nusinersen versus sham control in infantile-onset spinal muscular atrophy. *N Engl J Med*. 2017;377(18):1723-1732.
9. Mercuri E, Darras BT, Chiriboga JW, et al, for the CHERISH Study Group. Nusinersen versus sham control in later-onset spinal muscular atrophy. *N Engl J Med*. 2018;378:625-635.
10. Day JW, Swoboda KJ, Darras BT, et al [poster]. Nusinersen experience in teenagers and young adults with spinal muscular atrophy. Presented at: the Muscular Dystrophy Association Clinical Conference; Arlington, VA; March 11-14, 2018.
11. De Vivo DC, Bertini E, Swoboda KJ, et al. Nusinersen initiated in infants during the presymptomatic stage of spinal muscular atrophy: interim efficacy and safety results from the Phase 2 NURTURE study. *Neuromuscular Dis*. 2019;29:842-856.
12. Finkel RS, Castro D, Farrar MA, et al. Interim report on the safety and efficacy of longer-term treatment with nusinersen in infantile-onset spinal muscular atrophy (SMA): updated results from the SHINE study. Presented at: the American Academy of Neurology 71st annual meeting. Philadelphia, PA; May 4-11, 2019.
13. Darras BT, Chiriboga CA, Iannaccone ST, et al. Nusinersen in later-onset spinal muscular atrophy. *Neurology*. 2019;92(1):e2492-e2506.
14. Michelson D, Cifaloni E, Ashwal S, et al. Evidence in focus: nusinersen use in spinal muscular atrophy. Report of the Guideline Development, Dissemination, and implementation Subcommittee of the American Academy of Neurology. *Neurology*. 2018;91(20):923-933.
15. Hoy SM. Nusinersen: a review of 5q spinal muscular atrophy. *CNS Drugs*. 2018;32:689-696.
16. Shieh PB, Acsadi G, Mueller-Felber W, et al. Safety and efficacy of nusinersen in infants/children with spinal muscular atrophy (SMA): Part 1 of the Phase 2 EMBRACE Study. Presented at: the American Academy of Neurology 70th annual meeting. Los Angeles, CA; April 21-27, 2018.
17. Pane M, Palermo C, Messina S, et al, for the Italian EAP Working Group. Nusinersen in type 1 SMA infants, children and young adults: preliminary results on motor function. *Neuromuscular Disord*. 2018;28(7):582-585.
18. Wurster CD, Winter B, Bollinsky K, et al. Intrathecal administration of nusinersen in adolescent and adult SMA type 2 and 3 patients. *J Neurol*. 2019;266(1):183-194.
19. Stolte B, Totzeck A, Kizina K, et al. Feasibility and safety of intrathecal treatment with nusinersen in adult patients with spinal muscular atrophy. *Ther Adv Neurol Disord*. 2018;11:1-9.
20. Glascock J, Sampson J, Haidet-Phillips A, et al. Treatment algorithm for infants diagnosed with spinal muscular atrophy through newborn screening. *J Neuromusc Dis*. 2018;5:145-158.
21. Mercuri E, Rinkel RS, Muntoni F, et al, for the SMA Care Group. Diagnosis and management of spinal muscular atrophy: Part 1: Recommendations for diagnosis, rehabilitation, orthopedic and nutritional care. *Neuromuscul Disord*. 2018;28(2):103-115. Available at: <https://air.unimi.it/retrieve/handle/2434/702816/1424447/main.pdf>. Accessed on May 19, 2020.
22. Prior TW, Leach ME, Finanger E. Spinal Muscular Atrophy. 2000 Feb 24 [Updated 2019 Nov 14]. In: Adam MP, Ardinger, HH, Pagon RA, et al, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK1352/>. Accessed on May 19, 2020.
23. Hagenacker T, Wurster CD, Funther R, et al. Nusinersen in adults with 5q spinal muscular atrophy: a non-interventional, multicenter, observational cohort study. *Lancet Neurol*. 2020;19:317-325.
24. Jochmann E, Steinbach, R, Jochmann T, et al. Experiences from treating seven adult 5q spinal muscular atrophy patients with nusinersen. *Ther Adv Neurol Disord*. 2020;13:1-11.

25. Osmanovic A, Ranxha G, Kumpe M, et al. Treatment expectations and patient-reported outcomes of nusinersen therapy in adult spinal muscular atrophy. *J Neurol*. 2020 May 2. [Online ahead of print].
26. Walter MC, Wenninger S, Thiele S, et al. *J Neuromuscul Dis*. Safety and treatment effects of nusinersen in longstanding adult 5q-SMA Type 3 – a prospective observational study. *J Neuromuscul Dis*. 2019;6(4):453-465.
27. Veerapandiyam A, Eichinger K, Guntrum D, et al. Nusinersen for older patients with spinal muscular atrophy: a real-world clinical setting experience. *Muscle Nerve*. 2020;61:218-242.
28. Nusinersen (Spinraza®). In: DRUGDEX® System Micromedex. Greenwood Village, CO: Thompson Reuters (Healthcare) Inc. [Updated; December 6, 2019; Accessed June 11, 2020].