

Subcutaneous Immune Globulins (Cutaquig, Cuvitru, Hizentra, Hyqvia, Xembify)

When requesting Subcutaneous Immune Globulin (Cutaquig, Cuvitru, Hizentra, Hyqvia, Xembify), the individual requiring treatment must be diagnosed with an FDA-approved indication and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-Approved Indications

Subcutaneous immune globulin (SCIG) injection is indicated for the treatment of:

- Primary humoral immunodeficiency including but not limited to, congenital agammaglobulinemia, common variable immunodeficiency, x-linked agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies.
- Chronic inflammatory demyelinating polyneuropathy (CIDP) as maintenance therapy to prevent relapse of neuromuscular disability and impairment

Coverage Guidelines

Primary humoral immunodeficiency

Must have one of the following subtypes of primary immunodeficiencies:

- Common variable immunodeficiency (CVID)
- X-linked agammaglobulinemia
- Congenital agammaglobulinemia
- Other agammaglobulinemia due to absence of B-cells
- Severe combined immunodeficiency (SCID)
- Wiskott-Aldrich syndrome
- Hyper-Immunoglobulin M (IgM) syndrome
- IgG level lower than 250 mg/dL
- DiGeorge syndrome
- Ataxia telangiectasis,
- Other combined immunodeficiencies with significant hypogammaglobulinemia
- Unspecified hypogammaglobulinemia (or unspecified IgG deficiency)

For an initial authorization request:

- Is 18 years of age or older (*applies to Hyqvia only*)
- The requested product is prescribed by or in consultation with one of the following physician specialist who treats patients with PID: allergist, immunologist, otolaryngologist (ear, nose, and throat [ENT] physician), pulmonologist, or infectious disease physician

- Has frequent and severe infections (*applies to other combined immunodeficiencies with significant hypogammaglobulinemia or antibody production defect indication only*)
- For CVID, unspecified hypogammaglobulinemia, or other immunodeficiencies with significant hypogammaglobulinemia indications:
 - Is 2 years of age or older (*excludes Hyqvia*)
 - The total serum IgG level is below the normal range (age-adjusted and according to the normal reference range for the reporting laboratory) ; AND
 - The patient has impaired antibody response (i.e., failure to produce antibodies to specific antigen); OR
 - The patient has recurrent infections; OR
 - The patient has an IgG subclass deficiency or a diagnosis of selective antibody deficiency (SAD) with impaired antibody response AND recurrent infections; OR
 - The delay caused by pre-vaccination and post-vaccination antibody measurement can be deleterious to the individual's health in regards to impaired antibody response

For re-authorization request:

- The patient is diagnosed with a primary immunodeficiency and is continuing to receive benefit from the product (e.g., increased IgG levels, preventing or controlling infections)

Chronic inflammatory demyelinating polyneuropathy (excludes Hyqvia):

For an initial authorization request:

- Is 18 years of age or older
- The request product is prescribed by or in consultation with a neurologist
- Electrodiagnostic studies to support the diagnosis of CIDP

For a re-authorization request:

- Must have a clinically significant improvement in neurological symptoms (e.g., improvement in disability; nerve conduction study results improved or stabilized; physical examination showing improvement in neurological symptoms, strength, and sensation)

Approval duration (initial and renewal): 12 months

Dosing Recommendations

Cutaquig

Dosage for patients switching from IVIG

- Initial weekly dose = previous IVIG dose (in grams) x 1.40/No. of weeks between IVIG

Dosage for patients switching from SCIG:

- The weekly dose of Cuvitru is recommended to be the same as the weekly dose of prior SCIG treatment

Cuvitru

Dosage for patients switching from IVIG or Hyqvia:

- Initial weekly dose = previous IVIG or Hyqvia dose (in grams)/No. of weeks between IVIG or Hyqvia doses x 1.30
- Frequent dosing (2-7 times per week): divide the calculated weekly dose by the desired number of times per week
- Biweekly dosing: multiply the calculated weekly dose by 2

Dosage for patients switching from SCIG:

- The weekly dose of Cuvitru is recommended to be the same as the weekly dose of prior SCIG treatment
- Frequent dosing (2-7 times per week): divide the calculated weekly dose by the desired number of times per week
- Biweekly dosing: multiply the calculated weekly dose by 2

Hyqvia

Initial treatment interval/dosage ramp-up schedule:

- For patients previously on another immune globulin treatment, administer the first dose approximately one week after the last infusion of their previous treatment
- Increase the dose and frequency from a 1-week dose to a 3- or 4-week dose (see table below)

Week	Infusion Number	Dose Interval	Example for 30 grams per 4 weeks
1	1 st infusion	1-week-dose	7.5 grams
2	2 nd infusion	2-week-dose	15 grams
3	No infusion		
4	3 rd infusion	3-week-dose	22.5 grams
5	No infusion		
6	No infusion		
7	4 th infusion (if required)	4-week-dose	30 grams

Dosage for patients switching from IVIG:

- Administer Hyqvia at the same dose and frequency as the previous intravenous treatment, after the initial dose ramp-up

Dosage for patient naïve to SCIG treatment or switching from another SCIG:

- Administer Hyqvia at 300 to 600 mg/kg at 3 to 4 week interval, after initial ramp-up

Hizentra

For primary immunodeficiency:

- Hizentra can be administered at regular intervals from daily up to every 2 weeks
- For weekly or frequent dosing, start Hizentra 1 week after the last IVIG infusion or SCIG infusion
- For biweekly dosing, start Hizentra 1 or 2 weeks after the last IVIG infusion or 1 week after the last weekly SCIG infusion

Dosage for patients switching to Hizentra from IVIG:

- For weekly dosing: start Hizentra 1 week after last IVIG infusion
Initial weekly dose = previous IVIG dose (in grams)/No. of weeks between IVIG doses x 1.37
- For biweekly dosing: Multiply the calculated Hizentra weekly dose by 2
- For frequent dosing (2 to 7 times per week): Divide the calculated weekly dose by the desired number of times per week

Dosage for patients switching to Hizentra from SCIG:

- The previously weekly SCIG dose should be maintained
- For biweekly dosing: Multiply the previous weekly dose by 2
- For frequent dosing (2 to 7 times per week): Divide the previous weekly dose by the desired number of times per week.

For chronic inflammatory demyelinating polyneuropathy (CIDP):

- Initiate Hizentra 1 week after the last IVIG infusion
- The recommended dose is 0.2 g/kg (1 mL/kg) per week. In the clinical study, a dose of 0.4 g/kg (2 mL/kg) per week was also safe and effective to prevent CIDP relapse

Xembify

Dosage for patients switching to Xembify from IVIG:

- Initial weekly dose (grams) = prior IVIG dose (in grams) x 1.37/No. of weeks between IVIG doses
- Frequent dosing (2-7 times per week) = divide the calculated weekly dose by the desired number of times per week

Dosage for patients switching to Xembify from SCIG:

- Weekly dose (grams) should be the same as the weekly dose or prior SCIG treatment (grams)

References

1. Gammagard Liquid 10% [prescribing information]. Lexington, MA: Baxalta US; July 2017.
2. Gammaked™ 10% injection [prescribing information]. Fort Lee, NJ: Kedrion Biopharma, Inc. (manufactured by Grifols Therapeutics, Inc, Research Triangle Park, NC); June 2018.
3. Gamunex®-C 10% liquid [prescribing information]. Research Triangle Park, NC: Grifols USA, LLC (manufactured by Grifols Therapeutics Inc, Research Triangle Park, NC); June 2018.

4. Hizentra® for subcutaneous infusion [prescribing information]. Kankakee, IL: CSL Behring LLC (manufactured by CSL Behring AG, Bern, Switzerland); March 2018.
5. HyQvia immune globulin infusion 10% with recombinant human hyaluronidase [prescribing information]. Lexington, MA: Baxalta US Inc.; January 2019.
6. McLean HQ, Fiebelkorn AP, Temte JL, Wallace GS; Centers for Disease Control and Prevention. Prevention of measles, rubella, congenital rubella syndrome, and mumps, 2013: summary recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep*. 2013;62:1-34.
7. Xembify 20% solution [prescribing information]. Research Triangle Park, NC: Grifols Therapeutics LLC; July 2019.
8. Ochs HD, Gupta S, Kiessling P, et al; Subcutaneous IgG Study Group. Safety and efficacy of self-administered subcutaneous immunoglobulin in patients with primary immunodeficiency diseases. *J Clin Immunol*. 2006;26:265-273.
9. Gardulf A, Nicolay U, Asensio O, et al. Rapid subcutaneous IgG replacement therapy is effective and safe in children and adults with primary immunodeficiencies--a prospective, multi-national study. *J Clin Immunol*. 2006;26:177-185.
10. Berger M, Murphy E, Riley P and Bergman GE and the VIRTUE Trial Investigators. Improved quality of life, immunoglobulin G levels, and infection rates in patients with primary immunodeficiency diseases during self-treatment with subcutaneous immunoglobulin G. *South Med J*. 2010;103:856-863.
11. McCormak PL. Immune globulin subcutaneous (human) 20%: in primary immunodeficiency disorders. *Drugs*. 2012;72:1087-1097.
12. Cuvitru subcutaneous 20% solution [prescribing information]. Westlake Village, CA: Baxalta US Inc.; September 2016.
13. Cutaquiq 16.5% solution [prescribing information]. Hoboken, NJ: Octapharma USA, Inc.; December 2018.
14. Berger M. Subcutaneous administration of IgG. *Immunol Allergy Clin North Am*. 2008;28:779-802, viii.
15. Perez EE, Orange JS, Bonilla F, et al. Update on the use of immunoglobulin in human disease: A review of evidence. *J Allergy Clin Immunol*. 2017;139(3S):S1-S46. Available at: [http://www.jacionline.org/article/S0091-6749\(16\)31141-1/pdf](http://www.jacionline.org/article/S0091-6749(16)31141-1/pdf). Accessed on July 16, 2019.
16. Donofrio PD, Berger A, Brannagan TH, et al. Consensus statement: the use of intravenous immunoglobulin in the treatment of neuromuscular conditions report of the AANEM ad hoc committee. *Muscle Nerve*. 2009;40:890-900.
17. Patwa HS, Chaudhry V, Katzberg H, et al. Evidence-based guideline: Intravenous immunoglobulin in the treatment of neuromuscular disorders. Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 2012;78(13):1009-15. Available at <http://n.neurology.org/content/78/13/1009.full>. Accessed on July 16, 2019.
18. Harbo T, Andersen H, Hess A, et al. Subcutaneous versus intravenous immunoglobulin in multifocal motor neuropathy: a randomized, single-blinded cross-over trial. *Eur J Neurol*. 2009;16:631-638.
19. Eftimov F, Vermeulen M, de Hann RJ, et al. Subcutaneous immunoglobulin therapy for multifocal motor neuropathy. *J Peripher Nerv Syst*. 2009;14:93-100.
20. Harbo T, Anderson H and Jakobsen J. Long-term therapy with high doses of subcutaneous immunoglobulin in multifocal motor neuropathy. *Neurology*. 2010;75:1377-1380.
21. Vacca A, Melaccio A, Sportelli A, et al. Subcutaneous immunoglobulins in patients with multiple myeloma and secondary hypogammaglobulinemia: a randomized trial. *Clin Immunol*. 2018;191:110-15.
22. Kovacs E, Dando K, Nagy-Vince M, et al. Long-term management of refractory myasthenia gravis with subcutaneous immunoglobulin. *Ther Adv Neurol Disord*. 2017; 10(11):363-366.
23. van Schaik IN, Bril V, van Geloven N, et al; PATH study group. Subcutaneous immunoglobulin for maintenance treatment in chronic inflammatory demyelinating polyneuropathy (PATH): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Neurol*. 2018;17(1):35-46.
24. Bonilla FA, Khan DA, Ballas ZK, et al. Practice parameter for the diagnosis and management of primary immunodeficiency. *J Allergy Clin Immunol*. 2015;136:1186-1205.e1-78.
25. Immune Deficiency Foundation. Patient & Family Handbook for Primary Immunodeficiency diseases. 5th edition. 2015. Available at: <http://primaryimmune.org/wp-content/uploads/2016/03/IDF-Patient-Family-Handbook-5th-Edition-2015-Reprint-Web.pdf>. Accessed on July 16, 2019.
26. Bonilla FA, Barlan I, Chapel H, et al. International Consensus Document (ICON): Common variable immunodeficiency disorders. *J Allergy Clin Immunol Pract*. 2016;4(1): 38-59
27. Wasserman RL, Melamed I, Stein MR, et al; the IGSC, 10% with rHuPH20 Study Group. Recombinant human hyaluronidase-facilitated subcutaneous infusion of human immunoglobulins for primary immunodeficiency. *J Allergy Clin Immunol*. 2012;130:951-957
28. Sanford M. Human immunoglobulin 10% with recombinant human hyaluronidase: replacement therapy in patients with primary immunodeficiency disorders. *BioDrugs*. 2014;28:411-420.
- Lejeune A, Martin L, Santibanez S, et al. Postexposure prophylaxis with intravenous immunoglobulin G prevents infants from getting measles. *Acta Paediatr*. 2017;106(1):174-177.
29. Berger M. Subcutaneous IgG in neurologic diseases. *Immunotherapy*. 2014;6:71-83.
30. Haddad E, Berger M, Wang EC, et al. Higher doses of subcutaneous IgG reduce resource utilization in patients with primary

- immunodeficiency. *J Clin Immunol*. 2012;32:281-289.
31. Hadden RD, Marreno F. Switch from intravenous to subcutaneous immunoglobulin in CIDP and MMN: improved tolerability and patient satisfaction. *Ther Adv Neurol Disord*. 2015;8:14-19.
 32. Hagan JB, Fasano MB, Spector S, et al. Efficacy and safety of a new 20% immunoglobulin preparation for subcutaneous administration, IgPro20, in patients with primary immunodeficiency. *J Clin Immunol*. 2010;30:734-745.
 33. Jolles S, Bernatowska E, de Gracia J, et al. Efficacy and safety of Hizentra® in patients with primary immunodeficiency after a dose-equivalent switch from intravenous or subcutaneous replacement therapy. *Clin Immunol*. 2011;141:90-102.