



CLINICAL GUIDELINES

Peripheral Vascular Disease (PVD) Guidelines

Version 2.0.2022

Effective March 1, 2022



eviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Imaging requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

eviCore's Clinical Review Criteria ("CRC") and related content is made available for the limited uses of: reference; and individual use, only limited to facilitating the determination of medically necessary and appropriate clinical treatment by clinicians for specific delegated patients under their care. The CRC and related content is proprietary information of eviCore, and copyrighted to the full extent of the law. Except as expressly permitted, you may not modify, copy, reproduce, republish, upload, post, transmit, hyperlink to or from, or distribute in any way the CRC, nor may you sell, transfer, distribute, assign, lease, reproduce, or otherwise use the CRC in commerce, in a manner that competes with us or infringes upon our rights, or for any public or commercial endeavor without our prior and express written consent.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2022 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Peripheral Vascular Disease (PVD) Imaging Guidelines	
Abbreviations and Glossary for the PVD Imaging Guidelines	3
PVD-1: General Guidelines	4
PVD-2: Screening for Suspected Peripheral Artery Disease/Aneurysmal Disease	10
PVD-3: Cerebrovascular and Carotid Disease	14
PVD-4: Upper Extremity Peripheral Vascular Disease	21
PVD-5: Pulmonary Artery Hypertension	23
PVD-6: Aortic Disorders, Renal Vascular Disorders and Visceral Artery Aneurysms	24
PVD-7: Lower Extremity Peripheral Vascular Disease	39
PVD-8: Imaging for Hemodialysis Access	46
PVD-9: Arteriovenous Malformations (AVMs)	48
PVD-10: Nuclear Medicine	50
PVD-11: Venous Imaging General Information	51
PVD-12: Acute Limb Swelling	56
PVD-13: Chronic limb swelling due to chronic deep venous thrombosis/May Thurner's syndrome	60
PVD-14: Chronic limb swelling due to venous insufficiency/Venous stasis changes/Varicose veins	63
PVD-15: Venous stasis ulceration	66
PVD-16: IVC filters	68
PVD-17: Post iliac vein stent/angioplasty	70

Abbreviations and Glossary for the PVD Imaging Guidelines

(See also: Cardiac Imaging Guidelines Glossary)

AAA	abdominal aortic aneurysm
ABI	Ankle brachial index: a noninvasive, non-imaging test for arterial insufficiency – (see toe-brachial index below). This testing can also be done after exercise if resting results are normal.
Claudication	or Intermittent claudication : usually a painful cramping sensation of the legs with walking or severe leg fatigue
CLI	Critical Limb Ischemia
CTA	computed tomography angiography
CTV	computed tomography venography
DLCO	diffusion capacity: defined as the volume of carbon monoxide transferred into the blood per minute per mmHg of carbon monoxide partial pressure
DVT	deep venous thrombosis
ECG	Electrocardiogram
ENT	Ears, Nose, Throat
HbA1C	hemoglobin A1C: test used to determine blood sugar control for individuals with diabetes
MRA	magnetic resonance angiography
MRV	magnetic resonance venography
PAD	peripheral artery disease
PAH	pulmonary artery hypertension
PFT	pulmonary function tests
PVD	peripheral vascular disease
SVC	superior vena cava
TIA	transient ischemic attack
TTE	transthoracic echocardiogram
Toe-Brachial Index	useful in individuals with ABI above the normal range due to non-compressible posterior tibial or dorsalis pedis arteries
V/Q Scan	ventilation and perfusion scan

PVD-1: General Guidelines

PVD-1.0: General Guidelines	5
PVD-1.1: General Information	6
PVD-1.2: Procedure Coding	7
PVD-1.3: General Guidelines – Imaging	9

PVD-1.0: General Guidelines

- A pertinent clinical evaluation, or meaningful technological contact (telehealth visit, telephone call, electronic mail or messaging), including medical treatments and a relevant history and physical, are required prior to considering advanced imaging, including:
 - ◆ Palpation of pulses
 - ◆ Evaluation of lower extremities for presence of non-healing wounds or gangrene
 - ◆ Associated skin changes such as thickened nails, absence of hair in the feet or calves, cool extremities
 - ◆ Evaluation for the presence of arterial bruits
 - ◆ Appropriate laboratory studies
 - ◆ Non-advanced imaging modalities, such as recent ABIs (within 60 days) after symptoms started or worsened
- ABI should be measured first:
 - ◆ If normal, then further vascular studies are generally not indicated.
 - ◆ If clinical suspicion for PAD remains high with normal ABI's, exercise ABI's (CPT® 93924) can be performed on a treadmill to elicit ischemia
 - ◆ The TBI (toe-brachial index) is used to establish the diagnosis of PAD in the setting of non-compressible arteries (ABI >1.40) and may also be used to assess perfusion in individuals with suspected CLI (rest pain and/or non-healing wound)
- If a prior imaging study (Ultrasound, MRA, CTA, Catheter angiogram, etc.) has been completed for a condition, a follow-up, additional, or repeat study for the same condition is generally not indicated unless there has been a change in the individual's condition, previous imaging showed an indeterminate finding, or eviCore healthcare guidelines support routine follow-up imaging.
- Runoff studies (CPT® 75635 for CTA or CPT® 74185, CPT® 73725, and CPT® 73725 for MRA) image from the umbilicus to the feet
 - ◆ CTA Abdomen and lower extremities should be reported as CPT® 75635, rather than using the individual CPT® codes for the abdomen, pelvis, and legs
 - ◆ MRA Abdomen, MRA Pelvis and MRA Lower extremities should be reported as CPT® 74185, CPT® 73725, and CPT® 73725. The CPT® code for MRA Pelvis (CPT® 72198) should not be included in this circumstance.

PVD-1.1: General Information

- Risk factors for vascular disease include:
 - ◆ Diabetes
 - ◆ Cigarette smoking
 - ◆ Hypertension
 - ◆ Hyperlipidemia
 - ◆ Age >50, with at least one risk factor, are considered “at risk” for vascular disease
 - ◆ See also: **PV-17: Impotence/Erectile Dysfunction** in the Pelvis Imaging Guidelines.
- Signs and symptoms of peripheral arterial disease
 - ◆ Claudication (Cramping pain in the legs, most notably back of the calves but can involve hips or thighs, after walking which is relieved with rest but recurs at a predictable distance)
 - Symptoms that are not consistent with claudication include
 - Generalized leg pain
 - Nocturnal cramps
 - Pain that is not easily relieved after a few minutes of rest
 - Burning pain in feet
 - ◆ Critical limb ischemia
 - Rest pain: Pain in the foot (not leg) at rest, particularly at night when the leg is elevated. Pain is relieved by dangling the leg off the bed or moving to an upright position
 - Non healing wounds: Wounds present for >2 weeks with little to no evidence of healing
 - ◆ Erectile dysfunction can be associated with vascular disease
- Claudication and critical limb ischemia have different natural histories. Claudication generally follows a benign indolent course. 70% of individuals with claudication will have the same symptoms after five years with no progression. Critical limb ischemia, on the other hand, is associated with a high rate of limb loss (25%) and death (35%) one year after presentation
- Simultaneous venous and arterial systems evaluation are unusual but are occasionally needed
- Post angioplasty/reconstruction: follow-up imaging is principally guided by symptoms. See also:
 - ◆ **PVD-6.8: Post Aortic Endovascular/Open Surgery Surveillance Studies**
 - ◆ **PVD-7.3: Post-Procedure Surveillance Studies**

PVD-1.2: Procedure Coding

Non-Invasive Physiologic Studies of Extremity Arteries	CPT®
➤ Limited bilateral noninvasive physiologic studies of upper or lower extremity arteries. ➤ Non-invasive physiologic studies of upper or lower extremity arteries, single level, bilateral (e.g., ankle/brachial indices, Doppler waveform analysis, volume plethysmography, transcutaneous oxygen tension measurement).	93922
➤ Complete bilateral noninvasive physiologic studies of upper or lower extremity arteries, 3 or more levels. ➤ Non-invasive physiologic studies of upper or lower extremity arteries, multiple levels or with provocative functional maneuvers, complete bilateral study (e.g., segmental blood pressure measurements, segmental Doppler waveform analysis, segmental volume plethysmography, segmental transcutaneous oxygen tension measurements, measurements with postural provocative tests, measurements with reactive hyperemia).	93923

- CPT® 93922 and CPT® 93923 can be requested and reported only once for the upper extremities and once for the lower extremities.
- CPT® 93922 and CPT® 93923 should not be ordered on the same request nor billed together for the same date of service.
- CPT® 93924 and CPT® 93922 and/or CPT® 93923 should not be ordered on the same request and should not be billed together for the same date of service.
- ABI studies performed with handheld dopplers, where there is no hard copy output for evaluation of bidirectional blood flow, are not reportable by these codes.

Non-Invasive Physiologic Studies of Extremity Arteries	CPT®
Non-invasive physiologic studies of lower extremity arteries, at rest and following treadmill stress testing, complete bilateral study.	93924

Arterial Duplex – Upper and Lower Extremities	CPT®
Duplex scan of lower extremity arteries or arterial bypass grafts; complete bilateral.	93925
➤ A complete duplex scan of the lower extremity arteries includes examination of the full length of the common femoral, superficial femoral and popliteal arteries. ➤ The iliac, deep femoral, and tibioperoneal arteries may also be examined.	
Duplex scan of lower extremity arteries or arterial bypass grafts; unilateral or limited study.	93926
➤ The limited study is reported when only one extremity is examined or when less than a full examination is performed (e.g. only one or two vessels or follow-up).	
Duplex scan of upper extremity arteries or arterial bypass grafts; complete bilateral.	93930
➤ A complete duplex of the upper extremity arteries includes examination of the subclavian, axillary, and brachial arteries. ➤ The radial and ulnar arteries may also be included.	
Duplex scan of upper extremity arteries or arterial bypass grafts; unilateral or limited study.	93931
➤ The limited study is reported when only one extremity is examined or when less than a full examination is performed (e.g. only one or two vessels or follow-up).	

Cerebrovascular Artery Studies	CPT®
Duplex scan of extracranial arteries; complete bilateral study.	93880
Duplex scan of extracranial arteries; unilateral or limited study.	93882
➤ This study is often referred to as a “carotid ultrasound” or “carotid duplex”. ➤ Typically, it includes evaluation of the common, internal, and external carotid arteries.	

Transcranial Doppler Studies		CPT®
Transcranial Doppler study of the intracranial arteries; complete study		93886
Transcranial Doppler study of the intracranial arteries; limited study		93888
Transcranial Doppler vasoreactivity study		93890
Transcranial Doppler study of the intracranial arteries; emboli detection without intravenous microbubble injection		93892
Transcranial Doppler study of the intracranial arteries; emboli detection with intravenous microbubble injection		93893
Venous Studies - Extremities		CPT®
Duplex scan of extremity veins, including responses to compression and other maneuvers; complete bilateral study.		93970
Duplex scan of extremity veins, including responses to compression and other maneuvers; unilateral or limited study.		93971
➤ These codes are used to report studies of lower or upper extremity veins. ➤ A complete bilateral study of the lower extremity veins includes examination of the common femoral, proximal deep femoral, great saphenous and popliteal veins. Calf veins may also be included. ➤ A complete bilateral study of upper extremity veins includes examination of the subclavian, jugular, axillary, brachial, basilica, and cephalic veins. Forearm veins may also be included.		
Visceral Vascular Studies		CPT®
Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; complete study.		93975
Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; limited study		93976
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study		93978
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; unilateral or limited study		93979
Duplex for Hemodialysis Access		CPT®
Duplex scan of hemodialysis access (including arterial inflow, body of access and venous outflow).		93990
Duplex scan of arterial inflow and venous outflow for preoperative vessel assessment prior to creation of hemodialysis access; complete bilateral study		93985
Duplex scan of arterial inflow and venous outflow for preoperative vessel assessment prior to creation of hemodialysis access; complete unilateral study		93986

PVD-1.3: General Guidelines – Imaging

➤ **Imaging Studies:**

- ◆ Carotid studies (MRA Neck or CTA Neck) capture the area from the top of the aortic arch (includes the origin of the innominate artery, common carotid artery, and subclavian artery, which gives off the vertebral artery) to the base of the skull.
- ◆ CTA or MRA Abdomen (CPT® 74175 or CPT® 74185) images from the diaphragm to the umbilicus or iliac crest
- ◆ CTA or MRA Chest (CPT® 71275 or CPT® 71555) images from the base of the neck to the dome of the liver
- ◆ Runoff studies (CPT® 75635 for CTA or CPT® 74185, CPT® 73725, and CPT® 73725 for MRA) image from the umbilicus to the feet
 - CTA Abdomen and lower extremities should be reported as CPT® 75635, rather than using the individual CPT® codes for the abdomen, pelvis, and legs
 - MRA Abdomen, MRA Pelvis and MRA Lower extremities should be reported as CPT® 74185, CPT® 73725, and CPT® 73725. The CPT® code for MRA Pelvis (CPT® 72198) should not be included in this circumstance
- ◆ Studies used to quantify plaque morphology in noncoronary vessels (CPT® 0710T, CPT® 0711T, CPT® 0712T, CPT® 0713T) are considered experimental, investigational, or unproven.

References

1. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the management of patients with lower extremity peripheral artery disease. *J Am Coll Cardiol*. 2017 Mar 69 (11):1467-1508.
2. Perlstein TS and Creager MA. The ankle-brachial index as a biomarker of cardiovascular risk: it's not just about the legs. *Circulation*. 2009 Nov 29; 120 (21):2033-2035.

PVD-2: Screening for Suspected Peripheral Artery Disease/Aneurysmal Disease

PVD-2.1: Asymptomatic Screening	11
PVD-2.2: Screening for Vascular related genetic connective tissue Disorders	11
PVD-2.3: Screening for TAA with bicuspid aortic valves	12

PVD-2.1: Asymptomatic Screening

- Routine screening of asymptomatic individuals for PAD is not advised. Those with CVD risk factors should be placed on best medical management and should be questioned on symptoms of PAD at annual physicals
- Resting ABI's may be indicated in individuals with abnormal pulse exams
- Currently, there is no evidence to demonstrate that screening all individuals with PAD for asymptomatic atherosclerosis in other arterial beds improves clinical outcome

PVD-2.2: Screening for Vascular related genetic connective tissue Disorders

- Vascular related genetic connective tissue Disorders include:
 - ◆ Familial Aneurysm Syndromes
 - ◆ Fibromuscular Dysplasia
 - ◆ Spontaneous Coronary Artery Dissection (SCAD)
 - ◆ Ehlers-Danlos
 - ◆ Marfan
 - ◆ Loeys-Dietz
- Screening for Familial Syndromes in individuals with a positive family history (1st degree relative with dissection/TAA) but no known genetic syndrome/mutation, otherwise known as **Suspected Familial Aneurysm Syndrome**.
 - ◆ ECHO (CPT® 93306, CPT® 93307, or CPT® 93308) and chest x-ray for all First-degree relatives (parents, siblings, children) of individuals with TAA and/or dissection.
 - ◆ Any imaging listed can be performed if these studies identify a TAA or are equivocal or do not visualize the ascending aorta adequately.
 - ◆ Studies can be repeated at 2 year intervals if negative
- Initial imaging for individuals with documented SCAD/fibromuscular dysplasia/Marfan/Loeys-Dietz/Ehlers-Danlos type IV:
 - ◆ On initial diagnosis full vascular imaging should be performed from head to pelvis with:
 - CTA/MRA Head
 - Carotid Duplex
 - CTA Chest or CT Chest with contrast
 - CTA Abdomen/Pelvis or MRA Abdomen/Pelvis
 - ◆ If there are no identified aneurysms or dissections, repeat imaging can be obtained at 2 year intervals

- Surveillance imaging
 - ◆ If an aneurysm is identified in individuals with fibromuscular dysplasia, then the aneurysm can be surveilled per the typical timeframe as described in **PVD-6.2: Thoracic Aortic Aneurysm**, **PVD-6.3: Abdominal Aortic Aneurysm** and **PVD-6.4: Iliac Artery Aneurysm** and **PVD-6.5: Visceral Artery Aneurysm**.
 - ◆ Follow-Up of aneurysms in individuals with documented SCAD/Marfan's/Loeys-Dietz/Ehlers-Danlos type IV.
 - Imaging can be performed every 6 months once an aneurysm has been identified until a decision has been made to repair.
 - Intracranial aneurysm – CTA or MRA Head (CPT® 70496 or 70544)
 - Aneurysm of a cervical artery – Carotid duplex or CTA or MRA neck if unable to fully visualize with carotid duplex
 - Thoracic aorta – CTA Chest (CPT® 71275) or CT Chest with (CPT® 71260) or without (CPT® 71250)
 - Abdominal aneurysm – Abdominal duplex (CPT® 93975/93976/93978/93979/76770/76775)
 - Visceral aneurysm – These can be difficult to visualize on duplex. If not visible on duplex, can obtain a CTA or MRA Abdomen.

PVD-2.3: Screening for TAA with bicuspid aortic valves

- Screening in individuals with bicuspid aortic valve:
 - ◆ Screening, any requested imaging from the “**Table of Thoracic Aorta Imaging Options**” in **PVD-6.2: Thoracic Aortic Aneurysm** and/or ECHO (CPT® 93306, CPT® 93307, or CPT® 93308).
 - Additional imaging such as Cardiac MRI, Cardiac CT, or CCTA is NOT generally indicated.
 - There is no evidence-based data to support screening relatives of individuals with bicuspid aortic valve for TAA except with echocardiogram.
 - ◆ Follow-up per TAA Follow-Up guidelines in **PVD-6.2: Thoracic Aortic Aneurysm (TAA)**
- If no dilatation of the aortic root or ascending thoracic aorta is found, there is no evidence-based data to support continued surveillance imaging.

References

1. Demo E, Rigelsky C, Rideout AL, et al. Genetics and Precision Medicine: Heritable Thoracic Aortic Disease. *Med Clin North Am*. 2019;103(6):1005-1019. doi:10.1016/j.mcna.2019.08.001.
2. Firnhaber JM et al. Lower Extremity Peripheral Arterial Disease: Diagnosis and Treatment. *Am Fam Physician*. 2019;362-369.
3. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients with Lower Extremity Peripheral Artery Disease: Executive Summary. *J Am Coll Cardiol*. 2017;69(11):1465-1508. doi:10.1016/j.jacc.2016.11.008.
4. Gornik HL, Persu A, Adlam D, et al. First International Consensus on the diagnosis and management of fibromuscular dysplasia [published correction appears in Vasc Med. 2019 Oct;24(5):475. *Vasc Med*. 2019;24(2):164-189. doi:10.1177/1358863X18821816.
5. Hayes SN, Kim ES, Saw J, et al. Spontaneous Coronary Artery Dissection: Current State of the Science: A Scientific Statement from the American Heart Association. *Circulation*. 2018;137(19). doi:10.1161/cir.0000000000000564.
6. Hiratzka LF, Bakris GL, Beckman JA, et al. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with thoracic aortic disease. *J Am Coll Cardiol*. 2010; 55: e27-e129.
7. MacCarrick G, Black J, Bowdin S, et al. Loeys–Dietz syndrome: a primer for diagnosis and management. *Genet Med* (2014).16:576–587 doi:10.1038/gim.2014.11.
8. Mohler ER, Gornik HL, Gerhard-Herman M, Misra S, Olin JW, Zierler RE. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 Appropriate Use Criteria for Peripheral Vascular Ultrasound and Physiological Testing Part I: Arterial Ultrasound and Physiological Testing. *J Am Coll of Cardiol*. 2012;60(3):242-276. doi:10.1016/j.jacc.2012.02.009.
9. MS Conte, FB Pomposelli, DG Clair, et al. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: Management of asymptomatic disease and claudication. *J Vasc Surg*. 2015. Vol 6:1S-41S.
10. Olin JW, Gornik HL, Bacharach JM, et al. Fibromuscular Dysplasia: State of the Science and Critical Unanswered Questions. *Circulation*. 2014;129(9):1048-1078. doi:10.1161/01.cir.0000442577.96802.8c.
11. Persu A, Niepen PVD, Touzé E, et al. Revisiting Fibromuscular Dysplasia. *Hypertension*. 2016;68(4):832-839. doi:10.1161/hypertensionaha.116.07543
12. Saydah SH. Poor Control of Risk Factors for Vascular Disease among Adults with Previously Diagnosed Diabetes. *JAMA*. 2004;291(3):335-342. doi:10.1001/jama.291.3.335.
13. Schoenhoff F, Schmidli J, Czerny M, Carrel TP. Management of aortic aneurysms in patients with connective tissue disease. *J Cardiovasc Surg (Torino)*. 2013;54(1 Suppl 1):125-134.
14. US Preventive Services Task Force, Curry SJ, Krist AH, et al. Screening for Peripheral Artery Disease and Cardiovascular Disease Risk Assessment with the Ankle-Brachial Index:US Preventive Services Task Force Recommendation Statement. *JAMA*. 2018;320(2):177. doi:10.1001/jama.2018.8357.

PVD-3: Cerebrovascular and Carotid Disease	
PVD-3.1: Initial Imaging	15
PVD-3.2: Surveillance Imaging with NO History of Carotid Surgery or Intervention	17
PVD-3.3: Surveillance Imaging WITH History of Carotid Surgery or Intervention	18

PVD-3.1: Initial Imaging

- Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) should generally be used to evaluate possible carotid artery disease, prior to considering advanced imaging, when ANY of the following apply:
 - ◆ Hemispheric neurologic symptoms including stroke, TIA, or amaurosis fugax.
 - ◆ Known or suspected retinal arterial emboli or Hollenhorst plaque
 - ◆ Suspected carotid dissection
 - ◆ Pulsatile neck masses
 - ◆ Carotid or cervical bruit
 - ◆ Abnormal findings on physical exam of the carotid arteries (e.g. aneurysm or absent carotid pulses)
 - ◆ Preoperative evaluation of individuals with evidence of severe diffuse atherosclerosis, scheduled for major cardiovascular surgical procedures
 - ◆ Preoperative evaluation of individuals prior to elective cardiovascular surgery in individuals older than 65 years of age and in those with peripheral artery disease, history of cigarette smoking, history of stroke or TIA, or carotid bruit
 - ◆ Suspected Subclavian Steal Syndrome
 - See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging Guidelines
 - ◆ Blunt neck trauma
 - ◆ Neurologic complaints after chiropractic neck manipulation
 - ◆ Vasculitis potentially involving carotid arteries, i.e., Takayasu's arteritis and fibromuscular dysplasia (FMD)
- Carotid ultrasound screening in asymptomatic individuals due only to risk factors is not indicated
- New signs and symptoms consistent with carotid artery disease (e.g. TIA, amaurosis fugax, change in nature of a carotid bruit) are an indication to re-image the cervical vessels (regardless of when the previous carotid imaging was performed) using one of the following:
 - ◆ Duplex ultrasound (CPT® 93880 bilateral study or CPT® 93882 unilateral study),
 - ◆ MRA Neck with contrast (CPT® 70548) or without and with contrast (CPT® 70549)
 - ◆ CTA Neck (CPT® 70498)
- For Typical Symptoms of TIA/Stroke or Carotid Dissection:
 - ◆ See **HD-21: Stroke/TIA** in the Head Imaging Guidelines

- CTA or MRA Neck can be approved for suspected vertebrobasilar pathology:
 - ◆ Symptoms include:
 - Vertigo associated with nausea and vomiting
 - Diplopia
 - Loss of vision in one or both eyes
 - Dysarthria
 - Bifacial numbness
 - Bilateral extremity weakness and/or numbness
 - Acute changes in mental status
 - Loss of consciousness
 - Ataxia
 - ◆ MRA or CTA of **both** Neck and Head are required to visualize the entire vertebral-basilar system for evaluation of posterior circulation disease. See **HD-1.5: General Guidelines – CT and MR Angiography in the Head Imaging Guidelines**
 - ◆ For suspected carotid artery dissection, a CTA or MRA neck can be approved in individuals with ANY of the following mechanisms of injury, or risk factors, for arterial dissection:
 - Chiropractic manipulation of neck
 - Whiplash injury
 - Fibromuscular dysplasia/Marfan's
 - Stroke in the young (age ≤50)
 - See **HD-21: Stroke/TIA** in the Head Imaging Guidelines
 - ◆ Surveillance imaging, post-stenting or known vertebrobasilar disease, interval determined by Vascular Specialist, Neurologist, or Neurosurgeon or any provider in consultation with a vascular specialist, neurologist, or neurosurgeon for ANY of the following:
 - Asymptomatic
 - Unchanged symptoms
 - New or worsening symptoms
- **After Intracranial Hemorrhage:**
 - ◆ Initial Imaging see **HD-13.1: Head Trauma** in the Head Imaging Guidelines
 - ◆ Surveillance Imaging
 - Interval determined by neurosurgeon or neurologist or any provider in consultation with a neurologist or neurosurgeon.
- For Suspected Subclavian Steal Syndrome:
 - ◆ Initial imaging should be a carotid duplex
 - If initial duplex demonstrates high grade stenosis or occlusion of the subclavian artery, advanced imaging is NOT indicated unless the individual is symptomatic with arm claudication or signs of hypo-perfusion of the vertebral artery with recurrent dizziness
 - ◆ Surveillance of subclavian arterial disease is NOT indicated if there has not been any intervention such as a carotid-subclavian bypass or subclavian stent
Advanced imaging, see **CH-27: Subclavian Steal Syndrome – General** in the Chest Imaging Guidelines

PVD-3.2: Surveillance Imaging with NO History of Carotid Surgery or Intervention

- Surveillance imaging is indicated once a year for individuals with fibromuscular dysplasia of the extracranial carotid arteries.
- Reporting standards for carotid stenosis varies widely. The most commonly used criteria, however, is noted in the chart below published by the Society of Radiology in 2003

% Stenosis	Primary parameters		Additional Parameters	
	ICA PSV (cm/sec)	Plaque estimate (%)	ICA/CCA PSV ratio	ICA/EDV (cm/sec)
Normal	< 125	None	< 2.0	< 40
< 50	< 125	< 50	< 2.0	< 40
50-69	125-230	≥ 50	2.0-4.0	40-100
≥ 70 but less than near occlusion	> 230	> 50	> 4.0	> 100
Near occlusion	High, low, or undetectable	Visible	Variable	Variable
Total occlusion	undetectable	Visible- no detectable lesion	Not applicable	Not applicable

- If normal study, no routine follow-up imaging is indicated
- If <50% carotid stenosis
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) can be performed every **two** years
- Between 50% and 70% carotid stenosis
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) can be performed annually.
 - ◆ A repeat duplex (CPT® 93880 bilateral or CPT® 93882 unilateral) may be performed in three to six months until stability is reached when **one** of the following occurs:
 - Change in the character of the bruit
 - Duplex demonstrates rapid progression, including:
 - Doubling of peak systolic velocities
 - Increase of the ICA/CCA ratio
 - Heavy calcification
 - Thrombus
 - Ulcerated plaque
 - Echolucent plaque
 - ◆ A one-time CTA Neck (CPT® 70498) is indicated to confirm degree of stenosis in individuals with ulcerated plaque or heavy calcification seen on duplex.

- Carotid stenosis $\geq 70\%$ or ICA/CCA ratio >4
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) or MRA Neck with contrast (CPT® 70548) or CTA Neck (CPT® 70498) can be performed at the following intervals:
 - Every 6 months until one of the following occurs:
 - Intervention is performed
 - Decision is made to not intervene
- MRA Neck with contrast (CPT® 70548) or CTA Neck (CPT® 70498) is indicated if duplex Ultrasound shows $\geq 70\%$ occlusion/stenosis of the internal carotid artery or the ICA/CCA ratio is >4.0 , even with a lower percentage of stenosis.
 - ◆ If carotid stent is planned
 - MRA Head (CPT® 70544, or CPT® 70545, or CPT® 70546) or CTA Head (CPT® 70496) can be added

PVD-3.3: Surveillance Imaging WITH History of Carotid Surgery or Intervention

- Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) is indicated post carotid surgery or intervention at the following intervals:
 - ◆ 1 month after procedure
 - ◆ Every 6 months for 2 years after procedure
 - ◆ Then annually
- If $\geq 70\%$ residual carotid stenosis is seen on duplex at 1 month after procedure
 - ◆ Duplex ultrasound (CPT® 93880 bilateral or CPT® 93882 unilateral) or CTA Neck (CPT® 70498) is indicated at the following intervals:
 - Every 3-6 months for one year
 - Then annually or until decision is made to re-intervene.
- If $\geq 70\%$ residual carotid stenosis is seen on duplex at any time post procedure, then
 - ◆ CTA Neck (CPT® 70498) is indicated for further evaluation and at six month intervals until decision is made to re-intervene.

Practice Notes

- Carotid intima-media thickness using duplex ultrasound imaging (Category III code 0126T) is not recommended in clinical practice for risk assessment for a first ASCVD event. Although outcomes data are lacking, Texas has adopted this method in Texas Heart Attack Preventive Screening Bill (HR 1290)
- Texas Heart Attack Preventive Screening Law (HR 1290) mandates that insurers in Texas cover either a calcium scoring study (CPT® 75571 or HCPCS S8092) or a carotid intima-media thickness study (ultrasound—Category III code 0126T) every five years for certain populations. To qualify, the following must apply:
 - ◆ Must be a member of a fully-insured Texas health plan.
 - ◆ Must be male sex assigned at birth age 45 to 75 or female sex assigned at birth age 55 to 75.
 - ◆ Must have either diabetes or a Framingham cardiac risk score of intermediate or higher.
 - ◆ Must not have had a calcium scoring study or a carotid intima-media thickness study within the past 5 years
- MRA Neck (CPT® 70548) or CTA Neck (CPT® 70498) may be indicated if ultrasound is technically difficult or confirmation of the degree of stenosis on ultrasound is needed because an interventional procedure is being considered.

References

1. Aboyans V, Ricco J-B, Bartelink M-LEL, et al. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS). *European Heart Journal*. 2017;39(9):763-816. doi:10.1093/eurheartj/ehx095.
2. Aburahma AF, Srivastava M, Aburahma Z, et al. The value and economic analysis of routine postoperative carotid duplex ultrasound surveillance after carotid endarterectomy. *J Vasc Surg*. 2015;62(2):378-384. doi:10.1016/j.jvs.2015.03.023.
3. ACR–AIUM–SPR–SRU. AIUM Practice Parameter for the Performance of an Ultrasound Examination of the Extracranial Cerebrovascular System. *Journal of Ultrasound in Medicine*. 2016;35(9):1-11. doi:10.7863/ultra.35.9.1.
4. Brott TG, Halperin JL, Abbara S, et al. 2011 ASA/ACCF/AHA/AANN/AANS/ACR/ASNR/CNS/SAIP/SCAI/SIR/SNIS/SVM/SVS Guideline on the Management of Patients with Extracranial Carotid and Vertebral Artery Disease. *J Am Coll Cardiol*. 2011;57(8). doi:10.1016/j.jacc.2010.11.006.
5. Derdeyn CP, Zipfel GJ, Albuquerque FC, et al. Management of Brain Arteriovenous Malformations: A Scientific Statement for Healthcare Professionals from the American Heart Association/American Stroke Association. *Stroke*. 2017;48(8). doi:10.1161/str.0000000000000134.
6. Gerhard-Herman M, Gardin JM, Jaff M, Mohler E, Roman M, Naqvi TZ. Guidelines for Noninvasive Vascular Laboratory Testing: A Report from the American Society of Echocardiography and the Society of Vascular Medicine and Biology. *J Am Soc Echocardiogr*. 2006;19(8):955-972. doi:10.1016/j.echo.2006.04.019.
7. Goff DC, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk. *Circulation*. 2014;129(25 suppl 2):s49-s73. doi:10.1161/01.cir.0000437741.48606.98.
8. Mohler ER, Gornik HL, Gerhard-Herman M, Misra S, Olin JW, Zierler RE. ACCF/ACR/AIUM/ASE/ASN/ICAVL/SCAI/SCCT/SIR/SVM/SVS 2012 Appropriate Use Criteria for Peripheral Vascular Ultrasound and Physiological Testing Part I: Arterial Ultrasound and Physiological Testing. *J Am Coll Cardiol*. 2012;60(3):242-276. doi:10.1016/j.jacc.2012.02.009.
9. Moresoli P, Habib B, Reynier P, et al. Carotid Stenting versus Endarterectomy for Asymptomatic Carotid Artery Stenosis. *Stroke*. 2017;48(8):2150-2157. doi:10.1161/strokeaha.117.016824.
10. Naylor A, Ricco J-B, Borst GD, et al. Editor's Choice – Management of Atherosclerotic Carotid and Vertebral Artery Disease: 2017 Clinical Practice Guidelines of the European Society for Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg*. 2018;55(1):3-81. doi:10.1016/j.ejvs.2017.06.021.
11. Ricotta JJ, Aburahma A, Ascher E, et al. Updated Society for Vascular Surgery guidelines for management of extracranial carotid disease. *J Vasc Surg*. 2011;54(3):1-31. doi:10.1016/j.jvs.2011.07.031.
12. Reutern G-MV, Goertler M-W, Bornstein NM, et al. Grading Carotid Stenosis Using Ultrasonic Methods. *Stroke*. 2012;43(3):916-921. doi:10.1161/strokeaha.111.636084.
13. Rogers RK, Bishu K. Optimal Treatment of Extracranial Carotid Artery Disease: Carotid Endarterectomy, Carotid Stenting, or Optimal Medical Therapy. *Current Cardiology Reports*. 2015;17(10). doi: 10.1007/s11886-015-0636-2.
14. Salmela MB, Mortazavi S, Jagadeesan BD, et al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria® Cerebrovascular Disease. *J Am Coll Radiol*. 2017;14(5S):S34-S61. doi:10.1016/j.jacr.2017.01.051.
15. Shakarchi JA, Lowry D, Nath J, et al. Duplex ultrasound surveillance after carotid artery endarterectomy. *Journal of Vascular Surgery*. 2016;63(6):1647-1650. doi:10.1016/j.jvs.2016.01.054.

PVD-4: Upper Extremity Peripheral Vascular Disease

PVD-4.1: Upper Extremity PVD – Imaging

- Signs and symptoms of arterial insufficiency include but are not limited to:
 - ◆ Arm or hand claudication, cramping or fatigue of the unilateral extremity with use or with raising limb overhead that is relieved with rest and is reproducible. See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging Guidelines
 - ◆ Systolic blood pressure differential between arms of <15mmHg. See **CH-27: Subclavian Steal Syndrome** in the Chest Imaging Guidelines
 - ◆ Bluish discoloration of the hand or fingers
 - ◆ Unilateral cold painful pulseless hand
 - ◆ Non healing wound (>2 weeks with no healing or evidence of healing) or frank gangrene
- For signs and symptoms of arterial insufficiency, appropriate studies include:
 - ◆ Arterial ultrasound upper extremities (CPT® 93930 or CPT® 93931) or
 - ◆ CTA Upper extremity (CPT® 73206) or MRA of Upper extremity (CPT® 73225) and/or
 - ◆ CTA Chest (CPT® 71275) or MRA Chest (CPT® 71555)
- For suspected Fibromuscular Dysplasia of the brachial artery, appropriate studies include:
 - ◆ MRA Upper extremity (CPT® 73225)
 - ◆ CTA Upper extremity (CPT® 73206)
 - ◆ Arterial Ultrasound (CPT® 93930 bilateral study or CPT® 93931 unilateral study)
- Arterial Duplex (CPT® 93931) can be obtained following upper extremity arterial revascularization at:
 - ◆ Baseline (within one month)
 - ◆ 6 months
 - ◆ Then annually if stable
 - ◆ Anytime for new or worsening symptoms
- For symptoms of venous insufficiency including but not limited to unilateral pain and swelling of the upper extremity
 - ◆ Venous duplex upper extremities (CPT® 93970 or CPT® 93971) should be performed initially
 - ◆ If duplex ultrasound is non-diagnostic:
 - MRV Upper extremity (CPT® 73225) and/or MRV Chest (CPT® 71555), or
 - CTV Upper extremity (CPT® 73206) and/or CTV Chest (CPT® 71275)
 - CTV Upper extremity (CPT® 73206) or MRV Upper extremity (CPT® 73225), and/or CTV Chest (CPT® 71275) or MRV Chest (CPT® 71555) can be performed if there is a history of exertion with the limb such as with weight lifting or in the presence of central venous access (port, PICC line, to name a few) with a negative venous duplex. See **CH-31.1: Thoracic Outlet Syndrome** in the Chest Imaging Guidelines.

- For Superior Vena Cava Syndrome (upper extremity and facial symptoms) **one** of the following can be approved when stenting of the SVC is being considered:
 - ◆ CT Chest with contrast (CPT® 71260)
 - ◆ MRV Chest (CPT® 71555)
 - ◆ CTV Chest (CPT® 71275)

References

1. Desjardins B, Rybicki FJ, Kim HS, et al. ACR Appropriateness Criteria® Suspected Upper Extremity Deep Vein Thrombosis. *J Am Coll Radiol*. 2012;9(9):613-619. doi:10.1016/j.jacr.2012.05.021.
2. Kraaijpoel N, Es NV, Porreca E, Büller HR, Nisio MD. The diagnostic management of upper extremity deep vein thrombosis: A review of the literature. *Thrombosis Research*. 2017;156:54-59. doi:10.1016/j.thromres.2017.05.035.
3. Skeik N, Soo-Hoo SS, Porten BR, et al. Arterial Embolisms and Thrombosis in Upper Extremity Ischemia. *Vascular and Endovascular Surgery*. 2015;49(5-6):100-109. doi:10.1177/1538574415596740.
4. Hughes K, Cubangbang M, Blackman K, et al. Upper Extremity Bypass for Chronic Ischemia—A National Surgical Quality Improvement Program Study Database Study. *Vascular and Endovascular Surgery*. 2013;47(3):192-194. doi:10.1177/1538574413478472.

PVD-5: Pulmonary Arterial Hypertension

PVD-5.1: Pulmonary Arterial Hypertension – Imaging

- Pulmonary arterial hypertension (PAH) is a complex, chronic diseases which requires extensive evaluation, including ECG (right ventricular hypertrophy with/without strain, right atrial dilatation); chest x-ray; arterial blood gas, PFT's, CT angiography or V/Q scan. Imaging is based on suspected etiology.
- Transthoracic echocardiogram (TTE) (CPT® 93306) should be performed initially as it can help determine the probability of pulmonary hypertension. Echo may be accompanied by:
 - ◆ Stress echocardiogram (CPT® 93350 or CPT® 93351) , especially in the setting of concomitant valvular disease to assess for treatment
 - ◆ Left and/or right heart catheterization for direct measurement of pressures
 - ◆ High-resolution CT Chest (CPT® 71250) to rule out restrictive lung disorders such as idiopathic pulmonary fibrosis in the setting of hypoxemia
- For suspected acute and/or chronic pulmonary embolism one of the following:
 - ◆ CTA Chest (CPT® 71275)
 - ◆ MRA Chest (CPT® 71555)
 - ◆ V/Q scan (CPT® 78580-Pulmonary Perfusion Imaging CPT® 78582-Pulmonary Ventilation (e.g., Aerosol or Gas) and Perfusion Imaging). If requested can add to V/Q scan one of the following:
 - SPECT imaging (CPT® 78803)
 - SPECT/CT imaging (CPT® 78830)
- See in specific subsections:
 - ◆ CD-2.2: Transthoracic Echocardiogram (TTE)-Indications, CD-7.4: Right Heart Catheterization (RHC), CD-11.3.12: Severe Pulmonary artery hypertension (PHT) and Eisenmenger syndrome in the Cardiac Imaging Guidelines
 - ◆ PEDCD-2.3: Congenital Heart Disease Modality Considerations, PEDCD-7.1: Pediatric Pulmonary Hypertension - General in the Pediatric Cardiac Imaging Guidelines
 - ◆ **CH-25: Pulmonary Embolism (PE)** in the Chest Imaging Guidelines.

References

1. Barbosa EJM, Gupta NK, Torigian DA, et al. Current role of imaging in the diagnosis and management of pulmonary hypertension. *Am J Roentgenol*. 2012; 198 (6): 1320-1331.
2. Galiè A, Torbicki A, Barst R, et al. Guidelines on diagnosis and treatment of pulmonary arterial hypertension: the task force on diagnosis and treatment of pulmonary arterial hypertension of the European Society of Cardiology. *Eur Heart J*. 2004 Dec 1; 25 (24): 2243-2278.
3. Peña E, Dennie C, Veinot J, et al. Pulmonary hypertension: how the radiologist can help. *RadioGraphics*. 2012; 32 (1):9-32
4. Mazurek A, Dziuk M, Witkowska-Patena E, Piszczek S, Gizewska A. The Utility of Hybrid SPECT/CT Lung Perfusion Scintigraphy in Pulmonary Embolism Diagnosis. *Respiration*. 2015;90(5):393-401. doi:10.1159/000439543.
5. Yaghi S, Novikov A, Trandafirescu T. Clinical update on pulmonary hypertension. *J Investig Med*. 2020;68(4):821-827. doi:10.1136/jim-2020-001291.

PVD-6: Aortic Disorders, Renal Vascular Disorders and Visceral Artery Aneurysms	
PVD-6.1: Aortic Disorders General Information	25
PVD-6.2: Thoracic Aortic Aneurysm (TAA)	25
PVD-6.3: Abdominal Aortic Aneurysm (AAA)	27
PVD-6.4: Iliac Artery Aneurysm (IAA)	28
PVD-6.5: Visceral Artery Aneurysm	28
PVD-6.6: Renovascular Hypertension/Renal Artery Stenosis	29
PVD-6.7: Aortic Dissection and Other Aortic Conditions	30
PVD-6.8: Post Aortic Endovascular/Open Surgery Surveillance Studies	32
PVD-6.8.1: Post-operative surveillance after TEVAR for any indication	32
PVD-6.8.2: Post-operative surveillance after abdominal EVAR (endovascular aneurysm repair)	33
PVD-6.8.3: Endovascular (Stent) Iliac Repair	33
PVD-6.9: Large Vessel Vasculitis	34
PVD-6.9.1: Inflammatory Aortitis	34
PVD-6.9.2: Giant Cell Arteritis	35
PVD-6.9.3: Takayasu Arteritis	36

PVD-6.1: Aortic Disorders General Information

Duplex ultrasound for visceral vascular studies	CPT®
Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; complete study.	93975
Duplex scan of arterial inflow and venous outflow of abdominal, pelvic, scrotal contents and/or retroperitoneal organs; limited study.	93976
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study.	93978
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; unilateral or limited study.	93979
Ultrasound, abdominal aorta, real time, with image documentation, screening study for abdominal aortic aneurysm (AAA) for AAA screening	76706

- In clinical practice, CT, CTA, MRA are usually preferred to evaluate for stenosis of these vessels rather than ultrasound (Exception: Duplex ultrasound is indicated to rule out testicular or ovarian torsion or to evaluate an abdominal bruit or a pulsatile abdominal mass).
- Mesenteric Ischemia
 - ◆ See **AB-6: Mesenteric/Colonic Ischemia** in the Abdomen Imaging Guidelines.

PVD-6.2: Thoracic Aortic Aneurysm (TAA)

The thoracic aorta is generally divided into two segments: the ascending aorta which includes the aortic root, aortic arch and ends just distal to the left subclavian artery and the descending aorta which starts just distal to the left subclavian artery to the level of the diaphragm.

- Advanced imaging with a CT or MR is preferred imaging for this diagnosis. Transesophageal echocardiogram (TEE) can also be indicated for initial imaging of ascending and descending thoracic aortic aneurysms. For repeat imaging or established thoracic aneurysms, TEE is indicated **only** when imaging with CT or MR is contraindicated.
- Given the diversity of studies, pathology, and provider preference, **ONE** of the imaging studies in the Table of Thoracic Aorta Imaging Options can be approved for Thoracic Aortic Aneurysm (TAA) as indicated in this section

Table of Thoracic Aorta Imaging Options	CPT®
CT Chest with contrast	71260
CT Chest without contrast	71250
CTA Chest	71275
MRA Chest	71555
TEE	93312- 93314

- For TAA associated with a dissection, please see section **PVD-6.7 Aortic Dissection and Other Aortic Conditions**
- For suspected TAA, any requested imaging from the **Table of Thoracic Aorta Imaging Options** above:
 - ◆ Abnormalities identified on chest x-ray (abnormality including widened mediastinum, suspicious calcifications) or other imaging studies (fluoroscopy, MRI Spine, etc.) abnormality.
- For known TAA accompanied with chest pain or back pain and suspicion of rupture, any requested imaging from the **Table of Thoracic Aorta Imaging Options** above.
- For planning for pre–thoracic endovascular repair (TEVAR) of thoracic aorta disease.
 - ◆ CTA Chest, and/or Abdomen, and/or Pelvis (CPT® 71275, CPT® 74175, CPT® 72191, CPT® 74174); or
 - ◆ MRA Chest, and/or Abdomen, and/or Pelvis (CPT® 71555, CPT® 74185, CPT® 72198).
- For follow-up of ascending aortic aneurysms CTA Chest (CPT® 71275) or CT Chest (CPT® 71250 or CPT® 71260) or MRA chest (CPT® 71555)
 - ◆ Operative treatment is reasonable for asymptomatic individuals when the diameter of the arch exceeds 5.5 cm.
 - ◆ For individuals with ascending aortic aneurysms <4.0 cm in diameter
 - Repeat imaging annually
 - ◆ For individuals with ascending aortic aneurysms ≥4.0 cm
 - Repeat imaging 6 months.
 - ◆ TEE is indicated **only** when imaging with CT or MR is contraindicated
- For follow-up of descending aortic aneurysms, any requested imaging from the “Table of Thoracic Aorta Imaging Options” above for the following:
 - ◆ “Medically” treated/observation.
 - 3.5 to 4.4 cm TAA can be followed annually.
 - ≥4.5 cm TAA can be followed every 6 months.
 - ≥3.0 cm TAA when there is concern for growth can have a one-time 3-month interval advanced imaging.
 - ◆ TEE is indicated **only** when imaging with CT or MR is contraindicated
- Screening in the presence of other aortic aneurysms.
 - ◆ In an individual with a known TAA, screening for AAA is indicated with an abdominal duplex. See **PVD-6.3: Abdominal Aortic Aneurysm (AAA)**.
 - ◆ In an individual with a known AAA, screening for TAA is not supported by sufficient evidence.
- Screening in individuals with bicuspid aortic valve or familial TAA syndromes. See **PVD-2.3: Screening for TAA with bicuspid aortic valve**. See **PVD-2.2: Screening for Vascular related genetic connective tissue Disorders**

PVD-6.3: Abdominal Aortic Aneurysm (AAA)

- Ultrasound abdominal aorta with any of the studies from the table of **Duplex ultrasound for visceral vascular studies** in **PVD-6.1: Aortic Disorders General Information** is the preferred imaging study to:
 - ◆ Screen for AAA and
 - ◆ Survey known AAA or
 - ◆ To evaluate a pulsatile abdominal mass
- **Obese Individual (BMI ≥ 35):** CT Abdomen and Pelvis with contrast (CPT® 74177) or without contrast (CPT® 74176) can be substituted for US using the same timeline as a non-obese individual. Ultrasound abdominal aorta should ideally first be attempted to see if the image quality is adequate
- Screening
 - ◆ One-time screening recommendations for AAA (Ultrasound CPT® 76706)
 - Individuals 65 to 75 years of age with a history of tobacco use
 - Individuals older than 75 years with a history of tobacco use and in otherwise good health who have not previously received a screening ultrasound examination
 - All first-degree relatives of individuals who present with an AAA and are between 65 and 75, or in those older than 75 in good health
 - ◆ Medicare covers a one-time AAA screening ultrasound (CPT® 76706) if there are at least **one** of the following risk factors
 - Family history of AAA
 - An individual with **all** of the following:
 - Male sex assigned at birth
 - Age is 65 to 75
 - Lifetime smoking history of at least 100 cigarettes
 - ◆ AAA screening is reasonable with ultrasound (CPT® 76706, 93975, 93976, 93978, or 93979) if there is a documented thoracic aortic aneurysm; however, there is insufficient evidence to support the use of advanced imaging to screen for a thoracic aortic aneurysm in individuals with known abdominal aortic aneurysm.
- Surveillance recommendations for AAA (CPT® 76706, 93975, 93976, 93978, 93979)
 - ◆ >2.5 cm but <3.0 cm: 10 years
 - ◆ 3.0 cm to 3.9 cm: 3 year intervals
 - ◆ 4.0 cm to 4.9 cm: every 12 months
 - ◆ 5.0 cm to 5.4 cm: every 6 months
 - ◆ >5.4 cm or aortic diameter has increased in size by 0.5 cm in six months, or at least 1 cm in a year may undergo more frequent monitoring and should be evaluated by a Vascular Specialist

- Additional Imaging
 - ◆ CT Abdomen and Pelvis with contrast (CPT® 74177), CT Abdomen and Pelvis without contrast (CPT® 74176), or CTA Abdomen and Pelvis (CPT® 74174), or CTA Abdomen (CPT® 74175), or CTA Pelvis (CPT® 72191).
 - Suspected or known AAA with recent-onset abdominal or back pain, particularly in the presence of a pulsatile epigastric mass or significant risk factors for AAA
 - Pre-operative imaging for AAA repair

PVD-6.4: Iliac Artery Aneurysm (IAA)

- Ultrasound (CPT® 76882 or CPT® 93925) for evaluation of a suspected IAA
 - ◆ CT Pelvis with contrast (CPT® 72193) if ultrasound is equivocal.
 - ◆ Ultrasound for follow-up imaging annually if an aneurysm is >2cm
- Additional Imaging
 - ◆ CT Abdomen and Pelvis with contrast (CPT® 74177), CT Abdomen and Pelvis without and with contrast (CPT® 74178), or CTA Abdomen and Pelvis (CPT® 74174) for preoperative imaging if endovascular or open repair is being considered

Practice Notes

- Isolated IAA's are rare and are typically associated with AAA
- Approximately one third to one half of isolated IAA's are bilateral at time of presentation
- Abdominal Aortic aneurysm rupture usually occurs at a diameter of 5 cm or larger, whereas common iliac aneurysms that are less than 3 cm in diameter almost never rupture.

PVD-6.5: Visceral Artery Aneurysm

- Treatment is generally indicated for aneurysms >2cm.
- Workup for suspected visceral artery aneurysm (spleen, kidney, liver or intestines) if calcifications seen on plain film imaging can include:
 - ◆ Ultrasound (CPT® 76700, 76705, 93975, 93976, 93978, or 93979), or
 - ◆ CTA Abdomen (CPT® 74175), or
 - ◆ CT Abdomen with contrast (CPT® 74160).

- Ultrasound (CPT® 76700, 76705, 93975, 93976, 93978, or 93979) **or** CTA Abdomen (CPT® 74175) **or** CT Abdomen with contrast (CPT® 74160) for further monitoring based on the intervals below or as determined by a vascular specialist or any provider in consultation with a vascular specialist:
 - ◆ Splenic artery aneurysms:
 - <20mm can be imaged every three years
 - If >25mm, they should be referred for treatment, either stent, excision or splenectomy
 - ◆ For all other visceral artery aneurysms:
 - Initial evaluation with six-month follow-up for one year
 - Further follow-up annually if no significant enlargement is seen
- CTA Abdomen (CPT® 74175), MRA Abdomen (CPT® 74185), or CT Abdomen with contrast (CPT® 74160) are indicated following stent placement at:
 - ◆ 1 month
 - ◆ 6 months
 - ◆ 12 months
 - ◆ Then every year

Practice Notes

- Splenic artery aneurysms, the most common (60%), tend to exhibit very slow rates of growth, while the other visceral artery aneurysms are more unpredictable in their rate of growth with a greater tendency to rupture.
- Visceral Artery Aneurysms are defined by an increase of more than 50% of the original arterial diameter
- Vascular specialty consultation is beneficial in order to determine the time frame to intervention

PVD-6.6: Renovascular Hypertension/Renal Artery Stenosis

Renal artery revascularization has **not** been shown to be more effective than medical therapy in most situations and should not be pursued except in extreme cases, or if there is concern for Takayasu arteritis or fibromuscular dysplasia

- MRA Abdomen without or with contrast (CPT® 74185) or CTA Abdomen with contrast (CPT® 74175) or US kidney retroperitoneal (CPT® 76775) and/or Doppler (CPT® 93975, 93976, 93978, or 93979 if expertise is available) are indicated when there is documentation of **any** of the following:
 - ◆ The individual is adherent to full doses of three blood pressure medications (including a diuretic) yet has still not achieved goal
 - ◆ Sudden and persistent worsening of previously controlled hypertension
 - ◆ Onset of hypertension younger than 30 years of age
 - ◆ Malignant hypertension with coexistent evidence of acute end-organ damage (acute renal failure, new visual or neurological disturbance and/or advanced retinopathy) or flash pulmonary edema
 - ◆ Individuals who develop hypertension ($\geq 140/90$) within the first 20 weeks of pregnancy when hypertension persists >12 weeks post-partum

- ◆ New or worsening renal function/increasing creatinine (especially after the administration of an ACE inhibitor or with angiotensin receptor blocking agent)
- ◆ Unexplained atrophic kidney or discrepancy in size between kidneys of greater than 1.5 cm
- Gadolinium agents may be contraindicated in individuals with severe renal disease or on dialysis due to the risk of developing nephrogenic systemic sclerosis
- Carotid duplex (CPT® 93880) is reasonable to screen for carotid involvement in individuals with documented or highly suspicious renal artery stenosis due to fibromuscular dysplasia (mostly women between 15 and 50 years of age).
- CTA Abdomen (CPT® 74175) or MRA Abdomen (CPT® 74185) to screen for renovascular fibromuscular dysplasia in hypertensive individuals with documented cervicocephalic fibromuscular dysplasia. The assessment of other vascular beds should be considered if supported by suggestive symptoms or medical history.

PVD-6.7: Aortic Dissection and Other Aortic Conditions

Imaging for Aortic conditions	CPT®
CT Chest, and/or Abdomen, and/or Pelvis	71260 74177 74160 72193
CTA Chest, and/or Abdomen, and/or Pelvis	71275 74175 72191 74174
MRA Chest, and/or Abdomen, and/or Pelvis	71555 74185 72198

- Classic symptoms of sharp, severe acute onset of retrosternal or interscapular chest pain is seen in 96% and is best adapted to the emergent setting. Chest x-ray is imprecise; any suspicion should be considered since up to 10% of individuals with aortic dissection present without classic symptoms.
- CTA or MRA of the entire aorta (including arch branches) and extending through the femoral arteries for suspected aortic dissection. **Any** of the following studies can be used if acute dissection is suspected:

- ◆ CT Chest (CPT® 71260 or CPT® 71270) and/or one of the following:
 - CT Abdomen (CPT® 74160 or CPT® 74170) with or without and with contrast
 - CT Pelvis (CPT® 72193 or CPT® 72194) with or without and with contrast
 - CT Abdomen and Pelvis (CPT® 74177 or CPT® 74178) with or without and with contrast
- ◆ CTA Chest (CPT® 71275) and/or **one** of the following:
 - CTA Abdomen (CPT® 74175)
 - CTA Pelvis (CPT® 72191)
 - CTA Abdomen and Pelvis (CPT® 74174)
- ◆ MRA Chest and/or Abdomen and/or Pelvis (CPT® 71555 and/or CPT® 74185 and/or CPT® 72198)
- Chronic Aortic Dissections- 1/3 of individuals with chronic type B dissections that were not treated via open or endovascular repair will go on to develop aneurysmal disease requiring subsequent intervention.
 - ◆ Advanced imaging of the affected segment of the aorta with any of the studies in above table **Imaging for Aortic conditions** can be performed as follows:
 - In individuals with a persistent false lumen or initial aortic diameter of $\geq 4\text{cm}$:
 - Every 6 months for two years until stability has been reached
 - Then annually
 - In individuals with initial aortic diameter of $< 4\text{cm}$ and/or a thrombosed false lumen:
 - Annually
 - Any time if the individual is symptomatic with chest pain, back pain or has any evidence of end organ ischemia: renal dysfunction, mesenteric ischemia or acute limb ischemia
- In individuals with Marfan syndrome/Loeys-Dietz/Ehlers-Danlos
 - ◆ As aneurysmal expansion within a dissection can occur rapidly, post-dissection imaging in these individuals is indicated as follows:
 - 1 month
 - 3 months
 - 6 months
 - 12 months
 - yearly thereafter
 - ◆ Depending on the location of the dissection the following may be approved:
 - CTA or MRA Head (CPT® 70496 or CPT® 70544)
 - Carotid duplex or CTA Neck or MRA Neck (CPT® 93980, CPT® 70498, or CPT® 70547)
 - CTA or MRA Chest (CPT® 71275 or CPT® 71555)
 - CTA Abdomen and Pelvis (CPT® 74174); or CTA or MRA Abdomen (CPT® 74175 or CPT® 74185); or CTA or MRA Pelvis (CPT® 72191 or CPT® 72198)

PVD-6.8: Post Aortic Endovascular/Open Surgery Surveillance Studies

- Aortic root/ascending aortic aneurysm repair post-operative echocardiography (TEE/TTE) can be obtained:
 - ◆ Every three months for the first year
 - ◆ Every six months during the second year
 - ◆ Annually thereafter
- For individuals who have had open descending thoracic aortic aneurysm repair, one of the following, CT Chest with contrast (CPT® 71260), CT Chest without contrast (CPT® 71250) or CTA Chest (CPT® 71275), can be obtained
 - ◆ 3-6 months postoperatively
 - ◆ 12 months postoperatively
 - ◆ And then every two years thereafter
- Open Aortic Abdominal Aneurysm Repair
 - ◆ Non-contrast enhanced CT of the entire aorta at 5-year intervals (CPT® 74176).
 - ◆ Imaging as requested to assess for suspected infection of the graft

PVD-6.8.1: Post-operative surveillance after TEVAR for any indication

Imaging for post-operative abdominal TEVAR	CPT®
CT Chest, and/or Abdomen, and/or Pelvis	71260 74177 74160 72193
CTA Chest, and/or Abdomen, and/or Pelvis	71275 74175 72191 74174
MRA Chest, and/or Abdomen, and/or Pelvis	71555 74185 72198

- ANY of the above studies listed in the table can be performed at one month, six months, twelve months and then annually
- Abdomen and Pelvis imaging is indicated only if TEVAR performed for a dissection that extends into the abdomen or pelvis

PVD-6.8.2: Post-operative surveillance after abdominal EVAR (endovascular aneurysm repair)

Imaging for post-operative abdominal EVAR	CPT®
CT Abdomen and/or Pelvis with contrast	74160 72193 74177
CT Abdomen and/or Pelvis without and with contrast	74170 72194 74178
CTA Abdomen and/or Pelvis	74175 72191 74174
MRA Abdomen and/or Pelvis	74185 72198

- CT as per above coding as requested and color duplex ultrasound (CPT® 93975, CPT® 93976, CPT® 93978, or CPT® 93979) one month after EVAR
- If no endoleak, or sac enlargement, repeat **either** preferred CT or duplex ultrasound (**but not both**) at 12 months
- If a type II endoleak is observed 1 month after EVAR, may approve **BOTH at 6 months**:
 - ◆ Any of the above CT with contrast
 - ◆ Color duplex US
- If no endoleak or AAA enlargement is detected at 1 year after EVAR annual surveillance with:
 - ◆ Color duplex US
 - ◆ If DGUS is not available, any of the above CT can be performed
- If a type II endoleak is associated with an aneurysm sac that is shrinking or stable in size:
 - ◆ Continue surveillance with color duplex US every 6 months for 2 years
 - ◆ Annually thereafter.
- If US detects a new endoleak, graft migration, or aneurysm sac growth > 5mm:
 - ◆ Any of the above CT scan as requested.
 - ◆ Non-contrast CT of the entire aorta at 5-year intervals (CPT®74176)

PVD-6.8.3: Endovascular (Stent) Iliac Repair

Imaging for endovascular iliac repair (stent)	CPT®
CT Pelvis	72193 72194
CTA Pelvis	72191
MRA Pelvis	72198

- One of the above studies can be performed for endovascular iliac repair (stent)
- If performed in conjunction with EVAR, surveillance can follow the same schedule as EVAR.
- For isolated iliac artery aneurysm repair, surveillance can be performed with an arterial duplex (CPT® 93975, CPT® 93976, CPT® 93978, or CPT® 93979) or CT or MR as above if duplex unavailable:
 - ◆ Post-operatively within the first month
 - ◆ 6 months after endovascular treatment
 - ◆ Annually

PVD-6.9: Large Vessel Vasculitis

- Large vessel vasculitis is generally sub-grouped into three areas
 - ◆ Aortitis (Inflammatory Aortitis)
 - ◆ Giant Cell Vasculitis
 - ◆ Takayasu Arteritis

PVD-6.9.1: Inflammatory Aortitis

Imaging for Inflammatory Aortitis	CPT®
CTA Chest	71275
MRA Chest	71555
CTA Pelvis	72191
MRA Pelvis	72198
CTA Abdomen and Pelvis	74174
CTA Abdomen	74175
MRA Abdomen	74185

- Initial imaging with CTA or MRA of the affected body region is considered medically necessary after the following workup:
 - ◆ Lab studies: CBC, CMP, elevated inflammatory markers such as ESR or CRP
 - ◆ Clinical history suggestive of disease listed below in practice notes
- Follow up imaging with CTA or MRA of the affected body region is considered medically necessary for:
 - ◆ Change in signs/symptoms
 - ◆ Known aneurysm monitoring
 - See **HD-12: Aneurysm and AVM** in the Head Imaging Guidelines
 - See **PVD-6.2: Thoracic Aortic Aneurysms**
 - See **PVD-6.3: Abdominal Aortic Aneurysms**

Practice Notes

- Aortitis may be congenital (Marfan's, Hypermobility Syndromes, others) or acquired, including traumatic, atherosclerotic (dissecting aneurysm, other), infectious (syphilis, tuberculosis, other), neoplastic or inflammatory (Ankylosing Spondylitis, Giant Cell Arteritis, Cogan's, Relapsing Polychondritis, Behcet's Syndrome, Polyarteritis Nodosa, Granulomatous Polyangitis, Lupoid, idiopathic, other).

PVD-6.9.2: Giant Cell Arteritis

Imaging for Giant Cell Arteritis	CPT®
CTA Chest	71275
MRA Chest	71555
CTA Pelvis	72191
MRA Pelvis	72198
CTA Abdomen and Pelvis	74174
CTA Abdomen	74175
MRA Abdomen	74185
MRA Head without contrast	70544
MRA Head with contrast	70545
MRA Neck without contrast	70547
MRA Neck with contrast	70548
CTA Neck	70498
PET/CT skull base to mid-thigh	78815

Most commonly encountered vasculitis in adults. Although classically thought of as a disease of the temporal arteries, aortic arch involvement is now recognized as a frequent complication (up to 50% of individuals) and responsible for many of the more serious morbidities encountered such as blindness.

- GCA may be subdivided into two basic types; Cranial and Extra-cranial
 - ◆ **Cranial GCA** is the more common type with temporal artery involvement. For predominantly Cranial GCA:
 - US (CPT® 93880 or CPT® 93882) of the temporal (and or axillary) arteries is the preferred modality. Ultrasound should be considered prior to advanced imaging.
 - MRA Head and/or MRA Neck (CPT® 70544, or CPT® 70545, or CPT® 70547, or CPT® 70548) may be considered when:
 - Vascular trained ultrasonography is not available
 - US is negative or equivocal with a clinical suspicion of GCA
 - For symptoms of stroke or TIA, see **HD-21: Stroke/TIA** in the Head Imaging Guidelines
 - CT and PET are **not** currently recommended for the assessment of inflammation of cranial arteries.
 - ◆ **Extra-cranial GCA**: less commonly encountered. None of the “classic” clinical signs or symptoms of cranial GCA are present initially but may develop later.
 - Extra-cranial GCA is characterized by at least two or more of the following:
 - Jaw and/or upper extremity claudication
 - Fever/weight loss or “FUO” symptoms
 - New murmurs
 - Pulse asymmetry
 - Abdominal pain
 - Pulsatile mass
 - High inflammatory markers such as CRP or ESR > 50 mm/h

- Imaging for aortic root, arch or abdomen involvement:
 - MRA Chest (CPT® 71555), MRA Neck (CPT® 70547), MRA Abdomen (CPT® 74185), CTA Chest (CPT® 71275), CTA Neck (CPT® 70498) or CTA Abdomen (CPT® 74175)
 - PET (CPT® 78815) may be indicated if MRA or CTA are non-diagnostic and there is still suspicion for aortic root, arch or abdomen involvement
- Follow up imaging is considered medically necessary for:
 - ◆ One-time documentation of remission or disease control
 - ◆ Change in signs/symptoms suggesting progression of disease
 - ◆ Although individuals with GCA can develop aortic aneurysms over time screening in the absence of signs or symptoms is not medically necessary
 - ◆ In individuals with known thoracic or abdominal aortic aneurysm:
 - See **PVD-6.2: Thoracic Aortic Aneurysm (TAA)** for thoracic aneurysm surveillance
 - See **PVD-6.3: Abdominal Aortic Aneurysm (AAA)** for abdominal aneurysm surveillance.
- Follow up imaging is not routinely recommended for individuals in clinical and biochemical remission or without aneurysm/complication.

PVD-6.9.3: Takayasu Arteritis

Imaging for Takayasu Arteritis	CPT®
CTA Chest	71275
CTA Pelvis	72191
CTA Abdomen and Pelvis	74174
CTA Abdomen	74175
MRA Chest	71555
MRA Pelvis	72198
MRA Abdomen	74185

- Initial imaging is considered medically necessary for signs and symptoms suggestive of disease such as absent radial pulse, difficulty obtaining BP in one arm, or unexplained hypertension.
- Any of the following modalities may be indicated for evaluation of Takayasu arteritis:
 - ◆ MRA of the affected body area(s) (contrast as requested)
 - ◆ CTA of the affected body area(s) (contrast as requested)
 - ◆ Ultrasound with Doppler of the affected body area(s)
- For follow-up imaging, see **PEDPVD-3.2: Large Vessel Vasculitis** in the Pediatric Peripheral Vascular Disease Imaging Guidelines.

References

1. Bonci G, Steigner ML, et al. ACR Appropriateness Criteria® Thoracic Aorta Interventional Planning and Follow-Up. *J Am Coll Radiol*. 2017;14(11S):S570-S583. doi:10.1016/j.jacr.2017.08.042.
2. Kalva SP, Dill KE, et al. ACR Appropriateness Criteria® nontraumatic aortic disease. *J Thorac Imaging*. 2014;29(5):W85-W88. doi:10.1097/RTI.0000000000000107.
3. Albornoz G, Coady MA, Roberts M, et al. Familial thoracic aortic aneurysms and dissections--incidence, modes of inheritance, and phenotypic patterns. *Ann Thorac Surg*. 2006;82(4):1400-1405. doi:10.1016/j.athoracsur.2006.04.098.
4. Anderson JL, Halperin JL, Albert N, et al. Management of Patients with Peripheral Artery Disease (Compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations). *J Am Coll Cardiol*. 2013;61(14):1555-1570. doi:10.1016/j.jacc.2013.01.004.
5. Bennett SJ, Dill KE, Hanley M, et al. ACR Appropriateness Criteria® Suspected Thoracic Aortic Aneurysm. *Journal of the American College of Radiology*. 2018;15(5). doi:10.1016/j.jacr.2018.03.031.
6. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg*. 2018;67(1). doi:10.1016/j.jvs.2017.10.044.
7. Elefteriades JA. Natural history of thoracic aortic aneurysms: indications for surgery, and surgical versus nonsurgical risks. *Ann Thorac Surg* 2002; 74: S1877-S1880.
8. Erben Y, Brownstein AJ, Rajaei S. Natural History of and Management of Splanchnic Artery Aneurysms in a Single Tertiary Referral Center. *J Vasc Surg* 2018 Oct; 68(4): 1079-1087.
9. Expert Panel on Cardiac Imaging. ACR Appropriateness Criteria® Acute Chest Pain -- Suspected Aortic Dissection. American College of Radiology (ACR); 2014.
10. Collard M, Sutphin PD, Kalva SP, et al. Expert Panel on Vascular Imaging. ACR Appropriateness Criteria® Abdominal Aortic Aneurysm Follow-up (without Repair). *J Am Coll Radiol*. 2019;16(5S):S2-S6. doi:10.1016/j.jacr.2019.02.005.
11. Corey MR, Ergul EA, Cambria RP. The Natural History of Splanchnic Aneurysms and Outcome after Operative Intervention. *J Vasc Surg*. 2016 April 63 (4):949-57.
12. Dejaco C, Ramiro S, Duftner C, et al. EULAR recommendations for the use of imaging in large vessel vasculitis in clinical practice. *Annals of the Rheumatic Diseases*. 2018;77(5):636-643. doi:10.1136/annrheumdis-2017-212649.
13. Diercks D, Promes S, Schuur J, et al. American College of Emergency Physicians. Clinical policy: critical issues in the evaluation and management of adult patients with suspected acute nontraumatic thoracic aortic dissection. *Ann Emerg Med*. 2015 Jan; 65 (1) :32-42.
14. Francois CJ, Skulborstad EP, Majdalany BS, et al. Expert Panels on Vascular Imaging and Interventional Radiology. ACR Appropriateness Criteria® Abdominal Aortic Aneurysm: Interventional Planning and Follow-Up. *J Am Coll Radiol*. 2018;15(5S):S2-S12. doi:10.1016/j.jacr.2018.03.008.
15. Harvin HJ, Verma N, Nikolaidis P, et al. ACR Appropriateness Criteria® Renovascular Hypertension. *J Am Coll Radiol*. 2017;14(11). doi:10.1016/j.jacr.2017.08.040.
16. Hiratzka LF, Bakris GL, Beckman JA, et al. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM Guidelines for the diagnosis and management of patients with thoracic aortic disease. A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, American Association for Thoracic Surgery, American College of Radiology, American Stroke Association, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of Thoracic Surgeons, and Society for Vascular Medicine [published correction appears in *J Am Coll Cardiol*. 2013 Sep 10;62(11):1039-40]. *J Am Coll Cardiol*. 2010;55(14):e27-e129. doi:10.1016/j.jacc.2010.02.015.
17. Kallianos KG, Burris NS. Imaging Thoracic Aortic Aneurysm. *Radiol Clin North Am*. 2020;58(4):721-731. doi:10.1016/j.rcl.2020.02.009.
18. Loren F, Hiratzka MD, et al, 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM Guidelines for the Diagnosis and Management of Patients with Thoracic Aortic Disease. *Circulation* 2010; 121: e266-e369.
19. Masuda Y1, Yamada Z, Morooka N, Watanabe S, Inagaki Y. Prognosis of patients with medically treated aortic dissections. *Circulation*. 1991 Nov;84(5 Suppl):III7-13.
20. Moser M, Setaro JF. Resistant or Difficult-to-Control Hypertension. *New England Journal of Medicine*. 2006;355(4):385-392. doi:10.1056/nejmcp041698.
21. Persu A, Giavarini A, Touzé E, et al. European consensus on the diagnosis and management of fibromuscular dysplasia. *Journal of Hypertension*. 2014;32(7):1367-1378. doi:10.1097/hjh.0000000000000213.
22. Sakamoto I, Sueyoshi E, Hazama S, et al. Endovascular treatment of iliac artery aneurysms. *Radiographics* October 2005;25:S213-S227.
23. Shiga T, Wajima Z, Apfel CC, Inoue T, Ohe Y. Diagnostic Accuracy of Transesophageal Echocardiography, Helical Computed Tomography, and Magnetic Resonance Imaging for Suspected Thoracic Aortic Dissection: Systematic Review and Meta-analysis. *Arch Intern Med* 2006; 166 (13): 1350-1356.
24. Smith T, Quencer KB. Best Practice Guidelines: Imaging Surveillance After Endovascular Aneurysm Repair. *American Journal of Roentgenology*. 2020;214(5):1165-1174. doi:10.2214/ajr.19.22197.

25. Tadros TM, Klein MD, Shapira OM. Ascending aortic dilatation associated with bicuspid aortic valve. *Circulation* 2009; 119: 880-890.
26. Taimen K, Salomäki SP, Hohenthal U, et al. The Clinical Impact of Using 18F-FDG-PET/CT in the Diagnosis of Suspected Vasculitis: The Effect of Dose and Timing of Glucocorticoid Treatment. *Contrast Media & Molecular Imaging*. 2019;2019:1-8. doi:10.1155/2019/9157637.
27. van Bogerijen GH, Tolenaar JL, Rampoldi V, et.al. Predictors of aortic growth in uncomplicated type B aortic dissection. *J Vasc Surg*. 2014 Apr;59(4):1134-43. doi: 10.1016/j.jvs.2014.01.042.
28. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018;71(6). doi:10.1161/hyp.0000000000000065.
29. Zierler RE, Jordan WD, Lal BK, et al. The Society for Vascular Surgery practice guidelines on follow-up after vascular surgery arterial procedures. *J Vasc Surg* . 2018;68(1):256-284. doi:10.1016/j.jvs.2018.04.018.

PVD-7: Lower Extremity Peripheral Vascular Disease	
PVD-7.1: Claudication	40
PVD-7.2: Popliteal Artery Entrapment Syndrome	41
PVD-7.3: Post-Procedure Surveillance Studies	41
PVD-7.3.1: For suprainguinal disease	42
PVD-7.3.2: For infrainguinal bypass	43
PVD-7.4: Lower Extremity Artery Aneurysms	44
PVD-7.5: Arterial Imaging for Free Flaps in Reconstructive Surgery	45

PVD-7.1: Claudication

- Resting ABI for initial evaluation of suspected PAD. This can be accomplished at the bedside as part of the physical examination or requested as CPT® 93922 (limited Doppler ultrasound) or CPT® 93923 (multi-level complete Doppler ultrasound)
 - ◆ CPT® 93923 may be performed once
 - ◆ Follow-up studies may be performed with CPT® 93922
 - ◆ Post-exercise ABI (CPT® 93924) can be performed if the resting ABI is >0.89 and PAD is still highly suspected clinically.
- History and physical suggestive of PAD include:
 - ◆ History
 - Claudication- reproducible calf or thigh cramping with exertion that is relieved completely with rest
 - Critical limb ischemia
 - Rest pain suggestive of ischemia-pain in the ball of foot when the leg is in an elevated position particularly at night
 - Distal non-healing wound or punched out ulcer with sharply demarcated edges present for >2 weeks with no evidence of healing, i.e. presence of granulation tissue
 - ◆ Physical Examination
 - Abnormal lower extremity pulse examination
 - Vascular bruit
 - Non-healing lower extremity wound
 - Lower extremity gangrene
 - Other suggestive lower extremity physical findings (e.g., elevation pallor/dependent rubor)
 - Atrophic nails, hair loss, shiny skin
- If resting ABI (CPT® 93922) is normal (0.9 to 1.3) and disease is still suspected:
 - ◆ Differentiate from “pseudoclaudication”. See **SP-9: Lumbar Spinal Stenosis** in the Spine Imaging Guidelines
 - ◆ Re-measure ABI after exercise (CPT® 93924)
 - ◆ A toe-brachial index may be used as further screening in individuals with ABI's greater than 1.3
 - ◆ Advanced imaging is necessary only if there is consideration for invasive therapy not to confirm diagnosis
- Duplex ultrasound (CPT® 93925 bilateral study or CPT® 93926 unilateral study) and Doppler studies are adjuncts to abnormal ABI that may be used to identify location and extent of disease once there has been a decision for revascularization.

- MRA Aorta and Pelvic vessels, and Lower extremities (CPT® 74185, CPT® 73725 and CPT® 73725), **or** CTA with run-off (CPT® 75635) to further evaluate the lower extremity arteries for the purpose of preoperative planning for ANY of the following:
 - ◆ Intermittent claudication (i.e. non-limb threatening ischemia) AND either:
 - Failed 3 months' conservative medical therapy (physician supervised walking/exercise program plus medical therapy), **or**
 - Functional disability (e.g. exercise impairment sufficient to threaten the individual's employment or to require significant alterations in the individual's lifestyle)
 - ◆ Potentially limb-threatening vascular disease evidenced by:
 - Skin breakdown
 - Non-healing ischemic ulcers
 - Resting leg pain
 - Gangrene
 - ◆ Blue Toe Syndrome:
 - Emboli from aortic plaque or mural thrombus
 - Hyperviscosity syndrome
 - Hypercoagulable states
 - Vasculitis
 - ◆ Note: MRA Pelvis (CPT® 72198) should not be requested/billed with CPT® 74185, CPT® 73725 and CPT® 73725

Practice Notes

Claudication symptoms usually remain stable (70% to 80% of individuals) and do not worsen or improve at rapid rates. Repeat studies to assess the efficacy of medical therapy are not indicated unless there is a negative change in clinical status for the purpose of preoperative planning such as worsening claudication or progression to critical limb ischemia

PVD-7.2: Popliteal Artery Entrapment Syndrome

- Diagnosis of popliteal artery stenosis or occlusion due to compression by adjacent muscle and tendons seen in young men (ages 20 to 40).
 - ◆ Ultrasound (CPT® 93926 unilateral study), CTA Lower extremity (CPT® 73706), or MRA Lower extremity (CPT® 73725).
 - ◆ CT or MRI Lower Extremity (contrast as requested) if requested by the operating surgeon

PVD-7.3: Post-Procedure Surveillance Studies

- Scheduled Interval
 - ◆ ABI (CPT® 93922) following any revascularization procedure
 - ABI (CPT® 93922) or Duplex ultrasound (CPT® 93926 unilateral study) at each routine follow up is indicated generally after a history/physical has been performed
 - Further imaging studies such as CTA or MRA are indicated for worsening symptoms, an abnormal duplex or a significant reduction (>0.15) in the ABI

Indication	Imaging
➤ Suprainguinal Revascularization Both Open and Endovascular Therapy including Aortobifem/iliofem/fem-fem bypass/iliac angioplasty/stent	➤ Clinical examination and ABI (CPT® 93922) with arterial duplex (CPT® 93978 or CPT® 93979) at: ◆ 1 month ◆ 6 months ◆ 12 months ◆ Then annually
➤ Infrainguinal Open Revascularization (Femoral-popliteal, femoral-tibial, femoral-distal bypass) ◆ With vein or autologous conduit	➤ Clinical exam and ABI (CPT® 93922) with arterial duplex (CPT® 93978 or CPT® 93979) ◆ Post-operatively ◆ 3 months ◆ 6 months ◆ 12 months ◆ Then annually
◆ With Prosthetic conduit (PTFE/Dacron)	➤ Clinical exam and ABI (CPT® 93922) with arterial duplex (CPT® 93978 or CPT® 93979) ◆ Post-operatively ◆ 6 months ◆ 12 months ◆ Then annually
➤ Infrainguinal Endovascular Revascularization Femoropopliteal angioplasty/stent	➤ Clinical exam and ABI (CPT® 93922) with arterial duplex (CPT® 93978 or CPT® 93979) ◆ 1 month ◆ 3 month ◆ Every 6 months for two years ◆ Then annually

PVD-7.3.1: For suprainguinal disease

- One of the following studies: Arterial duplex, CTA Abdomen and Pelvis, CT Abdomen and Pelvis with contrast, CTA Aorta with lower extremity runoff, MRI Abdomen and Pelvis, MRA Abdomen and Pelvis, or MRA Aorta with lower extremity runoff for:
 - ◆ Worsening signs or symptoms
 - ◆ Reduction of ABI >0.15
 - ◆ Peak systolic velocities or PSV ratio suggestive of high grade stenosis or in-stent re-stenosis

PVD-7.3.2: For infrainguinal bypass

- CTA Lower Extremity or MRA Lower Extremity for:
 - ◆ Worsening signs or symptoms
 - ◆ Reduction of ABI>0.15
 - ◆ Duplex suggestive of threatened graft
- If intervention was performed for a non-healing wound and wound has gone on to heal, no additional imaging is recommended for surveillance. Repeat arterial duplex imaging can be obtained for worsening clinical signs and symptoms such as the presence of a new wound or rest pain

References

1. Dick F, Ricco J-B, Davies A.H, Cao P, et. al. Chapter VI: Follow-up after Revascularization. *European Journal of Vascular and Endovascular Surgery*, Volume 42, Supplement 2, 2011 pp.S75-90.
2. Lane TRA et al. Post-Operative Surveillance after Open Peripheral Arterial Surgery. *Eur J Vasc Endovasc Surg*. Volume 42, Issue 1, July 2011. Pages 59-77.
3. Brooke BS, Beck AW, Kraiss LW, et al. Association of Quality Improvement Registry Participation with Appropriate Follow-up after Vascular Procedures. *JAMA Surgery*. 2018;153(3):216. doi:10.1001/jamasurg.2017.3942.
4. Rooke TW, Hirsch AT, Misra S, Sidawy AN, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society for Vascular Medicine, and Society for Vascular Surgery. American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines., Society for Cardiovascular Angiography and Interventions., Society of Interventional Radiology., Society for Vascular Medicine., Society for Vascular Surgery. *J Vasc Surg*. 2011 Nov; 54(5):e32-58
5. Conte MS, Pomposelli FB, and Clair DG. Society for Vascular Surgery practice guidelines for atherosclerotic occlusive disease of the lower extremities: Management of asymptomatic disease and claudication. *J Vasc Surg*. March 2015;61(3):2S-41S.e1
6. Bui TD, Mills JL Sr, Inhnat DM et al. The natural history of duplex-detected stenosis after femoropopliteal endovascular therapy suggests questionable clinical utility of routine duplex surveillance. *J Vasc Surg*. 2012 Feb 55:2:346-352
7. Troutman DA, Madden NJ, Dougherty MJ, et al. Duplex Ultrasound Diagnosis of Failing Stent Grafts Placed for Occlusive Disease. *J Vasc Surg* 2014;70:1–5.
8. Humphries MD, Pevec WC, Laird JR, et.al. Early Duplex Scanning after Infringuinal Endovascular Therapy. *J Vasc Surg* 2011; 53:353–8.
9. Zierler RE, Jordan WD, Lal BK, et al. The Society for Vascular Surgery practice guidelines on follow-up after vascular surgery arterial procedures. *J Vasc Surg*. 2018 Jul;68(1):256-284. doi: 10.1016/j.jvs.2018.04.018.
10. Cooper K, Majdalany BS, Kalva SP, et al. ACR Appropriateness Criteria[®] Lower Extremity Arterial Revascularization—Post-Therapy Imaging. *J Am Coll Radiol*. 2018;15:S104-S115.
11. Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the management of patients with lower extremity peripheral artery disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2017; 135: e726-e779.
12. Ahmed O, Hanley M, Bennett SJ, et al. ACR Appropriateness Criteria[®] Vascular claudication -Assessment for Revascularization. *J Am Coll Radiol*. 2017 May;14(5S):S372-S379. doi: 10.1016/j.jacr.2017.02.037.
13. Macedo TA, Johnson M, Hallett Jr JW, and Breen JF. Popliteal artery entrapment syndrome: role of imaging in the diagnosis. *AJR Am J Roentgen*. 2003 Nov;181(5):1259-1265. doi:10.2214/ajr.181.5.1811259.

PVD-7.4: Lower Extremity Artery Aneurysms

- For Iliac artery aneurysm see **PVD-6.4: Iliac Artery Aneurysm (IAA)**
- Femoral artery aneurysm
 - ◆ Initial imaging
 - Ultrasound (CPT® 93925 bilateral study or CPT® 93926 unilateral study).
 - ◆ Surveillance imaging
 - Symptomatic true femoral aneurysms smaller than 2.5 cm in diameter
 - Ultrasound (CPT® 93926 unilateral study) annually
 - Symptomatic true femoral aneurysms larger than 2.5 cm
 - Ultrasound (CPT® 93926 unilateral study) every 6 months
 - ◆ Other imaging
 - CTA Lower extremity or MRA Lower extremity without or with contrast when:
 - Preoperative study for individuals with no plans for invasive angiography
 - Technically limited or abnormal ultrasound results
- Popliteal artery aneurysm
 - ◆ Initial imaging
 - Ultrasound (CPT® 93925 bilateral study or CPT® 93926 unilateral study) and Ultrasound to assess for a contralateral popliteal aneurysm and abdominal aortic aneurysm (CPT® 76770 or CPT® 76775)
 - ◆ Surveillance imaging
 - Ultrasound (CPT® 93926 unilateral study) annually
 - Post-interventional functional testing (ABI) (CPT® 93922) may be useful as clinically indicated
 - ◆ Other imaging
 - CTA or MRA for:
 - Preoperative study for individuals with no plans for invasive angiography
 - Technically limited or abnormal ultrasound results

References

1. Hall HA, Minc S, Babrowski T. Peripheral artery aneurysm. *Surg Clin North Am.* 2013;93(4):911-ix. doi:10.1016/j.suc.2013.04.008.
2. von Stumm M, Teufelsbauer H, Reichenspurner H, Debus ES. Two Decades of Endovascular Repair of Popliteal Artery Aneurysm--A Meta-analysis. *Eur J Vasc Endovasc Surg.* 2015;50(3):351-359. doi:10.1016/j.ejvs.2015.04.036
3. Kim TI, Sumpio BE. Management of Asymptomatic Popliteal Artery Aneurysms. *Int J Angiol.* 2019;28(1):5-10. doi:10.1055/s-0038-1676792.
4. Corriere MA, Guzman RJ. True and false aneurysms of the femoral artery. *Semin Vasc Surg.* 2005;18(4):216-223. doi:10.1053/j.semvascsurg.2005.09.008

PVD-7.5: Arterial Imaging for Free Flaps in Reconstructive Surgery

- For breast reconstruction preoperative planning. See **BR-3: Breast Reconstruction** in the Breast Imaging Guidelines
- For head and neck reconstruction, CTA or MRA lower extremity (CPT® 73706 or 73725) may be approved for evaluation of perforator anatomy for planned fibular flap

References

1. Garvey, PB et al. A Prospective Study of Preoperative Computed Tomographic Angiographic Mapping of Free Fibula Osteocutaneous Flaps for Head and Neck Reconstruction. *Plastic and Reconstructive Surgery*. 2013;130(4):541e-49e.
2. Smith JM et al. An overview of methods for vascular mapping in the planning of free flaps. *J Plast Reconstr Aesthet Surg* 2010 Sep;63(9):e674-82.
3. Abou-Foul AK, et al. Anatomical variants of lower limb vasculature and implications for free fibula flap: Systematic review and critical analysis. *Microsurgery*. 2016 Feb;36(2):165-72.

PVD-8: Imaging for Hemodialysis Access

PVD-8.1: Arterial Evaluation and Venous Mapping Prior to AV Fistula	47
PVD-8.2: Hemodialysis access imaging	47

PVD-8.1: Arterial Evaluation and Venous Mapping Prior to AV Fistula

- Imaging prior to AV fistula creation:
 - ◆ For vessel mapping CPT® 93985 or 93986
 - ◆ MRA Upper Extremity (CPT® 73225) may be needed if duplex imaging is equivocal
- Arterial evaluation to assess arterial suitability (size, degree of stenosis and calcification) prior to AV fistula creation may be indicated
 - ◆ CPT® 93930 or CPT® 93931 can be used to report upper extremity arterial evaluation
- Venous mapping (CPT® 93970 or CPT® 93971) to assess venous suitability prior to AV fistula creation may be indicated

PVD-8.2: Hemodialysis access imaging

- Indications for Duplex ultrasound (CPT® 93990) of hemodialysis access include but are not limited to:
 - ◆ Individuals with decreased flow rates during hemodialysis.
 - ◆ Development of arm swelling or discomfort after access placement surgery or a hemodialysis session.
 - ◆ Prolonged immaturity of a surgically created AV fistula.
 - ◆ Suspected pseudoaneurysm.
 - ◆ Suspected AV fistula or graft stenosis.
 - ◆ Known or suspected fluid collection adjacent to an AV fistula or graft.
 - ◆ One Duplex US (CPT® 93990) can be performed after a surgically created AV fistula for assessment, although it is not generally needed.
- Central venous stenosis can cause new dialysis access to fail to mature or cause the premature failure of existing fistulas/grafts.
- CT Chest with contrast (CPT® 71260), or CTA Chest (CPT® 71275), or MRA Chest (CPT® 71555) is indicated when there is documentation of either:
 - ◆ Signs and symptoms of central venous stenosis including:
 - Arm swelling
 - Presence of numerous collateral veins
 - Prolonged bleeding from dialysis puncture sites
 - ◆ A history of pacemaker placement or previous tunneled dialysis graft, regardless of signs and symptoms.

References

1. Sidawy AN, Spergel LM, Besarab A, et al. The Society for Vascular Surgery: Clinical practice guidelines for the surgical placement and maintenance of arteriovenous hemodialysis access. *J Vasc Surg*. 2008;48(5). doi:10.1016/j.jvs.2008.08.042.
2. Schmidli J, Widmer MK, Basile C, et al. Editor's Choice – Vascular Access: 2018 Clinical Practice Guidelines of the European Society for Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg*. 2018;55(6):757-818. doi:10.1016/j.ejvs.2018.02.001
3. Kundu S. Central venous disease in hemodialysis patients: prevalence, etiology and treatment. *Journal of Vascular Access*. 2010;11(1):1-7.

PVD-9: Arteriovenous Malformations (AVMs)

PVD-9.1: Arteriovenous Malformations (AVMs)

Arteriovenous malformations are characterized by a network of multiple abnormal vascular channels interposed between enlarged feeding arteries and draining veins. The arteriovenous fistula has a single communication interposed between a feeding artery and a draining vein. The normal capillary bed is absent in both lesions. Both lesions may have an aggressive clinical course and are characterized by a reddish pulsatile mass which has a thrill or bruit. Though often recognized at birth, these lesions may grow and present near adolescence.

- See **PEDPVD-2.5: Arteriovenous Malformations (AVMs) and Fistulas** in the Pediatric Peripheral Vascular Disease Imaging Guidelines
- Ultrasound with Doppler is indicated as an initial examination for superficial lesions.
 - ◆ Large lesion characterization may be limited by ultrasound imaging window.
 - ◆ Ultrasound is also limited in evaluating AVM relationship to airway or bony structures.
- MRI without contrast or without and with contrast of the affected body part is also indicated for evaluation of AVMs, and is useful in evaluating the extent of AVMs and their relationship to normal structures.
- MRA (contrast as requested) of the affected body part can be approved for evaluation and surveillance of known AVMs.
- It is unusual for both MRI and MRA to be necessary for routine treatment response or surveillance imaging of AVMs, but both may be approved for preoperative planning.
- CT and CTA can also be used to characterize AVMs and their relationship to normal structures, but is generally not better than MRI and has associated radiation risks.
 - ◆ CT with contrast and/or CTA (contrast as requested) of the affected body part can be approved when MRI and/or MRA is inconclusive or contraindicated.

References

1. Pizzo PA, Poplack DG, Krishnamurthy R, Daldrop-Link HE, Jones JY, et. al. Imaging studies in the diagnosis and management of pediatric malignancies. In: Principles and Practice of Pediatric Oncology. Vol 7. Philadelphia: Wolters Kluwer; 2016:185-234.
2. Martin KL. Vascular disorders. *Nelson Textbook of Pediatrics*, Chapter 669. eds Kliegman R, St. Geme JW III, Blum NJ, et al. 21st ed. Philadelphia, PA: Elsevier; 2020:3461-3469.
3. Blei F, Guarini A. Current workup and therapy of infantile hemangiomas. *Clinics in Dermatology*. 2014;32(4):459-470. doi:10.1016/j.clindermatol.2014.02.001.
4. Cahill AM, Nijs ELF. Pediatric Vascular Malformations: Pathophysiology, Diagnosis, and the Role of Interventional Radiology. *CardioVascular and Interventional Radiology*. 2011;34(4):691-704. doi:10.1007/s00270-011-0123-0.
5. Bagrodia N, Defnet AM, Kandel JJ. Management of lymphatic malformations in children. *Current Opinion in Pediatrics*. 2015;27(3):356-363. doi:10.1097/mop.0000000000000209.
6. Wassef M, Blei F, Adams D, et al. Vascular Anomalies Classification: Recommendations from the International Society for the Study of Vascular Anomalies. *Pediatrics*. 2015;136(1):e203-e214. doi:10.1542/peds.2014-3673.
7. Kutz AM, Aranibar L, Lobos N, Wortsman X. Color Doppler Ultrasound Follow-Up of Infantile Hemangiomas and Peripheral Vascularity in Patients Treated with Propranolol. *Pediatric Dermatology*. 2015;32(4):468-475. doi:10.1111/pde.12596.
8. Adams DM, Trenor CC, Hammill AM, et al. Efficacy and Safety of Sirolimus in the Treatment of Complicated Vascular Anomalies. *Pediatrics*. 2016;137(2). doi:10.1542/peds.2015-3257.
9. Snyder E, Puttgen K, Mitchell S, Ahlawat S, Tekes A. Magnetic Resonance Imaging of the Soft Tissue Vascular Anomalies in Torso and Extremities in Children: An Update with 2014 International Society for the Study of Vascular Anomalies Classification. *Journal of Computer Assisted Tomography*. 2017;42(2):167-177. doi:10.1097/rct.0000000000000675.
10. Merrow AC, Gupta A, Patel MN, Adams DM. 2014 Revised Classification of Vascular Lesions from the International Society for the Study of Vascular Anomalies: Radiologic-Pathologic Update. *RadioGraphics*. 2016;36(5):1494-1516. doi:10.1148/rg.2016150197.
11. Johnson CM, Navarro OM. Clinical and sonographic features of pediatric soft-tissue vascular anomalies part 1: classification, sonographic approach and vascular tumors. *Pediatric Radiology*. 2017;47(9):1184-1195. doi:10.1007/s00247-017-3885-y.
12. Johnson CM, Navarro OM. Clinical and sonographic features of pediatric soft-tissue vascular anomalies part 2: vascular malformations. *Pediatric Radiology*. 2017;47(9):1196-1208. doi:10.1007/s00247-017-3906-x.
13. Sadick M, Müller-Wille R, Wildgruber M, Wohlgemuth W. Vascular Anomalies (Part I): Classification and Diagnostics of Vascular Anomalies. *RöFo - Fortschritte auf dem Gebiet der Röntgenstrahlen und der bildgebenden Verfahren*. 2018;190(09):825-835. doi:10.1055/a-0620-8925.
14. Olivieri B, White CL, Restrepo R, et. al. Low-Flow Vascular Malformation Pitfalls: From Clinical Examination to Practical Imaging Evaluation—Part 2, Venous Malformation Mimickers. *American Journal of Roentgenology*. 2016;206(5):952-962. doi:10.2214/ajr.15.15794.
15. White CL, Olivieri B, Restrepo R, et.al. Low-Flow Vascular Malformation Pitfalls: From Clinical Examination to Practical Imaging Evaluation—Part 1, Lymphatic Malformation Mimickers. *American Journal of Roentgenology*. 2016;206(5):940-951. doi:10.2214/ajr.15.15793.

PVD-10: Nuclear Medicine

PVD-10.1: Nuclear Medicine Imaging indications

- Nuclear medicine studies are rarely used in the evaluation of peripheral vascular disorders, but are indicated in the following circumstances:
 - ◆ Lymphoscintigraphy (CPT® 78195) is indicated for evaluation of lower extremity lymphedema when a recent Doppler ultrasound is negative for valvular insufficiency.
 - ◆ Vascular flow imaging (CPT® 78445) is an obsolete study that has been replaced by MRA, CTA, or Duplex ultrasonography, and is not supported for any indication at this time.
 - ◆ Venous thrombosis imaging (CPT® 78456, CPT® 78457, and CPT® 75458) are obsolete studies that have been replaced by MRA, CTA, or Duplex ultrasonography, and are not supported for any indication at this time.
 - ◆ Indium 111 (¹¹¹In)–labeled white blood cell (WBC) or Gallium-67 citrate studies (CPT® 78800, CPT® 78801, CPT® 78802, or CPT® 78803) can be approved for evaluation of the following:
 - Mycotic aneurysms.
 - Vascular graft infection.
 - Infection of central venous catheter or other indwelling device.

PVD-11: Venous Imaging General Information

PVD-11.1: Abbreviations and glossary	52
PVD-11.2: Background Information	52
PVD-11.3: Procedure Coding	54
PVD-11.4: General Guidelines-Imaging	55

PVD-11.1: Abbreviations and glossary

Abbreviation	Definition
CTV	Computed Tomography Venography
DVT	Deep Venous Thrombosis
EVA	Endovenous ablation – a minimally invasive procedure using heat to obliterate the saphenous vein for the treatment of venous reflux
IVC	Inferior vena cava
May-Thurner's	Syndrome of compression of the left iliac vein via an overlying right common iliac artery. The pulsations of the artery into the vein against the 5 th lumbar vertebrae can predispose to DVT
MRV	Magnetic Resonance Venography
Phlebectomy	Removal of a vein usually through a small incision
Post thrombotic syndrome	Constellation of symptoms including chronic edema and pain that develops after a DVT
Sclerotherapy	Injection of an irritant into a vein to obliterate it
SEPS	Sub-fascial endoscopic perforator surgery
SVT	Superficial venous thrombosis
VVI	Venous Valvular Insufficiency – a study utilizing ultrasound to assess for the presence of reflux within the superficial and deep veins of the lower extremity.

PVD-11.2: Background Information

- A current clinical evaluation (within 60 days), including medical treatments, are required prior to considering advanced imaging, which includes:
 - ◆ Relevant history and physical examination including:
 - The affected limb(s), the extent of the edema (calf and/or thigh), pitting or non-pitting. With regard to venous insufficiency, presence or absence of hyperpigmentation or other skin changes, ulcerations if applicable, size of varicosities if present as well as distribution
 - Arterial examination to rule out phlegmasia alba/cerulea dolens which is comprised arterial flow secondary to extensive DVT if applicable
 - Appropriate laboratory studies, for example d-dimer, if applicable
 - Non-advanced imaging modalities, such as a venous duplex or venous valvular insufficiency study (VVI) after symptoms started or worsened
 - ◆ Other meaningful contact (telephone call, electronic mail or messaging) by an established patient can substitute for a face-to-face clinical evaluation.
- Venous disease can be classified into three categories:
 - ◆ Venous occlusive disease
 - ◆ Venous insufficiency
 - ◆ Venous malformations

Veno-occlusive disease

- Types of thrombotic disease:
 - ◆ Superficial venous thrombosis
 - ◆ Deep venous thrombosis
 - ◆ Iliac vein obstruction, unilateral or bilateral
 - ◆ May-Thurner's syndrome
- Signs/Symptoms of veno-occlusive disease is generally sudden onset of pain and edema in the limb.
- Risk factors include age >40, obesity, pregnancy, prolonged immobility, post-surgery, and malignancy among others.
- Procedures related to veno-occlusive disease include:
 - ◆ Thrombolysis
 - ◆ Thrombectomy
 - ◆ Post iliac vein stent/angioplasty See **PVD-17: Post iliac vein stent/angioplasty**

Venous insufficiency

- Types of venous insufficiency:
 - ◆ Superficial and deep venous reflux
 - ◆ Varicose veins
 - ◆ Reticular and spider veins
- Signs/symptoms of venous insufficiency include:
 - ◆ Chronic swelling in the leg that is relieved with elevation
 - ◆ Chronic swelling in the leg that is worse in the evenings
 - ◆ Aching or sense of heaviness in the leg
 - ◆ Hyperpigmentation of the calf particularly around the ankle
 - ◆ Itchy skin on legs and feet
 - ◆ Leather appearance of the skin of the calves
 - ◆ Skin ulcers in the calf particularly around the medial malleolus
 - ◆ Varicose veins
 - ◆ Spider veins/reticular veins/telangiectasias
- Procedures related to the venous insufficiency include:
 - ◆ Endovenous laser ablation utilizing either chemical, laser or radio-frequency
 - ◆ Saphenous vein high ligation and stripping
 - ◆ Phlebectomy, stab or powered
 - ◆ Sclerotherapy, liquid or foam

Venous malformations

- Types of venous malformations include
 - ◆ Arterio-venous malformations which can occur throughout the body
 - ◆ See **CH-24: Pulmonary AVM** in the Chest imaging guidelines
 - ◆ See **HD-12: Aneurysm and AVM** in the Head imaging guidelines
 - ◆ See **PV-11: Pelvic Pain/Dyspareunia, Female** in the Pelvic imaging guidelines
 - ◆ Klippel-Trenaunay which affects primarily the lower extremity venous circulation and is characterized by varicose veins, limb size discrepancies, and port-wine stains.
- Treatment includes:
 - ◆ Primarily embolization
 - ◆ Sclerotherapy
 - Klippel-Trenaunay: treatment can include phlebectomy and sclerotherapy of symptomatic varicose veins provided they meet the criteria for intervention.

PVD-11.3: Procedure Coding

Venous Studies – Extremities		CPT®
CTV Abdomen and Pelvis involves obtaining images from the diaphragm to just below the inguinal ligament after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.		74174
CTV Pelvis involves obtaining images from the top of the pelvic brim to the upper thighs or just below the inguinal ligament. The venogram portion is performed by obtaining images after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.		72191
MRV Abdomen and Pelvis involves taking images from the diaphragm to just below the inguinal ligament after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.		74185
MRV Pelvis involves obtaining images from the top of the pelvic brim to the upper thighs or just below the inguinal ligament. The venogram portion is performed by obtaining images after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.		72198
Duplex scan of extremity veins, including responses to compression and other maneuvers; complete bilateral study.		93970
Duplex scan of extremity veins, including responses to compression and other maneuvers; unilateral or limited study.		93971
➤ These codes are used to report studies of lower or upper extremity veins. ➤ A complete bilateral study of the lower extremity veins includes examination of the external iliac veins, common femoral, proximal deep femoral, great saphenous and popliteal veins. Calf veins may also be included. ➤ A complete bilateral study of upper extremity veins includes examination of the subclavian, jugular, axillary, brachial, basilic, and cephalic veins. Forearm veins may also be included.		
Duplex scan of aorta, inferior vena cava, iliac vasculature, or bypass grafts; complete study		93978
Duplex scan of extremity veins, including responses to compression and other maneuvers; unilateral or limited study.		93979

PVD-11.4: General Guidelines-Imaging

- Venous duplex (CPT® 93970, CPT® 93971) of the limb is the initial imaging of choice
 - ◆ Follow-up duplex imaging (CPT® 93970, CPT® 93971) is not generally indicated to document resolution and should only be obtained for new signs/symptoms or for concerns of propagation of thrombus when the treatment plan would change (Insertion of IVC filter, change of anticoagulation, etc.)
- Imaging studies
 - ◆ Venous duplex (CPT® 93970, CPT® 93971) should visualize the veins, with demonstration of the presence or absence of compressibility and venous flow.
 - ◆ Venous valvular insufficiency studies (CPT® 93970, CPT® 93971) visualize the veins of the lower extremity, assess for reflux (reversal of venous antegrade flow after valve closure) and measure its duration.
 - ◆ CTV or MRV of the Abdomen and Pelvis images with contrast involves taking images from the diaphragm to just below the inguinal ligament after a delay of a few minutes after IV contrast is administered to optimize filling and therefore visualization of the venous vasculature.

References

1. Ageno W, Haas S, Weitz JI, et al. Characteristics and Management of Patients with Venous Thromboembolism: The GARFIELD-VTE Registry. *Thromb Haemost.* 2019;119(2):319-327. doi:10.1055/s-0038-1676611.
2. Jones WS, Vemulapalli S, Parikh KS et al. Treatment Strategies for Patients with Lower Extremity Chronic Venous Disease (LECVD. Agency for Healthcare Research and Quality (US); 2017 Apr.
3. Gloviczki P, Comerota AJ, Dalsing MD, et al. The care of patients with varicose veins and associated chronic venous diseases: Clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum. *J Vasc Surg.* 2011. 53(16S):1-47. doi:10.1016/j.jvs.2011.01.079.
4. Wolpert L, Rahmani O, Stein B, et al. Magnetic Resonance Venography in the Diagnosis and Management of May-Thurner syndrome. *Vasc Endovascular Surg.* 2002 Jan-Feb;36(1):51-7. doi:10.1177/153857440203600109.

PVD-12: Acute Limb Swelling

PVD-12.1: Superficial venous thrombosis (SVT)	57
PVD-12.2: Acute deep venous thrombosis (DVT)	57
PVD-12.3: Follow-up imaging of known DVT	58
PVD-12.4: Follow-up imaging after venous surgery	58
PVD-12.5: Generalized bilateral lower extremity edema	58

PVD-12.1: Superficial venous thrombosis (SVT)

- Superficial venous thrombosis (SVT) refers to acute or chronic thrombosis of the superficial veins in both the upper (cephalic and basilic veins) and lower extremities (greater [great] saphenous vein, lesser [small] saphenous vein, gastrocnemius and soleal veins). Treatment: Elevation and warm compresses until pain and swelling subsides.
- The diagnosis of superficial venous thrombosis is generally made on the basis of physical examination.
 - ◆ Duplex ultrasound (CPT® 93970, CPT® 93971) is the initial imaging if the diagnosis is equivocal
 - ◆ Follow-up duplex ultrasound (CPT® 93970, CPT® 93971) is indicated only if thrombus in the superficial systems is encroaching onto the deep venous system (saphenofemoral or saphenopopliteal junction)

PVD-12.2: Acute deep venous thrombosis (DVT)

- Deep venous thrombosis is characterized by thrombosis of a deep vein in either the upper (brachial, axillary, subclavian veins) or the lower extremity (peroneal, posterior tibial, popliteal, femoral or iliac veins).
- Duplex ultrasound (CPT® 93970 bilateral study or CPT® 93971 unilateral study) is the initial imaging study for any suspected DVT
 - ◆ Deep venous thrombosis can present as
 - Symptomatic
 - Swelling
 - Pain
 - Warmth
 - Erythema
 - Pain with dorsiflexion of the foot (Homan's Sign)
 - Or with progression, such as phlegmasia cerulea dolens
 - Risk factors for DVT include age >40, obesity, malignancy, prolonged immobilization, hypercoagulability as well as those outlined in **CH-25: Pulmonary Embolism (PE)** in Chest Imaging Guidelines.
- CTA/CTV Abdomen and pelvis with contrast can be performed to rule out IVC thrombus secondary to the filter when there is acute bilateral lower extremity swelling in an individual with a history of an IVC filter in place.
- When there is concern for proximal DVT (iliofemoral):
 - ◆ Focused abdominal duplex can generally visualize the iliac veins and IVC to determine the absence or presence of iliac vein thrombus in an individual. If the results are equivocal or indeterminate:
 - CTV or MRV Abdomen and Pelvis with contrast (CPT® 74174 or CPT® 74185 and CPT® 73725) can be performed.
- For request concerning abdominal vein thrombosis, see **AB-43.2: Abdominal Veins other than Hepatic and Portal Veins** in the Abdomen Imaging Guidelines
- For proximal DVT's (iliac vein DVT's or in cases of phlegmasia (extensive DVT compromising arterial inflow), thrombectomy (rarely performed) or thrombolysis can be performed.

- If the cause of the DVT is found to be due to May-Thurner, iliac vein angioplasty followed by stenting of the left iliac vein is generally performed. See **PVD-13.3: May-Thurner Syndrome**

PVD-12.3: Follow-up imaging of known DVT

- Duplex ultrasound (CPT® 93970 bilateral study or CPT® 93971 unilateral study) can be repeated in order to rule out proximal extension of a calf vein DVT in those individuals who cannot be anticoagulated, most commonly after recent surgery. Time interval for follow-up study includes:
 - ◆ One week after the initial diagnosis.
 - ◆ Serial imaging (up to 3 studies) over the first three weeks if calf DVT is not treated.
- Imaging during or to terminate long-term anticoagulation therapy to determine venous recanalization is not supported by evidence. Repeat imaging to make decisions on whether or not to continue or terminate anticoagulation is not indicated.

PVD-12.4: Follow-up imaging after venous surgery

- Venous duplex (CPT® 93971 unilateral study) of the treated limb is indicated to rule out a DVT within seven days of endovenous ablation.
- Follow-up routine imaging is **not indicated** after other venous procedures including:
 - ◆ Saphenous vein ligation and stripping
 - ◆ Phlebectomy
 - ◆ Sclerotherapy

PVD-12.5: Generalized bilateral lower extremity edema

Bilateral lower extremity edema is multifactorial. Prior to any request for advanced imaging, a workup for causes of the edema should be instituted including echocardiogram to rule out congestive heart failure and laboratory studies to exclude renal insufficiency and liver disease. The following imaging is indicated based on the suspected cause of the edema:

- Suspected abdominal or pelvic pathology
 - ◆ Abdominal ultrasound or duplex is the initial imaging
 - ◆ CT Abdomen and Pelvis or CT pelvis either with or without contrast can be performed if abdominal US is equivocal or indeterminate
- Suspected chronic venous insufficiency
 - ◆ A venous duplex CPT® 93970 (bilateral) or CPT® 93971 (unilateral) is indicated to evaluate for venous reflux.
- Suspected lymphedema
 - ◆ When initial noninvasive studies, such as ultrasound, are negative for venous valvular insufficiency **either** of the following advanced imaging studies is indicated:
 - Lymphoscintigraphy (CPT® 78195)
 - MRI lymphangiography (CPT® 73718)

References

1. Lim W, Gal GL, Bates SM, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Advances*. 2018;2(22):3226-3256. doi:10.1182/bloodadvances.2018024828.
2. American College of Phlebology. Treatment of Superficial Venous Disease of the Lower Leg Guidelines. American Vein & Lymphatic Society. <http://www.phlebology.org/member-resources/publications/treatment-superficial-venous-disease-lower-leg-guidelines>. Published 2016.
3. American College of Phlebology. Duplex Ultrasound Imaging of Lower Extremity Veins in Chronic Venous Disease. American Vein & Lymphatic Society. <http://www.phlebology.org/member-resources/publications/duplex-ultrasound-imaging-lower-extremity-veins-chronic-venous-disease>. Published 2012.
4. Healy DA, Kimura S, Power D, et al. A Systematic Review and Meta-analysis of Thrombotic Events Following Endovenous Thermal Ablation of the Great Saphenous Vein. *European Journal of Vascular and Endovascular Surgery*. 2018;56(3):410-424. doi:10.1016/j.ejvs.2018.05.008.
5. Jacobs CE, Pinzon MM, Orozco J, et al. Deep Venous Thrombosis after Saphenous Endovenous Radiofrequency Ablation: Is it Predictable? *Annals of Vascular Surgery*. 2014;28(3):679-685. doi:10.1016/j.avsg.2013.08.012.
6. Liu N, Wang C, Sun M. Noncontrast three-dimensional magnetic resonance imaging vs lymphoscintigraphy in the evaluation of lymph circulation disorders: A comparative study. *J Vasc Surg*. 2005;41(1):69-75. doi:10.1016/j.jvs.2004.11.013.

PVD-13: Chronic limb swelling due to chronic deep venous thrombosis/May Thurner's syndrome

PVD-13.1: Chronic deep venous thrombosis general information	61
PVD-13.2: Post thrombotic syndrome	61
PVD-13.3: May-Thurner syndrome general information	62

PVD-13.1: Chronic deep venous thrombosis general information

- Chronic deep venous thrombosis is defined as an acute DVT that is greater than 14 days old.
- Individual with incompletely lysed or residual DVT can develop post-thrombotic syndrome that can be characterized as chronic edema, venous stasis changes, pain and in advanced cases venous stasis ulceration.
 - ◆ Incompletely lysed DVT can cause luminal narrowing of the vein restricting venous outflow leading to stenosis or occlusion and /or can lead to valve dysfunction resulting in reflux of venous blood retrograde towards gravity. Both pathologies ultimately lead to chronic edema which can cause chronic pain and venous stasis disease.
 - Imaging for post-thrombotic syndrome is not indicated unless there are signs and symptoms suggestive of a new acute DVT or for preoperative planning for iliac vein/stenting for suspected iliac vein stenosis or occlusion.
 - The mainstay of treatment for chronic deep venous thrombosis is compression stockings.
 - ◆ In individuals with a history of proximal (iliofemoral) DVT who have subsequently developed post thrombotic syndrome, imaging can be performed to evaluate for iliac venous obstruction which can result from incompletely lysed thrombus.
 - Initial imaging should be duplex (CPT® 93970 bilateral study or CPT® 93971 unilateral study) followed by either a CT or MR venogram Abdomen and Pelvis, or CT or MR venogram Pelvis, or venography for treatment planning purposes.
 - Selected individuals may be a candidate for iliac vein angioplasty/stenting.

PVD-13.2: Post thrombotic syndrome

- Imaging for post-thrombotic syndrome is indicated when:
 - ◆ There are signs and symptoms suggestive of a new acute DVT and NOT for chronic swelling that has not changed in severity or character
 - ◆ For preoperative planning for iliac vein/stenting in the setting of known iliac venous obstruction in those with a history of a proximal (iliofemoral) DVT.
- See **PVD-13.1: Chronic deep venous thrombosis general information**
- See **PVD-13.3: May-Thurner syndrome general information**
- See **PVD-15: Venous Stasis Ulceration**

PVD-13.3: May-Thurner syndrome general information

In approximately 25% of people, the right iliac artery overlies the left iliac vein over the fifth lumbar vertebrae and its pulsations can compress the vein increasing the risk of DVT in the left extremity.

- CTV or MRV Abdomen and Pelvis (CPT® 74174, or 74185 and 72198) OR CTV Pelvis or MRV Pelvis (CPT® 72191 or 72198) can be approved in individuals with a history of
 - ◆ Left lower extremity iliac DVT
 - ◆ Persistent left lower extremity edema OR varicose veins OR venous stasis ulcer despite treatment of superficial venous disease in that extremity
 - ◆ Persistent left lower extremity edema OR varicose vein OR venous stasis ulcer in the absence of saphenous vein reflux.
- Treatment is with iliac vein angioplasty/stenting
- Prophylactic treatment of May-Thurner's syndrome in the absence of acute or chronic DVT **OR** chronic left lower extremity edema and its sequelae such as varicose veins or venous stasis ulcers is **NOT** considered medically necessary.

References

1. Eberhardt RT and Raffetto JD. Chronic Venous Insufficiency. *Circulation*. 2005; 111: 2398-2409.
2. Partsch H, Flour M, Smith PC, et al. Indications for compression therapy in venous and lymphatic disease. Consensus based on experimental data and scientific evidence. *Int Angiol*. 2008;27:193-219.
3. Nayak L, Vedantham S. Multifaceted Management of the Postthrombotic Syndrome. *Seminars in Interventional Radiology*. 2012; 29:1.
4. Christof T, Kaltenmeier, Erben Y, et al. Systematic review of May-Thurner syndrome with emphasis on gender differences. *J Vasc Surg Venous Lymphat Disord*. 2018;6(3):399-407.
5. Rossi FH, Kambara AM, Izukawa NM, et al. Randomized double-blinded study comparing medical treatment versus iliac vein stenting in chronic venous disease. *J Vasc Surg Venous Lymphat Disord*. 2018;6(2):183-191.
6. Sermathanasawadi N, Pruekprasert K, Pitaksantayothin W, et al. Prevalence, risk factors, and evaluation of ilio caval obstruction in advanced chronic venous insufficiency. *J Vasc Surg Venous Lymphat Disord*. 2019 May;7(3):441-447.
7. Vedantham S, Kahn SR, Goldhaber SZ, Comerota AJ, et al. Endovascular therapy for advanced post-thrombotic syndrome: Proceedings from a multidisciplinary consensus panel. *Vasc Med*. 2016 Aug;21(4):400-7.

PVD-14: Chronic limb swelling due to venous insufficiency/Venous stasis changes/Varicose veins

PVD-14.1: Venous insufficiency - General	64
PVD-14.2: Venous Reflux	64
PVD-14.3: Varicose Veins	65
PVD-14.4: Spider veins/reticular veins	65

PVD-14.1: Venous insufficiency – General information

- Venous insufficiency is characterized by failure of the venous blood to flow in its normal antegrade path of flow and instead reflux backwards by the force of gravity usually secondary to malfunction of the venous valves.
- Risk factors include previous DVT, obesity, female sex assigned at birth, hereditary, and environmental factors such as prolonged standing on a hard surface.
- Venous insufficiency loosely includes the diagnosis of venous reflux, varicose veins, venous stasis ulcers and spider/reticular veins.

PVD-14.2: Venous Reflux

- Diagnosis is made with a venous valvular insufficiency study (CPT® 93970 bilateral study or CPT® 93971 unilateral study) which documents the presence of reflux (>500ms) in the greater saphenous vein as well as the size of the refluxing vein (3-15mm).
- Symptoms of venous reflux include chronic edema, pain, and venous stasis ulcerations. Symptoms of venous reflux can be ameliorated with compression therapy with graded compression stockings, elevation, avoidance of prolonged standing and weight loss. Venous reflux can be seen in both the deep and superficial venous systems. Reflux within the deep system is not amenable to intervention.
 - ◆ Treatment of deep venous reflux is via active compression with compression stocks, pneumatic pumps or specialized dressings such as Unna boots.
 - ◆ Treatment of superficial venous reflux is amenable to intervention in selected individuals who are symptomatic and have failed conservative therapy. A duplex ultrasound (CPT® 93970 bilateral study or CPT® 93971 unilateral study) demonstrating the presence of pathologic reflux within the greater and lesser saphenous veins should be undertaken within the last six months. Vein size should be documented.
 - ◆ Treatment of symptomatic superficial venous reflux is via endovenous laser radiofrequency ablation of the greater or lesser saphenous vein resulting in closure of the vein allowing for venous blood to be rerouted to the deep venous system.
 - ◆ Treatment of symptomatic superficial venous reflux can also be treated via saphenous vein ligation and stripping which has fallen out of favor but can be performed for a tortuous or enlarged (>15mm) greater or lesser saphenous vein. One complication of endovenous ablation is deep venous thrombosis.
 - ◆ A post ablation venous ultrasound (CPT® 93970 bilateral study or CPT® 93971 unilateral study) is indicated within seven days post procedure. If thrombus is noted within the saphenofemoral junction, repeat imaging can be performed within seven days to assess for propagation into the deep system.
 - ◆ Ultrasound mapping or monitoring techniques are considered medically necessary only to initially determine the extent and configuration of symptomatic varicosities or valvular insufficiency. Post procedure assessment by imaging techniques is inappropriate to confirm efficacy or outcome of the procedure.

PVD-14.3: Varicose Veins

- Varicose veins are defined as enlarged, tortuous veins visible under the skin. Symptoms associated with varicose veins include achiness and heaviness of the legs and pain/discomfort over the varicosities. Varicose veins can exist both in the absence and presence of venous reflux.
- Treatment involves conservative therapy such as compression stockings, avoidance of prolonged standing, intermittent elevation, weight loss (if applicable) and exercise which relieves the distention of the varicose veins ameliorating the symptoms.
- If the varicosities remain symptomatic despite conservative therapy, varicose veins are treated with sclerotherapy or phlebectomy generally on the basis of size.

PVD-14.4: Spider veins/reticular veins

- Spider veins are formed by the dilation of a cluster of blood vessels within the dermis – generally <3mm in diameter. Diagnosis is via physical examination. Spider veins are usually asymptomatic but can cause aching, burning and tenderness in the area overlying the abnormal veins. Spider veins can exist in the absence or presence of venous reflux. The presence of spider veins should not be an indication for treatment of venous reflux.
- Treatment of spider veins is generally cosmetic except in certain cases and can be treated with sclerotherapy.

References

1. Gloviczki P, Comerota AJ, Dalsing MC, et al. The care of patients with varicose veins and associated chronic venous diseases: clinical practice guidelines of the Society for Vascular Surgery and the American Venous Forum. *J Vasc Surg.* 2011;53(5 Suppl):2S-48S. doi:10.1016/j.jvs.2011.01.079.
2. Wittens C, Davies A, Bækgaard N, et al. Editor's Choice – Management of Chronic Venous Disease. *Eur J Vasc Endovasc Surg.* 2015;49(6):678-737. doi:10.1016/j.ejvs.2015.02.007.
3. Piazza G. Varicose veins. *Circulation.* 2014;130(7):582-587. doi:10.1161/CIRCULATIONAHA.113.008331.
4. Jones WS, Vemulapalli S, Parikh KS, et al. Treatment Strategies for Patients with Lower Extremity Chronic Venous Disease (LECVD). Technology Assessment Program. Project ID: DVTT0515. (Prepared by the Duke University Evidence-based Practice Center under Contract No. 290-2015-00004-I). Rockville, MD: Agency for Healthcare Research and Quality; April 2017.
5. National Institute for Health and Care Excellence (NICE). Varicose veins in the legs: the diagnosis and management of varicose veins. NICE Clinical Guideline: Methods, evidence and recommendations. London: National Institute for Health and Care Excellence (UK). July, 2013.
6. Kahn SR, M'lan CE, Lamping DL, et al. Relationship between clinical classification of chronic venous disease and patient-reported quality of life: results from an international cohort study. *J Vasc Surg.* 2004;39(4):823-828. doi:10.1016/j.jvs.2003.12.007.
7. Khilnani NM. Duplex ultrasound evaluation of patients with chronic venous disease of the lower extremities. *AJR Am J Roentgenol.* 2014;202(3):633-642. doi:10.2214/AJR.13.11465.
8. King T, Coulomb G, Goldman A, et al. Experience with concomitant ultrasound guided foam sclerotherapy and endovenous laser treatment in chronic venous disorder and its influence on Health Related Quality of Life: interim analysis of more than 1000 consecutive procedures. *International Angiology.* 2009;28 (4):289 -297.
9. Lurie F, Kistner RL. Trends in patient reported outcomes of conservative and surgical treatment of primary chronic venous disease contradict current practices. *Annals of Surgery.* 2011 Aug;254(2):363-367. DOI: 10.1097/sla.0b013e31821d4a5f.
10. Meissner MH, Moneta G, Burnand K, et al. The hemodynamics and diagnosis of venous disease. *J Vasc Surg.* 2007;46 Suppl S:4S-24S. doi:10.1016/j.jvs.2007.09.043.
11. Smith JJ, Garratt AM, Guest M, Greenhalgh RM, Davies AH. Evaluating and improving health-related quality of life in patients with varicose veins. *J Vasc Surg.* 1999;30(4):710-719. doi:10.1016/s0741-5214(99)70110-2.

PVD-15: Venous stasis ulceration

PVD-15.1: Venous stasis ulcers - General	67
PVD-15.2: Venous stasis ulcers - Treatment	67

PVD-15.1: Venous stasis ulcers - General

- Venous stasis ulcers can arise from maceration of the skin in individuals with venous insufficiency often with minimal trauma. The area over the medial malleolus is usually the most commonly affected area. The presence of chronic edema from either venous reflux, post-thrombotic syndrome or either etiology predisposes to the formation of venous stasis ulcerations.

PVD-15.2: Venous stasis ulcers - Treatment

- The mainstay of treatment is a sterile dressing +/- adjunctive wound care salves coupled with compression with either stockings or wraps to reduce edema.
- In select individuals with venous stasis ulcers felt to be due to superficial venous reflux, incompetent perforators, and/or significant varicosities, the following may be indicated:
 - ◆ Endovenous ablation with RF or with laser for treatment of saphenous vein reflux and incompetent perforators
 - ◆ Saphenous vein ligation and stripping for treatment of saphenous vein reflux and varicose veins
 - ◆ Phlebectomy for treatment of varicose veins
 - ◆ Sclerotherapy for treatment of incompetent perforators and varicose veins

References

1. Abdul-Haqq R, Almaroof B, Chen BL, Panneton JM, Parent FN. Endovenous Laser Ablation of Great Saphenous Vein and Perforator Veins Improves Venous Stasis Ulcer Healing. *Annals of Vascular Surgery*. 2013;27(7):932-939. doi:10.1016/j.avsg.2012.09.014.
2. Poblete H, Elias S. Venous Ulcers: New Options in Treatment: Minimally Invasive Vein Surgery. *The Journal of the American College of Certified Wound Specialists*. 2009;1(1):12-19. doi:10.1016/j.jcws.2008.10.004.
3. Lawrence PF, Alktaifi A, Rigberg D, Derubertis B, Gelabert H, Jimenez JC. Endovenous ablation of incompetent perforating veins is effective treatment for recalcitrant venous ulcers. *J Vasc Surg*. 2011;54(3):737-742. doi:10.1016/j.jvs.2011.02.068.
4. Harlander-Locke M, Lawrence PF, Alktaifi A, Jimenez JC, Rigberg D, Derubertis B. The impact of ablation of incompetent superficial and perforator veins on ulcer healing rates. *J Vasc Surg*. 2012;55(2):458-464. doi:10.1016/j.jvs.2011.08.054.

PVD-16: IVC filters

PVD-16.1: IVC filters - General	69
PVD-16.2: IVC filters - Treatment	69

PVD-16.1: IVC filters - General

- IVC filters are placed in individuals with known DVT that cannot be anti-coagulated, individuals with poor pulmonary reserve and high risk for DVT, or prophylaxis in trauma and surgical individuals.
- Most IVC filters inserted are retrievable and should be removed as soon as clinically feasible. After 12 months, removal of IVC filters can become technically more difficult.

PVD-16.2: IVC filters - Treatment

- IVC filter insertion
 - ◆ An initial venous duplex can be performed to assess for the presence of thrombus in the femoral vein which would affect the approach (transjugular or transfemoral)
 - ◆ Advanced imaging is not indicated
- CT Abdomen and Pelvis with contrast CPT® 74177 for ANY of the following:
 - ◆ A KUB demonstrates tilting of the filter or malposition of one of the filter thongs
 - ◆ New **bilateral** lower extremity swelling (venous duplex should be performed first)
 - ◆ Filter present for >12 months, with documentation stating intent to remove

PVD-17: Post iliac vein stent/angioplasty

PVD-17.1: Post iliac vein stenting/angioplasty

Iliac venous stents can be placed after thrombolysis for DVT associated with May-Thurner's syndrome, DVT associated with extrinsic compression or for post thrombotic iliac obstruction.

- Arterial duplex (CPT® 93975, 93976, 93978, 93979) can be obtained for:
 - ◆ Surveillance of iliac venous stents
 - ◆ Worsening signs or symptoms including increased edema when stent malfunction is suspected
 - ◆ Postoperatively within the first month, at six months, twelve months and then annually
- CTV or MRV Abdomen and Pelvis can be obtained for an abnormal or indeterminate duplex

References

1. Washington State Health Care Authority. Selected Treatments for Varicose Veins, A Health Technology Assessment. Prepared by Hayes, Inc. Final Report: April 2017.
2. Proebstle TM, Alm BJ, Göckeritz, et al. Five-year results from the prospective European multicenter cohort study on radiofrequency segmental thermal ablation for incompetent great saphenous veins. *BJS*. 2015; 102: 212-218. doi:10.1002/bjs.9679.
3. Weiss RA, Weiss MA, Eimpunth S, et al. Comparative Outcomes of Different Endovenous Thermal Ablation Systems on Great and Small Saphenous Vein Insufficiency: Long-Term Results. *Lasers in Surgery and Medicine*. 2015; 47: 156-160. Aortic Disorders General Information.
4. Brittenden J, Cotton SC, Elders A, et al. Clinical effectiveness and cost-effectiveness of foam sclerotherapy, endovenous laser ablation and surgery for varicose veins: results from the Comparison of LAser, Surgery and foam Sclerotherapy (CLASS) randomised controlled trial. *Health Technol Assess*. 2015;19(27):1-342. doi:10.3310/hta19270.
5. Biemans AA, Kockaert M, Akkersdijk GP, et al. Comparing endovenous laser ablation, foam sclerotherapy, and conventional surgery for great saphenous varicose veins. *J Vasc Surg*. 2013;58(3):727-34.e1. doi:10.1016/j.jvs.2012.12.074.
6. van der Velden SK, Biemans AA, De Maeseneer MG, et al. Five-year results of a randomized clinical trial of conventional surgery, endovenous laser ablation and ultrasound-guided foam sclerotherapy in patients with great saphenous varicose veins. *Br J Surg*. 2015;102(10):1184-1194. doi:10.1002/bjs.9867.
7. Rasmussen L, Lawaetz M, Serup J, et al. Randomized clinical trial comparing endovenous laser ablation, radiofrequency ablation, foam sclerotherapy, and surgical stripping for great saphenous varicose veins with 3-year follow-up. *J Vasc Surg Venous Lymphat Disord*. 2013;1(4):349-356. doi:10.1016/j.jvsv.2013.04.008.
8. Brittenden J, Cotton SC, Elders A, et al. A randomized trial comparing treatments for varicose veins. *N Engl J Med*. 2014; 371(13): 1218-1227. doi:10.1056/NEJMoa1400781.
9. Venermo M, Saarinen J, Eskelinen E, et al. Randomized clinical trial comparing surgery, endovenous laser ablation and ultrasound-guided foam sclerotherapy for the treatment of great saphenous varicose veins. *Br J Surg*. 2016;103(11):1438-1444. doi:10.1002/bjs.10260.