



CLINICAL GUIDELINES

Musculoskeletal Imaging Guidelines

Version 1.0.2022

Effective January 1, 2022



eviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Imaging requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

eviCore's Clinical Review Criteria ("CRC") and related content is made available for the limited uses of: reference; and individual use, only limited to facilitating the determination of medically necessary and appropriate clinical treatment by clinicians for specific delegated patients under their care. The CRC and related content is proprietary information of eviCore, and copyrighted to the full extent of the law. Except as expressly permitted, you may not modify, copy, reproduce, republish, upload, post, transmit, hyperlink to or from, or distribute in any way the CRC, nor may you sell, transfer, distribute, assign, lease, reproduce, or otherwise use the CRC in commerce, in a manner that competes with us or infringes upon our rights, or for any public or commercial endeavor without our prior and express written consent.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2021 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Musculoskeletal Imaging Guidelines	
Procedure Codes associated with Musculoskeletal Imaging	3
MS-1: General Guidelines	5
MS-2: Imaging Techniques	8
MS-3: 3D Rendering	13
MS-4: Avascular Necrosis (AVN)/Osteonecrosis	14
MS-5: Fractures	17
MS-6: Foreign Body	21
MS-7: Ganglion Cysts	23
MS-8: Gout/Calcium Pyrophosphate Deposition Disease (CPPD)/ Pseudogout/Chondrocalcinosis	25
MS-9: Infection/Osteomyelitis	27
MS-10: Soft Tissue Mass or Lesion of Bone	30
MS-11: Muscle/Tendon Unit Injuries/Diseases	33
MS-12: Osteoarthritis	36
MS-13: Chondral/Osteochondral Lesions	39
MS-14: Osteoporosis	41
MS-15: Rheumatoid Arthritis (RA) and Inflammatory Arthritis	43
MS-16: Post-Operative Joint Replacement Surgery	46
MS-17: Limb Length Discrepancy	49
MS-18: Anatomical Area Tables – General Information	51
MS-19: Shoulder	52
MS-20: Elbow	58
MS-21: Wrist	63
MS-22: Hand	68
MS-23: Pelvis	70
MS-24: Hip	72
MS-25: Knee	78
MS-26: Ankle	87
MS-27: Foot	92
MS-28: Nuclear Medicine	97

Procedure Codes associated with Musculoskeletal Imaging	
MRI/MRA	CPT®
MRI Upper Extremity, other than joint, without contrast	73218
MRI Upper Extremity, other than joint, with contrast	73219
MRI Upper Extremity, other than joint, without and with contrast	73220
MRI Upper Extremity, any joint, without contrast	73221
MRI Upper Extremity, any joint, with contrast	73222
MRI Upper Extremity, any joint, without and with contrast	73223
MR Angiography Upper Extremity without or with contrast	73225
MRI Lower Extremity, other than joint, without contrast	73718
MRI Lower Extremity, other than joint, with contrast	73719
MRI Lower Extremity, other than joint, without and with contrast	73720
MRI Lower Extremity, any joint, without contrast	73721
MRI Lower Extremity, any joint, with contrast	73722
MRI Lower Extremity, any joint, without and with contrast	73723
MR Angiography Lower Extremity without or with contrast	73725
MRI Pelvis without contrast	72195
MRI Pelvis with contrast	72196
MRI Pelvis without and with contrast	72197
CT/CTA	CPT®
CT Upper Extremity without contrast	73200
CT Upper Extremity with contrast	73201
CT Upper Extremity without and with contrast	73202
CT Angiography Upper Extremity without and with contrast	73206
CT Lower Extremity without contrast	73700
CT Lower Extremity with contrast	73701
CT Lower Extremity without and with contrast	73702
CT Angiography Lower Extremity without and with contrast	73706
CT Pelvis without contrast	72192
CT Pelvis with contrast	72193
CT Pelvis without and with contrast	72194
Ultrasound	CPT®
Ultrasound, complete joint (ie, joint space and peri-articular soft tissue structures) real-time with image documentation	76881
Ultrasound, limited, joint or other nonvascular extremity structure(s) (e.g., joint space, peri-articular tendon[s], muscle[s], nerve[s], other soft tissue structure[s], or soft tissue mass[es]), real-time with image documentation	76882
Ultrasound, pelvic (nonobstetric), real time with image documentation	76857

Nuclear Medicine	CPT®
Bone Marrow Imaging, Limited	78102
Bone Marrow Imaging, Multiple	78103
Bone Marrow Imaging, Whole Body	78104
Bone or Joint Imaging Limited	78300
Bone or Joint Imaging Multiple	78305
Bone Scan Whole Body	78306
Bone Scan 3 Phase Study	78315
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, single area (e.g., head, neck, chest, pelvis), single day imaging	78800
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, 2 or more areas (eg, abdomen and pelvis, head and chest), 1 or more days imaging or single area imaging over 2 or more days	78801
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); planar, whole body, single day imaging	78802
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT), single area (e.g., head, neck, chest, pelvis), single day imaging	78803
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT) with concurrently acquired computed tomography (CT) transmission scan for anatomical review, localization and determination/detection of pathology, single area (e.g., head, neck, chest, pelvis), single day imaging	78830
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT), minimum 2 areas (e.g., pelvis and knees, abdomen and pelvis), single day imaging, or single area imaging over 2 or more days	78831
Radiopharmaceutical localization of tumor, inflammatory process or distribution of radiopharmaceutical agent(s) (includes vascular flow and blood pool imaging, when performed); tomographic (SPECT) with concurrently acquired computed tomography (CT) transmission scan for anatomical review, localization and determination/detection of pathology, minimum 2 areas (e.g., pelvis and knees, abdomen and pelvis), single day imaging, or single area imaging over 2 or more days	78832

MS-1: General Guidelines

MS-1.0: General Guidelines

6

MS-1.0: General Guidelines

- Before advanced diagnostic imaging can be considered, there must be an initial face-to-face clinical evaluation as well as a clinical re-evaluation after a trial of failed conservative treatment; the clinical re-evaluation may consist of a face-to-face evaluation or other meaningful contact with the provider's office such as email, web or telephone communications.
- A face-to-face clinical evaluation is required to have been performed within the last 60 days before advanced imaging can be considered. This may have been either the initial clinical evaluation or the clinical re-evaluation.
- The initial face-to-face clinical evaluation should include a relevant history and physical examination, appropriate laboratory studies, and non-advanced imaging modalities. Other forms of meaningful contact (e.g., telephone call, electronic mail or messaging) are not acceptable as an initial evaluation.
- Prior to advanced imaging consideration, the results of plain x-rays performed after the current episode of symptoms started or changed is required for all musculoskeletal conditions, unless otherwise noted in the guidelines.
 - ◆ Initial plain x-ray can rule out those situations that do not often require advanced imaging, such as osteoarthritis, acute/healing fracture, dislocation, osteomyelitis, acquired/congenital deformities, and tumors of bone amenable to biopsy or radiation therapy (in known metastatic disease), etc.
 - ◆ X-ray may provide complementary clinical information regarding detailed bony anatomy, and may assist with preoperative planning when surgery is being contemplated.
 - ◆ X-ray may provide clinically significant details for soft tissue masses, such as soft tissue calcification, presence or absence of phleboliths, radiographic density, and effect on adjacent bone.
 - ◆ X-ray often has a larger field of view than MRI or CT and has the potential to identify more proximal or distal pathology in an extremity.
- Clinical re-evaluation is required prior to consideration of advanced diagnostic imaging to document failure of significant clinical improvement following a recent (within 3 months) six week trial of provider-directed conservative treatment. Clinical re-evaluation can include documentation of a face-to-face encounter or documentation of other meaningful contact with the requesting provider's office by the patient (e.g. telephone call, electronic mail or messaging).
- Provider-directed conservative treatment may include rest, ice, compression, and elevation (R.I.C.E.), non-steroidal anti-inflammatories (NSAIDs), narcotic and non-narcotic analgesic medications, oral or injectable corticosteroids, viscosupplementation injections, a provider-directed home exercise program, cross-training, and/or physical/occupational therapy or immobilization by splinting/casting/bracing.

- Orthopedic specialist evaluation can be helpful in determining the need for advanced imaging.
 - ◆ The need for repeat advanced imaging should be carefully considered and may not be indicated if prior imaging has been performed.
 - ◆ Serial advanced imaging, whether CT or MRI, for surveillance of healing or recovery from musculoskeletal disease is not supported by the medical evidence in the majority of musculoskeletal conditions.

References

1. Reinus WR. *Clinician's guide to diagnostic imaging*. New York, NY: Springer Science; 2014.
2. Visconti AJ, Biddle J, Solomon M. Follow-up imaging for vertebral osteomyelitis a teachable moment. *JAMA*. 2014;174(2):184. doi: 10.1001/jamainternmed.2013.12742.
3. Fabiano V, Franchino G, Napolitano M, Ravelli A, Dilillo D, Zuccotti GV. Utility of magnetic resonance imaging in the follow-up of children affected by acute osteomyelitis. *Curr Pediatr Res*. 2017;21(2):354-358.

MS-2: Imaging Techniques

MS-2.1: Plain X-Ray	9
MS-2.2: MRI or CT	9
MS-2.3: Ultrasound	10
MS-2.4: Contrast Issues	10
MS-2.5: Positron Emission Tomography (PET)	10
MS-2.6: Nuclear Medicine	10

MS-2.1: Plain X-Ray

- The results of an initial plain x-ray are required prior to advanced imaging in all musculoskeletal conditions/disorders, unless otherwise noted in the guidelines, to rule out those situations that do not often require advanced imaging, such as osteoarthritis, acute/healing fracture, dislocation, osteomyelitis, acquired/congenital deformities, and tumors of bone amenable to biopsy or radiation therapy (in known metastatic disease), etc.

MS-2.2: MRI or CT

- Magnetic Resonance Imaging (MRI) is often the preferred advanced imaging modality in musculoskeletal conditions because it is superior in imaging the soft tissues and can also define physiological processes in some instances [e.g. edema, loss of circulation (AVN), and increased vascularity (tumors)].
- Computed Tomography (CT) is preferred for imaging cortical bone anatomy; thus, it is useful for studying complex fractures (particularly of the joints), dislocations, and assessing delayed union or non-union of fractures, if plain X-rays are equivocal. CT may be the procedure of choice in patients who cannot undergo an MRI, such as those with pacemakers.
- Bilateral Imaging
 - ◆ Coding for bilateral imaging may vary from health care plan to health care plan. Not all coding options may be available for all health care plans.

Positional MRI

Positional MRI is also referred to as dynamic, standing, weight-bearing, or kinetic MRI. Currently, there is inadequate scientific evidence to support the medical necessity of this study. As such, it should be considered experimental or investigational.

Positional CT

Positional CT, also referred to as weight-bearing or cone beam CT, may be useful in imaging of the foot and ankle.

- If a request for foot or ankle imaging with positional CT meets medical necessity criteria for standard CT imaging (as defined in the condition-specific guidelines), the request may be approved.
 - ◆ Positional CT of anatomic areas other than the foot and ankle are considered experimental or investigational.

dGEMRIC Evaluation of Cartilage

Delayed gadolinium enhanced Magnetic Resonance Imaging of Cartilage (dGEMRIC) is a technique where an MRI estimates joint cartilage glycosaminoglycan content after penetration of the contrast agent in order to detect cartilage breakdown. Currently, there is inadequate scientific evidence to support the medical necessity of this study. As such, it should be considered experimental or investigational for the diagnosis and surveillance of, or preoperative planning related to chondral pathology.

MS-2.3: Ultrasound

- Ultrasound (US) uses sound waves to produce images that can be used to evaluate a variety of musculoskeletal disorders. As with US in general, musculoskeletal US is highly operator-dependent, and proper training and experience are required to perform consistent, high quality evaluations.

MS-2.4: Contrast Issues

- Most musculoskeletal imaging (MRI or CT) is without contrast; however, the following examples may be considered with contrast:
 - ◆ Tumors, osteomyelitis, and soft tissue infection (without and with contrast)
 - ◆ MRI arthrography (with contrast only)
 - ◆ MRI for rheumatoid arthritis and inflammatory arthritis (contrast as requested)
 - ◆ For patients with a contrast contraindication, if the advanced imaging recommendation specifically includes contrast, the corresponding advanced imaging study without contrast may be approved as an alternative, although the non-contrast study may not provide an adequate evaluation of the condition of concern.

MS-2.5: Positron Emission Tomography (PET)

- At the present time, there is inadequate evidence to support the medical necessity of PET for the routine assessment of musculoskeletal disorders. It should be considered experimental or investigational.
 - ◆ See also: **MS-16: Post-Operative Joint Replacement Surgery**

MS-2.6: Nuclear Medicine

- A bone scan is a nuclear medicine imaging study in which an amount of radioactive material is injected and images are obtained at different time intervals, depending on the condition. A bone scan is done to reveal problems with bone metabolism. Areas where bone cells are repairing themselves show the most activity. It can help diagnose a number of bone conditions, including cancer of the bone or metastasis, location of bone inflammation, fracture, and bone infection.
- Nuclear Medicine WBC Scan is performed using radioactive material which is tagged to the white blood cells. When injected into the body, the material attaches to sites of inflammation/infection. Once distributed in these areas, the sites of suspected infection/inflammation can be seen on nuclear imaging equipment. These can be imaged as a planar study, SPECT study, or SPECT/CT study.
- Bone Marrow Imaging is used in combination with a WBC Scan to help differentiate between true infection and physiological marrow uptake. The bone marrow scan provides a map of the normal physiological white cell uptake that is then compared to the white blood cell scan. Any discordance in white cell uptake (e.g., more WBC uptake than marrow uptake) between the two studies indicates a focus of infection.
- See: **MS-28: Nuclear Medicine** and condition specific guidelines for specific indications.

References

1. DeMuro JP, Simmons S, Smith K, et al. Utility of MRI in blunt trauma patients with a normal cervical spine CT and persistent midline neck pain on palpation. *Global Journal of Surgery*. 2013;1(1):4-7. doi:10.12691/js-1-1-2.
2. Hsu W, Hearty TM. Radionuclide imaging in the diagnosis and management of orthopaedic disease. *J Am Acad Orthop Surg*. 2012;20(3):151-159. doi:10.5435/JAAOS-20-03-151.
3. Kayser R, Mahlfeld K, Heyde CE. Partial rupture of the proximal Achilles tendon: a differential diagnostic problem in ultrasound imaging. *Br J Sports Med*. 2005;9(11):838–842. doi:10.1136/bjsm.2005.018416.
4. Ward RJ, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute hip pain-suspected fracture. *Am Coll Radiol (ACR)*; Date of Origin: 2013. <https://acsearch.acr.org/docs/3082587/Narrative/>.
5. Mosher TJ, Kransdorf MJ, Adler R, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute trauma to the ankle. *Am Coll Radiol (ACR)*; Date of Origin: 2013. <https://acsearch.acr.org/docs/69436/Narrative/>.
6. Small KM, Adler RS, Shah SH, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Atraumatic. *Am Coll Radiol (ACR)*; New 2018. <https://acsearch.acr.org/docs/3101482/Narrative/>.
7. Amini B, Beckmann NM, Beaman FD, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Traumatic. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69433/Narrative/>.
8. Hayes CW, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic elbow pain. *Am Coll Radiol (ACR)*; Date of Origin:1998. Last Review: 2017. <https://acsearch.acr.org/docs/69423/Narrative/>.
9. Wise JN, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic foot pain. *Am Coll Radiol (ACR)*; Date of Origin:1998. Last Review: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
10. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic hip pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
11. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic wrist pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
12. Bennett DL, Nelson JW, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® nontraumatic knee pain. *Am Coll Radiol (ACR)*;1995. Last Review: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
13. Murphey MD, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® osteonecrosis of the hip. *Am Coll Radiol (ACR)*;Date of Origin: 1995. Last Review: 2015. <https://acsearch.acr.org/docs/69420/Narrative/>.
14. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® acute hand and wrist trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
15. Bencardino JT, Stone TJ, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® stress (fatigue/insufficiency) fracture, including sacrum, excluding other vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
16. Luchs JS, Flug JA, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic ankle pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
17. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot. *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/%203094201/Narrative/>.
18. Kransdorf MJ, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® suspected osteomyelitis of the foot in patients with diabetes mellitus. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2019. <https://acsearch.acr.org/docs/69340/Narrative/>.
19. Zoga AC, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® soft-tissue masses. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
20. Morrison WB, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® primary bone tumors. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2019. <https://acsearch.acr.org/docs/69421/Narrative/>.
21. Weissman BN, Palestro CJ, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® imaging after total hip arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin:1998. Last Review: 2015. <https://acsearch.acr.org/docs/3094200/Narrative/>.
22. Hochman MG, Melenevsky YV, Metter DF, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® imaging after total knee arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69430/Narrative/>.

23. Gyftopoulos S, Rosenberg ZS, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® imaging after shoulder arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/3097049/Narrative/>.
24. Patel ND, Broderick DF, Burns J, et al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria®: low back pain. *Am Coll Radiol (ACR)*; Date of Origin:1996. Last Review: 2015. <https://acsearch.acr.org/docs/69483/Narrative/>.
25. Shetty VS, Reis MN, Aulino JM, et al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria®: head trauma. *Am Coll Radiol (ACR)*; Date of Origin:1996. Last Review: 2020. <https://acsearch.acr.org/docs/69481/Narrative/>.
26. Li X, Yi P, Curry EJ, Murakami AM. Ultrasonography as a diagnostic, therapeutic, and research tool in orthopaedic surgery. *J Am Acad Orthop Surg*. 2018;26(6):187-196. doi: 10.5435/JAAOS-D-16-00221.
27. de Cesar Netto C, Myerson MS, Day J, et. al. Consensus for the use of weightbearing CT in the assessment of progressive collapsing foot deformity. *Foot Ankle Int*. 2020;41(10):1277-1282.
28. Conti MS, Ellis SJ. Weight-bearing ct scans in foot and ankle surgery. *J Am Acad Orthop Surg*. 2020;28(14):e595-e603.
29. de Cesar Netto C, Schon LC, Thawait GK, et. al. Flexible adult acquired flatfoot deformity: comparison between weight-bearing and non-weight-bearing measurements using cone-beam computed tomography. *J Bone Joint Surg Am*. 2017;99(18):e98.

MS-3: 3D Rendering

- Indications for musculoskeletal 3-D image post-processing for preoperative planning when conventional imaging is insufficient for:
 - ◆ Complex fractures/dislocations (comminuted or displaced) of any joint.
 - ◆ Spine fractures, pelvic/acetabulum fractures, intra-articular fractures.
 - ◆ Preoperative planning for other complex surgical cases.
- The code assignment for 3-D rendering depends upon whether the 3-D post-processing is performed on the scanner workstation (CPT® 76376) or on an independent workstation (CPT® 76377).
 - ◆ 2-D reconstruction (i.e. reformatting axial images into the coronal plane) is considered part of the tomography procedure, is not separately reportable, and does not meet the definition of 3-D rendering.
 - ◆ It is not appropriate to report 3-D rendering in conjunction with CTA and MRA because those procedure codes already include the post-processing.
 - ◆ In addition to the term “3-D,” the following terms may also be used to describe 3-D post-processing:
 - Maximum intensity projection (MIP)
 - Shaded surface rendering
 - Volume rendering
- The 3-D rendering codes require concurrent supervision of image post-processing 3-D manipulation of volumetric data set and image rendering. Certain health plan payors do not reimburse separately for 3-D rendering while others may have differing indication/limitation criteria. In these cases, individual plan coverage policies may take precedence over eviCore guidelines.

Reference

1. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® acute hand and wrist trauma. *Am Coll Radiol (ACR)*; Date of Origin:1995. Last Review: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.

MS-4: Avascular Necrosis (AVN)/Osteonecrosis

MS-4.1: AVN

15

MS-4.1: AVN

- Classification systems use a combination of plain radiographs, MRI, and clinical features to stage avascular necrosis. MRI of the area of concern without contrast can be performed when plain x-ray findings are negative or equivocal and clinical symptoms warrant further investigation for suspected avascular necrosis.
- Advanced imaging for AVN confirmed by plain x-ray is appropriate in the following situations:
 - ◆ Femoral head collapse:
 - MRI Hip without contrast (CPT® 73721) or CT Hip without contrast (CPT® 73700) for preoperative planning. See: **MS-24: Hip**.
 - ◆ Distal Femur:
 - MRI Knee without contrast (CPT® 73721) if needed for treatment planning. See: **MS-25: Knee**.
 - ◆ Talus:
 - MRI Ankle without contrast (CPT® 73721) if needed for treatment planning. See: **MS-26: Ankle**.
 - ◆ Tarsal navicular (Kohler Disease):
 - MRI Foot without contrast (CPT® 73718) if needed for treatment planning. See: **MS-27: Foot**.
 - ◆ Metatarsal head (Frieberg's Infraction):
 - MRI Foot without contrast (CPT® 73718) if needed for treatment planning. See: **MS-27: Foot**.
 - ◆ Humeral head:
 - For preoperative planning prior to shoulder replacement: CT Shoulder without contrast (CPT® 73200) and/or MRI Shoulder without contrast (CPT® 73221). See: **MS-19: Shoulder**.
 - ◆ Lunate (Kienbock's Disease)/Scaphoid (Preiser's Disease):
 - CT Wrist without contrast (CPT® 73200) or MRI Wrist without contrast (CPT® 73221). See: **MS-21: Wrist**.
- Patients with acute lymphoblastic leukemia and known or suspected osteonecrosis should be imaged according to guidelines in: **PEDONC-3.2: Acute Lymphoblastic Leukemia**
- Known or suspected osteonecrosis in long-term cancer survivors should be imaged according to guidelines in: **PEDONC-19.4: Osteonecrosis in Long Term Cancer Survivors**

References

1. Calder JD, Hine AL, Pearse MF, Revell PA. The relationship between osteonecrosis of the proximal femur identified by MRI and lesions proven by histological examination. *J Bone Joint Surg Br.* 2008;90(2):154-158.
2. Karantanas AH, Drakonaki EE. The role of MR imaging in avascular necrosis of the femoral head. *Semin Musculoskelet Radiol.* 2011;15(3):281-300. doi:10.1055/s-0031-1278427
3. Karim AR, Cherian JJ, Jauregui JJ, et al. Osteonecrosis of the knee: review. *Ann Transl Med.* 2015;3(1). doi:10.3978/j.issn.2305-5839.2014.11.13.
4. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic hip pain. *Am Coll Radiol (ACR)*; Revised:2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
5. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic wrist pain. *Am Coll Radiol (ACR)*; Revised:2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
6. Bennett DL, Nelson JW, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® nontraumatic knee pain. *Am Coll Radiol (ACR)*; Date of Origin:1995. Last Review:2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
7. Murphey MD, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® osteonecrosis of the hip. *Am Coll Radiol (ACR)*; Date of Origin:1995. Last Review: 2015. <https://acsearch.acr.org/docs/69420/Narrative/>.

MS-5: Fractures

MS-5.1: Acute	18
MS-5.2: Suspected Occult/Stress/Insufficiency Fracture/Stress Reaction and Shin Splints	18
MS-5.3: Other Indications	19

MS-5.1: Acute

- CT or MRI without contrast if **ANY** of the following:
 - ◆ Complex (comminuted or displaced) fracture with or without dislocation on plain x-ray.
 - CT is preferred unless it is associated with neoplastic disease when MRI without/with contrast is preferred unless MRI contraindicated.
 - ◆ Patient presents initially to the requesting provider with a documented history of an acute traumatic event at least two weeks prior with a negative plain x-ray at the time of this face-to-face encounter and a clinical suspicion for an occult/stress/insufficiency fracture see: **MS-5.2: Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints.**
- MRI without contrast, MRI with contrast (arthrogram), or CT with contrast (arthrogram) of the area of interest if:
 - ◆ Plain x-rays are negative and an osteochondral fracture is suspected, OR
 - ◆ Plain x-ray and clinical exam suggest an unstable osteochondral injury. See also: **MS-13.1: Chondral/ Osteochondral Lesions, Including Osteochondritis Dissecans and Fractures**

MS-5.2: Suspected Occult/Stress/Insufficiency Fracture/Stress Reaction and Shin Splints

- MRI without contrast can be performed for suspected hip/femoral neck, tibia, pelvis/sacrum, tarsal navicular, proximal fifth metatarsal, or scaphoid occult/stress/insufficiency fractures, and suspected atypical femoral shaft fractures related to bisphosphonate use if the initial evaluation of history, physical exam and plain x-ray fails to establish a definitive diagnosis.
 - ◆ CT without contrast can be performed as an alternative to MRI for suspected occult/insufficiency fractures of the pelvis/hip and suspected atypical femoral shaft fractures related to bisphosphonate see: **MS-23: Pelvis** and **MS-24: Hip**, and suspected occult fractures of the scaphoid see: **MS-21: Wrist.**
 - ◆ For suspected fractures, when MRI cannot be performed, any one of the following is indicated:
 - Tc-99m Bone scan whole body (CPT® 78306) with SPECT of the area of interest (CPT® 78803) OR
 - Bone scan (CPT® 78315, 78300) OR
 - SPECT/CT (CPT® 78830)
 - See: **MS-28: Nuclear Medicine.**
 - ◆ Tc-99m Bone scan Foot (CPT® 78315) is indicated for suspected occult or stress fractures of the tarsal navicular if MRI cannot be performed see: **MS-27: Foot.**
- MRI or CT without contrast can be performed for all other suspected occult/stress/insufficiency fractures with either of the following:
 - ◆ Repeat plain x-rays remain non-diagnostic for fracture after a minimum of 10 days of provider-directed conservative treatment, or
 - ◆ Initial plain x-rays obtained a minimum of 14 days after the onset of symptoms are non-diagnostic for fracture

- MRI of the lower leg without contrast (CPT® 73718) for suspected shin splints when **BOTH** of the following are met:
 - ◆ Initial plain x-ray
 - ◆ Failure of a 6-week trial of provider-directed conservative treatment.
- For stress reaction, advanced imaging is not medically necessary for surveillance or “return to play” decisions regarding a stress reaction identified on an initial imaging study.
- MRI without contrast of the area of interest for stress fracture follow-up imaging for “return to play” evaluation at least 3 months after the initial imaging study for stress fracture.
- For periprosthetic fractures related to joint replacement see: **MS-16.1: Post-Operative Joint Replacement Surgery**, **MS-19: Shoulder**, **MS-20: Elbow**, **MS-24: Hip**, **MS-25: Knee**, and **MS-26: Ankle**.

MS-5.3: Other Indications

- CT or MRI without contrast is appropriate after recent (within 30 days) plain x-ray if **ONE** of the following is present:
 - ◆ Concern for delayed union or non-union of fracture, osteotomy, or joint fusions.
 - ◆ As part of preoperative evaluation for a planned surgery of a complex fracture with or without dislocation.

References

1. Bencardino JT, Stone TJ, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
2. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
3. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hand and Wrist Trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
4. Luchs JS, Flug JA, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Ankle Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
5. Ward RJ, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hip Pain-Suspected Fracture. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Last Review: 2018. <https://acsearch.acr.org/docs/3082587/Narrative/>.
6. Mosher TJ, Kransdorf MJ, Adler R, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Ankle. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Last Review: 2020. <https://acsearch.acr.org/docs/69436/Narrative/>.
7. Hayes CW, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Elbow Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2015. <https://acsearch.acr.org/docs/69423/Narrative/>.
8. Wise JN, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
9. Greene WB. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:568-570.
10. Galbraith RM, Lavalley ME. Medial tibial stress syndrome: conservative treatment options. *Curr Rev Musculoskelet Med*. 2009;2:127-133. doi:10.1007/s12178-009-9055-6.
11. Boks SS, Vroegindeweij D, Kroes BW, et al. MRI follow-up of posttraumatic bone bruises of the knee in general practice. *AJR Am J Roentgenol*. 2007;189 556-562. doi:10.2214/AJR.07.2276.
12. Kaeding CC, Yu JR, Wright R, et al. Management and return to play of stress fractures. *Clin J Sport Med*. 2005;15:442-7.
13. Sormaala MJ, Niva MH, Kiuru MJ, et al. Stress injuries of the calcaneus detected with magnetic resonance imaging in military recruits. *J Bone Joint Surg Am*. 2006;88:2237-42. doi:10.2106/JBJS.E.01447.
14. Shin AY, Morin WD, Gorman JD, et al. The superiority of magnetic resonance imaging in differentiating the cause of hip pain in endurance athletes. *Am J Sports Med*. 1996;24:168-76. doi:10.1177/036354659602400209.
15. Slocum KA, Gorman JD, Puckett ML, et al. Resolution of abnormal MR signal intensity in patients with stress fractures of the femoral neck. *AJR Am J Roentgenol*. 1997;168:1295-9. doi:10.2214/ajr.168.5.9129429.
16. Fredericson M, Bergman AG, Hoffman KL, et al. Tibial stress reaction in runners. Correlation of clinical symptoms and scintigraphy with a new magnetic resonance imaging grading system. *Am J Sports Med*. 1995;23(4):472-81. doi:10.1177/036354659502300418.

MS-6: Foreign Body

MS-6.1: Foreign Body – General

22

MS-6.1: Foreign Body – General

- Ultrasound (CPT® 76881 or 76882) or CT without contrast or MRI without and with contrast or MRI without contrast of the area of interest can be approved after plain x-rays rule out the presence of radiopaque foreign bodies.
 - ◆ Ultrasound (CPT® 76881 or 76882) is the preferred imaging modality for radiolucent (non-radiopaque) foreign bodies (e.g. wood, plastic).
 - ◆ CT without contrast is recommended when plain x-rays are negative and a radiopaque foreign body is still suspected, as CT is favored over MRI for the identification of foreign bodies
 - ◆ MRI without and with contrast is an alternative to US and CT for assessing the extent of infection associated with a suspected foreign body

References

1. Bancroft LW, Kransdorf MJ, Adler R, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Foot. *Am Coll Radiol (ACR)*; Date of Origin: 2010. Last Review: 2019. <https://acsearch.acr.org/docs/70546/Narrative/>.
2. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/%203094201/Narrative>.
3. Chan C, Salam GA. Splinter removal. *Am Fam Physician*. 2003;67(12):2557-2562.
4. Peterson JJ, Bancroft LW, Kransdorf MJ. Wooden foreign bodies: imaging appearance. (*AJR Am J Roentgenol*. 2002;178(3):557-562. doi:10.2214/ajr.178.3.1780557.
5. Jarraya M, Hayashi D, de Villiers RV, et al. Multimodality imaging of foreign bodies of the musculoskeletal system. (*AJR Am J Roentgenol*. 2014;203(1):W92-102. doi:10.2214/AJR.13.11743.

MS-7: Ganglion Cysts

MS-7.1: Ganglion Cysts – General

24

MS-7.1: Ganglion Cysts – General

- Plain x-ray is the initial imaging study for ganglion cysts.
- MRI without contrast or MRI without and with contrast or US (CPT® 76881 or 76882) is appropriate for surgical planning.
- Advanced imaging is not indicated for ganglions that can be diagnosed by history and physical examination.

References

1. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging ACR Appropriateness Criteria® Chronic wrist pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
2. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Soft-Tissue Masses. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Revised: 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
3. Freire V, Guerini H, Campagna R, et al. Imaging of hand and wrist cysts: a clinical approach. (*AJR Am J R Roentgenol*. 2012;199(5):W618-W628. doi:10.2214/AJR.11.8087.
4. Vo P, Wright T, Hayden F, Dell P, et al. Evaluating dorsal wrist pain: MRI diagnosis of occult dorsal wrist ganglion. *J Hand Surg Am*. 1995;20(4):667-670. doi:10.1016/S0363-5023(05)80288-6.
5. Teefey SA, Dahiya N, Middleton WD, et al. Ganglia of the hand and wrist: a sonographic analysis. *AJR Am J Roentgenol*. 2008;191(3):716-720. doi:10.2214/AJR.07.3438.

MS-8: Gout/Calcium Pyrophosphate Deposition Disease (CPPD)/ Pseudogout/Chondrocalcinosis

MS-8.1: Gout – General	26
MS-8.2: CPPD (Pseudogout/Chondrocalcinosis) – General	26

MS-8.1: Gout – General

- CT without contrast, MRI without contrast, or MRI without and with contrast of the area of interest is indicated when **BOTH** of the following are met:
 - ◆ Initial plain x-ray has been performed to rule out other potential disease processes
 - ◆ Infection or neoplasm is in the differential diagnosis for soft-tissue tophi.

Practice Notes

- Early stages of gout can be diagnosed clinically since radiographic findings are not present early in the disease course.

MS-8.2: CPPD (Pseudogout/Chondrocalcinosis) – General

- Calcium pyrophosphate deposition disease (CPPD), also called pseudogout, can often be diagnosed from plain x-rays; advanced diagnostic imaging is generally not medically necessary.

References

1. Hsu CY, Shih TT, Huang KM, et al. Tophaceous gout of the spine: MR imaging features. *Clin Radiol*. 2002;57(10):919.
2. Schumacher HR Jr, Becker MA, Edwards NL, et al. Magnetic resonance imaging in the quantitative assessment of gouty tophi. *Int J Clin Pract*. 2006;60(4):408. doi:10.1111/j.1368-5031.2006.00853.x.
3. McQueen FM, Doyle A, Reeves Q, Gao A. Bone erosions in patients with chronic gouty arthropathy are associated with tophi but not bone oedema or synovitis: new insights from a 3 T MRI study. *Rheumatology*. 2014;53(1):95-103. doi:10.1093/rheumatology/ket329.
4. Dore RK. Gout: What primary care physicians want to know. *J Clin Rheumatol*. 2008;14(5S):S47-S54. doi:10.1097/RHU.0b013e3181896c35.
5. Eggebeen AT. Gout: an update. *Am Fam Physician*. 2007;76(6):801-808.
6. Burns C, Wortmann RL. Gout. In: Imboden JB, Hellmann DB, Stone JH, eds. *CURRENT Diagnosis & Treatment: Rheumatology*. 3rd edition. New York: McGraw-Hill; 2013:332-338.
7. Jacobson JA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic extremity joint pain-suspected inflammatory arthritis. *Am Coll Radiol (ACR)*; 2017;14(5):S81-S89. [http://www.jacr.org/article/S1546-1440\(17\)30183-7/fulltext](http://www.jacr.org/article/S1546-1440(17)30183-7/fulltext).

MS-9: Infection/Osteomyelitis

MS-9.1: Infection – General	28
MS-9.2: Septic Joint	28

MS-9.1: Infection – General

- MRI without and with contrast after plain x-ray(s) and:
 - ◆ Plain x-ray(s) are negative or do not suggest alternative diagnoses such as neuropathic arthropathy or fracture, and soft tissue or bone infection (osteomyelitis) is suspected; *or*
 - ◆ Plain x-ray(s) are positive for osteomyelitis, and the extent of infection into the soft tissues and any skip lesions require evaluation.
- CT without and with contrast can replace an MRI:
 - ◆ To assess the extent of bony destruction from osteomyelitis; CT can guide treatment decisions.
 - ◆ For preoperative planning
 - ◆ If MRI is contraindicated
- Patients with suspected spinal infections
 - ◆ See: **SP-1.2: Red Flag Indications** for advanced imaging guidelines
- Patients with diabetic foot infections after plain x-ray(s)
 - ◆ See: **MS-27: Foot** for advanced imaging guidelines

MS-9.2: Septic Joint

- MRI of the joint, without and with contrast is appropriate when standard or image-guided arthrocentesis is contraindicated, unsuccessful, or non-diagnostic, and the clinical documentation satisfies ALL of the following criteria:
 - ◆ History and physical examination findings [One of the following]:
 - Development of an acutely hot and swollen joint (< 2 weeks)
 - Decreased range of motion due to pain
 - Documented fever
 - ◆ Laboratory tests [One of the following]:
 - Leukocytosis
 - Elevated ESR or C-reactive protein
 - Analysis of the joint fluid is non-diagnostic
 - ◆ Plain x-ray of the joint
- MRI without and with contrast is appropriate after plain x-rays if the arthrocentesis is diagnostic and if there is a confirmed septic joint, to evaluate the extent of infection into the soft tissues and any skip lesions that would require evaluation.
- CT with contrast can replace MRI without and with contrast if MRI is contraindicated.

Practice Notes

- Analysis of joint fluid is most often sufficient to diagnose a septic joint.

References

1. Coakley G, Mathews C, Field M, et al. BSR & BHP, BOA, RCGP and BSAC guidelines for management of the hot swollen joints in adults. *Rheumatology*. 2006;45(8):1039-1041. doi:10.1093/rheumatology/kel163a.
2. Karchevsky M, Schweitzer ME, Morrison WB, et al. MRI findings of septic arthritis and associated osteomyelitis in adults. (*AJR*) *Am J Roentgenol*. 2004;182(1):119-122. doi:10.2214/ajr.182.1.1820119.
3. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:918.
4. Staheli LT. Septic arthritis. In: Staheli LT, ed. *Fundamentals of Pediatric Orthopedics*. 4th edition. Philadelphia, PA: Lippincott Williams & Wilkins; 2008:110-111.
5. Kransdorf MJ, Weissman BN, Appel M, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected osteomyelitis of the foot in patients with diabetes mellitus. *Am Coll Radiol (ACR)*. 2012. <https://acsearch.acr.org/docs/69340/Narrative/>.
6. Beaman FD, von Herrmann PF, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected osteomyelitis, septic arthritis, or soft tissue infection (excluding spine and diabetic foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/3094201/Narrative/>.
7. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic wrist pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
8. Small KM, Adler RS, Shah SH, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Atraumatic. *Am Coll Radiol (ACR)*; New 2018. <https://acsearch.acr.org/docs/3101482/Narrative/>.
9. Amini B, Beckmann NM, Beaman FD, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Traumatic. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69433/Narrative/>.
10. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic hip pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
11. Reinus WR. *Clinician's Guide to Diagnostic Imaging*. 2014. Springer-Verlag New York.
12. Visconti AJ, Biddle J, Solomon M. Follow-up imaging for vertebral osteomyelitis a teachable moment. *JAMA Intern Med*. 2014;174(2):184. doi:10.1001/jamainternmed.2013.12742.
13. Fabiano V, Franchino G, Napolitano M, et al. Utility of magnetic resonance imaging in the follow-up of children affected by acute osteomyelitis. *Curr Pediatr Res*. 2017;21(2):354-358.
14. Patel ND, Broderick DF, Burns J, et al. Expert Panel on Neurologic Imaging. ACR Appropriateness Criteria®: Low back pain. *Am Coll Radiol (ACR)*. 2015; <https://acsearch.acr.org/docs/69483/Narrative/>.

MS-10: Soft Tissue Mass or Lesion of Bone

MS-10.1: Soft Tissue Mass	31
MS-10.2: Lesion of Bone	31

MS-10.1: Soft Tissue Mass

- History and physical exam should include documentation of: location, size, duration, growing or stable, solid/cystic, fixed/not fixed to the bone, discrete or ill-defined, and an association with pain.
- MRI without and with contrast or without contrast or US of the area of interest (CPT® 76881 or 76882) is appropriate when ANY of the following are met after plain x-ray:
 - ◆ Soft tissue mass(es)
 - ◆ Surgical planning
 - ◆ Known or suspected soft tissue mass in a patient with a cancer predisposition syndrome if a recent ultrasound is inconclusive. **Plain x-ray is not required for these patients.** See: **PEDONC-2: Screening Imaging in Cancer Predisposition Syndromes**
- CT with contrast or CT without and with contrast is appropriate when MRI is contraindicated or after a metal limiting MRI evaluation.
- Advanced imaging is not indicated for:
 - ◆ Subcutaneous lipoma with no surgery planned
 - ◆ Ganglia, see: **MS-7: Ganglion Cysts**
 - ◆ Sebaceous cyst

Practice Notes

- Plain x-rays can determine if an advanced imaging procedure is indicated, and if so, which modality is most appropriate. If non-diagnostic, these initial plain x-rays can provide complementary information if advanced imaging is indicated.

MS-10.2: Lesion of Bone

- History and physical exam should include documentation of: location, size, duration, growing or stable, discrete or poorly defined, and an association with pain.
- Complete radiograph of the entire bone containing the lesion of bone is required prior to consideration of advanced imaging. Many benign bone tumors have a characteristic appearance on plain x-ray and advanced imaging is not necessary.
- MRI without and with contrast, MRI without contrast, or CT without contrast may be indicated if ONE of the following applies:
 - ◆ Diagnosis uncertain based on plain x-ray appearance.
 - ◆ Imaging requested for preoperative planning.
- MRI without and with contrast or without contrast is appropriate when plain x-ray reveals an osteochondroma with clinical concern of malignant transformation.

➤ For Paget's Disease:

- ◆ Bone scan (See: **MS-28: Nuclear Medicine**) OR
- ◆ MRI (contrast as requested) can be considered if the diagnosis (based on plain x-rays and laboratory studies) is in doubt.
- ◆ MRI (contrast as requested) can be considered if malignant degeneration, which occurs in up to 10% of cases, is suspected.

References

1. Peterson JJ, Beaman FD, Fox MG, et al. ACR Practice Guideline. ACR-SSR Practice Guideline for the Performance and Interpretation of Magnetic Resonance Imaging (MRI) of bone and soft tissue tumors. *Am Coll Radiol*. Revised 2015. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-SoftTissue-Tumors.pdf>.
2. Zoga AC, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Soft-tissue masses. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
3. Hayes CW, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic elbow pain. *Am Coll Radiol (ACR)*; Revised 2015. <https://acsearch.acr.org/docs/69423/Narrative/>.
4. Musculoskeletal Tumor Society: Systematic Literature Review on the Use of Imaging Prior to Referral to a Musculoskeletal Oncologist. Rosemont, IL, Musculoskeletal Tumor Society, February 2018.
5. Schneider D, Hofmann MR, and Peterson JA. Diagnosis and treatment of Paget's Disease of Bone. *Am Fam Physician*. 2002;65(10):2069-2072. <https://www.aafp.org/afp/2002/0515/p2069.html>.
6. Theodorou DJ, Theodorou SJ, and Kakitsubata Y. Imaging of Paget Disease of bone and its musculoskeletal complications: review. (*AJR*) *Am J Roentgenol*. 2012;196(6):S64-S75.
7. Sinha S and Peach AH. Diagnosis and management of soft tissue sarcoma. *BMJ*. 2010;341:c7170. doi:10.1136/bmj.c7170.
8. Wu JS, Hochman MG. Soft-tissue tumors and tumorlike lesions: a systematic imaging approach. *Radiology*. 2009;253(2):297-316. doi:10.1148/radiol.2532081199.

MS-11: Muscle/Tendon Unit Injuries/Diseases

MS-11.1: Muscle/Tendon Unit Injuries/Diseases	34
MS-11.2: Acute Compartment Syndrome	34
MS-11.3: Chronic Exertional Compartment Syndrome	34

MS-11.1: Muscle/Tendon Unit Injuries/Diseases

- Plain x-ray is the initial imaging study for muscle/tendon unit injuries.
- MRI without contrast or US (CPT® 76881 or 76882) is supported for **EITHER** of the following:
 - ◆ Suspected partial tendon rupture of a specific (named) tendon
 - ◆ Complete tendon ruptures for preoperative planning (for example, Achilles tendon rupture, posterior tibial tendon rupture, humeral insertion of the pectoralis major rupture, proximal and distal biceps tendon rupture, patellar ligament/tendon rupture, proximal/distal hamstring tendon rupture).
- MRI is not medically necessary for muscle belly strains/muscle tears.
- See: **MS-19: Shoulder** for clinical suspicion of a partial or complete rotator cuff tear.
- See: **PN-6.2: Inflammatory Muscle Diseases** and **PEDMS-10.3: Inflammatory Muscle Diseases**

MS-11.2: Acute Compartment Syndrome

- Advanced imaging is not indicated. Diagnosis is made clinically and by direct measurement of compartment pressure and is a surgical emergency.

Practice Notes

- Noninvasive methods of measuring compartment pressures and diagnosing acute compartment syndrome are under study, but are currently experimental, investigational, and unproven.

MS-11.3: Chronic Exertional Compartment Syndrome

- Advanced imaging should only be considered when ruling out other potential causes of extremity pain following a plain x-ray and conservative treatment as indicated.

Practice Notes

- Direct measurement of compartment pressure remains the diagnostic standard. Noninvasive methods of measuring compartment pressures and diagnosing chronic exertional compartment syndrome are under study, but are currently experimental, investigational, and unproven.

References

1. Luchs JS, Flug JA, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® chronic ankle pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
2. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:452.
3. Kayser R, Mahlfeld K, Heyde CE. Partial rupture of the proximal Achilles tendon: a differential diagnostic problem in ultrasound imaging. *Br J Sports Med*. 2005;39:838-842. doi:10.1136/bjsm.2005.018416.
4. Rominger MB, Lukosch CJ, Bachmann GF. MR imaging of compartment syndrome of the lower leg: a case control study. *Eur Radiol*. 2004;14:1432-1439. doi:10.1007/s00330-004-2305-5.
5. McDonald S, Bearcroft P. Compartment syndromes. *Semin Musculoskelet Radiol*. 2010;14(2):236-244. doi:10.1055/s-0030-1253164.
6. Ringler MD, Litwiller DV, Felmlee JP, et al. MRI accurately detects chronic exertional compartment syndrome: a validation study. *Skeletal Radiology*. 2013;42:385-392. doi:10.1007/s00256-012-1487-1.
7. van den Brand JG, Nelson T, Verleisdonk EJ, van der Werken C. The diagnostic value of intracompartmental pressure measurement, magnetic resonance imaging, and near-infrared spectroscopy in chronic exertional compartment syndrome: a prospective study in 50 patients. *Am J Sports Med*. 2005;33:699-704. doi:10.1177/0363546504270565.
8. Heer ST, Callander JW, Kraeutler MJ, Mei-Dan O, Mulcahey MK. Hamstring Injuries. *The Journal of Bone and Joint Surgery*. 2019;101(9):843-853. doi:10.2106/jbjs.18.00261.

MS-12: Osteoarthritis

MS-12.1: Osteoarthritis

37

MS-12.1: Osteoarthritis

- Plain x-ray is the initial imaging study for osteoarthritis.
- CT without contrast is appropriate for treatment planning when congenital or significant atypical post-traumatic arthritic deformities are present in the shoulder, elbow, wrist, hip, knee, or ankle that would require further evaluation of the clinical significance of the deformity already identified on plain x-rays.
 - ◆ CT shoulder without contrast (CPT® 73200) and/or MRI shoulder without contrast (CPT® 73221) are considered medically necessary for preoperative planning prior to shoulder replacement
- Preoperative non-contrast CT/MRI requests (for either a diagnostic or unlisted CPT® code) of the shoulder, elbow, wrist, hip, knee, or ankle to be utilized as part of treatment planning for customized-to-patient joint replacement surgery or as an integral part of surgical planning using intraoperative navigation for joint replacement surgery (e.g. MAKOplasty) are considered medically necessary once the joint replacement surgery has been approved or if the joint replacement surgery does not require prior authorization.
 - ◆ Requests for preoperative imaging are considered not medically necessary if the surgery, customized-to-patient implant, and/or computer assisted navigation (e.g. MAKOplasty) has been deemed experimental, investigational, or unproven by the health plan
 - ◆ Benefits, coverage policies, and eligibility issues pertaining to each Health Plan may take precedence over eviCore's guidelines. Providers are urged to obtain written instructions and requirements directly from each payor. See: **Preface-4.3: Unlisted Procedures/Therapy Treatment Planning**
- MRI arthrogram or CT arthrogram is appropriate when joint sparing/salvage reconstructive surgery is planned for the following:
 - ◆ Suspected concomitant rotator cuff tear of the shoulder - See: **MS-19: Shoulder**
 - ◆ Suspected concomitant labral tear of the shoulder - See: **MS-19: Shoulder**
 - ◆ Suspected concomitant labral tear of the hip - See: **MS-24: Hip**
 - ◆ Suspected concomitant internal derangement of the knee - See: **MS-25: Knee**

Note:

- Refer to the Anatomic Area Tables **MS-19: Shoulder**, **MS-20: Elbow**, **MS-21: Wrist**, **MS-24: Hip**, **MS-25: Knee**, and **MS-26: Ankle** for the clinical imaging criteria regarding preoperative joint replacement surgery for each anatomic area.
- MRI knee without contrast (CPT® 73721) is appropriate in a patient with osteoarthritis for clinical suspicion of a symptomatic degenerative meniscus tear following plain x-rays and conservative treatment. See: **MS-25: Knee**

Practice Notes

Plain x-rays are performed initially and will reveal characteristic joint space narrowing, osteophyte formation, cyst formation, and subchondral sclerosis.

References

1. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
2. Bennett DL, Nelson JW, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Nontraumatic Knee Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
3. Manek NJ, Lane NE. Osteoarthritis: Current concepts in diagnosis and management. *Am Fam Physician* 2000 March;61(6):1795-1804. <https://www.aafp.org/afp/2000/0315/p1795.html>.
4. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:84.
5. Quatman CE, Hettrich CM, Schmitt LC, et. al. The Clinical Utility and Diagnostic Performance of MRI for Identification of Early and Advanced Knee Osteoarthritis: A Systematic Review. *Am J Sports Med*. 2011;39(7):1557–1568. doi:10.1177/0363546511407612.
6. Braun HJ, Gold GE. Diagnosis of osteoarthritis: imaging. *Bone*. 2012;51(2):278–288. doi:10.1016/j.bone.2011.11.019.

MS-13: Chondral/Osteochondral Lesions**MS-13.1: Chondral/Osteochondral Lesions, Including
Osteochondritis Dissecans and Fractures****40**

MS-13.1: Chondral/Osteochondral Lesions, Including Osteochondritis Dissecans and Fractures

- MRI without contrast, MRI with contrast (arthrogram), or CT with contrast (arthrogram) of the area of interest is indicated when **EITHER** of the following are met:
 - ◆ Plain x-rays are negative and an osteochondral fracture is still suspected
 - ◆ Plain x-ray and clinical exam suggest an unstable osteochondral injury
- See: **MS-26: Ankle** for suspected osteochondral injury of the ankle
- If plain x-rays show a non-displaced osteochondral fragment, follow-up imaging should be with plain x-rays. Advanced imaging is not necessary.
- MRI without contrast or CT without contrast is indicated when healing (including post-operative fixation) cannot be adequately assessed on follow-up plain x-rays.

References

1. Bridges MD, Berland LL, Cernigliaro JG, et al. ACR Practice Guideline. ACR-SSR Practice Guideline for the Performance and Interpretation of Magnetic Resonance Imaging (MRI). *Am Coll Radiol (ACR)*. 2017. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-perf-interpret.pdf?la=en>.
2. Bennett DL, Nelson JW, Weissman BN, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Nontraumatic Knee Pain. *Am Coll Radiol (ACR)*; 2012. Last Review: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
3. Rubin DA, Anderson MW, Hastreiter DM, et al. ACR Practice Guideline. ACR-SSR Practice Guideline for the Performance and Interpretation of Magnetic Resonance Imaging (MRI) of the elbow. *Am Coll Radiol (ACR)*. Revised 2016. <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/mr-elbow.pdf?la=en>.

MS-14: Osteoporosis

- Plain x-ray is not required for MS-14: Osteoporosis.
- Quantitative CT (CPT® 77078) can be approved for screening when DXA scanner is unavailable or known to be inaccurate for ANY of the following populations:
 - ◆ Women age ≥65 years
 - ◆ Men age >70 years
 - ◆ Women age <65 years who have additional risk factors for osteoporosis based on medical history and other findings:
 - Estrogen deficiency
 - A history of maternal hip fracture that occurred after age 50 years
 - Low body mass (<127 lb. or 57.6 kg)
 - History of amenorrhea (>1 year before age 42 years)
 - ◆ Women age <65 years or men age <70 years who have additional risk factors:
 - Current use of cigarettes
 - Loss of height, thoracic kyphosis
 - ◆ Individuals of any age with bone mass osteopenia or fragility fractures on imaging studies such as radiographs, CT, or MRI
 - ◆ Individuals age 50 years and older who develop a wrist, hip, spine, or proximal humerus fracture with minimal or no trauma, excluding pathologic fractures
 - ◆ Individuals of any age who develop 1 or more insufficiency fractures
 - ◆ Premenopausal females or males age 20 to 50 years with risk factors:
 - Individuals with medical conditions that could alter bone mineral density
 - Chronic renal failure
 - Rheumatoid arthritis and other inflammatory arthritides
 - Eating disorders, including anorexia nervosa and bulimia
 - Organ transplantation
 - Prolonged immobilization
 - Conditions associated with secondary osteoporosis, such as gastrointestinal malabsorption or malnutrition, sprue, osteomalacia, vitamin D deficiency, endometriosis, acromegaly, chronic alcoholism or established cirrhosis, and multiple myeloma
 - Individuals who have had gastric bypass for obesity
 - Individuals with an endocrine disorder known to adversely affect bone mineral density (e.g., hyperparathyroidism, hyperthyroidism, or Cushing syndrome)
 - Individuals receiving (or expected to receive) glucocorticoid therapy for >3 months
 - Hypogonadal men older than 18 years and men with surgically or chemotherapeutically induced castration
 - Individuals beginning or receiving long-term therapy with medications known to adversely affect BMD (e.g. anticonvulsant drugs, androgen deprivation therapy, aromatase inhibitor therapy, or chronic heparin)

Note: Repeat screening quantitative computed tomography (QCT) can be approved no sooner than every two years.

- Quantitative CT scan (CPT® 77078) can be approved for non-screening/monitoring when DXA scanner is unavailable or known to be inaccurate for ANY of the following circumstances:
- ◆ Follow-up in cases where QCT was the original study
 - ◆ Multiple healed vertebral compression fractures
 - ◆ Significant scoliosis
 - ◆ Advanced arthritis of the spine due to increased cortical sclerosis often with large marginal osteophytes. Obese patient over the weight limit of the dual-energy x-ray absorptiometry (DXA) exam table
 - ◆ Severely obese patients (BMI >35kg/m²)
 - ◆ Extremes in body height (i.e. very large and very small patients)
 - ◆ Patients with extensive degenerative disease of the spine
 - ◆ A clinical scenario that requires sensitivity to small changes in trabecular bone density (parathyroid hormone and glucocorticoid treatment monitoring).

Note: Repeat non-screening/monitoring QCT can be approved no earlier than one year following a change in treatment regimen, and only when the results will directly impact a treatment decision.

References

1. American Association of Clinical Endocrinologists (AACE) Menopause Guidelines Revision Task Force. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract.* 2016;22(Suppl 4):1-42.
<https://www.aace.com/files/postmenopausal-guidelines.pdf>.
2. Coleman F, de Buer SJ, LeBoff MS, et al. National Osteoporosis Foundation (NOF). Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int.* 2014;25(10):2359–2381. doi:10.1007/s00198-014-2794-2.
3. U.S. Preventive Services Task Force (USPSTF). Final Recommendation Statement Osteoporosis: Screening. January 2011.
4. Ward RJ, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Osteoporosis and Bone Mineral Density. *Am Coll Radiol (ACR)*; Revised 2016.
<https://acsearch.acr.org/docs/69358/Narrative/>.

MS-15: Rheumatoid Arthritis (RA) and Inflammatory Arthritis

MS-15.1: Rheumatoid Arthritis (RA) and Inflammatory Arthritis	44
MS-15.2: Pigmented Villonodular Synovitis (PVNS)	44

MS-15.1: Rheumatoid Arthritis (RA) and Inflammatory Arthritis

- Plain x-ray, physical exam and appropriate laboratory studies* are required prior to advanced imaging.
- MRI without contrast or MRI without and with contrast or US (CPT® 76881 or 76882) is appropriate for the most symptomatic joint, or of the dominant hand or wrist, in ALL of the following situations:
 - ◆ When diagnosis is uncertain prior to initiation of drug therapy.
 - ◆ To study the effects of treatment with disease modifying anti-rheumatic drug (DMARD) therapy.
 - ◆ To identify seronegative RA patients that might benefit from early DMARD therapy.
 - ◆ To determine change in treatment, such as:
 - Switching from standard DMARD therapy to tumor necrosis factor (TNF) therapy.
 - Changing to a different TNF drug therapy, then one MRI (contrast as requested) of a single joint can be performed.
 - Addition of other treatments, including joint injections
- MRI or US should NOT be considered for routine follow-up of treatment.

Practice Notes

- *Examples of appropriate laboratory studies may include: Lyme titers, rheumatoid factor (RF), anti-cyclic citrullinated peptide (anti-CCP), sedimentation rate (ESR), C-reactive protein (CRP), and antinuclear antibody (ANA)], joint fluid analysis

MS-15.2: Pigmented Villonodular Synovitis (PVNS)

- MRI of the affected joint without contrast, or CT of the affected joint with contrast (arthrogram) if MRI contraindicated is supported following plain x-rays.

References

1. Rubin DA, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Wrist Pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
2. Luchs JS, Flug JA, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Ankle Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
3. Hayes CW, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Elbow Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2015. <https://acsearch.acr.org/docs/69423/Narrative/>.
4. Jacobson JA, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Extremity Joint Pain-Suspected Inflammatory Arthritis. *Am Coll Radiol (ACR)*; New: 2016. <https://acsearch.acr.org/docs/3097211/Narrative/>.
5. Wise JN, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
6. Mintz DN, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
7. Boutry N, Morel M, Flipo RM, et al. Early rheumatoid arthritis: a review of MRI and sonographic findings. *AJR Am J Roentgenol*. 2007;189:1502-1509. doi:10.2214/AJR.07.2548.
8. Murphey MD, Rhee JH, Lewis RB, et al. Pigmented villonodular synovitis: radiologic-pathologic correlation. *Radiographics*. 2008;28:1493-1518. doi:10.1148/rg.285085134.
9. Conaghan P, Edmonds J, Emery P, et al. Magnetic resonance imaging in rheumatoid arthritis: summary of OMERACT activities, current status, and plans. *Journal of Rheumatology*. 2001;28(5):1158-1161. <http://www.jrheum.org/content/28/5/1158.long>.
10. Ostergaard M, McQueen FM, Bird P, et al. Magnetic resonance imaging in rheumatoid arthritis--advances and research priorities. *Journal of Rheumatology*. 2005;32(12):2462-2464. <http://www.jrheum.org/content/32/12/2462.long>.
11. McQueen FM. The use of MRI in early RA. *Rheumatology*. 2008;47(11):1597-1599. doi:10.1093/rheumatology/ken332.
12. Gossec L, Fautrel B, Pham T, et al. Structural evaluation in the management of patients with rheumatoid arthritis: development of recommendations for clinical practice based on published evidence and expert opinion. *Joint Bone Spine*. 2005;72(3):229-234. doi:10.1016/j.jbspin.2004.10.011.
13. Cohen SB, Potter H, Deodhar A, et al. Extremity magnetic resonance imaging in rheumatoid arthritis: updated literature review. *Arthritis Care & Research*. 2011;63(5):660-665. doi:10.1002/acr.20413.
14. Singh JA, Furst DE, Bharat A, et al. 2012 update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care & Research*. 2012;64(5):625-639. doi:10.1002/acr.21641.
15. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis & Rheumatism (Arthritis Care & Research)*. 2008;59:762-784. doi:10.1002/art.23721.

MS-16: Post-Operative Joint Replacement Surgery

MS-16.1: Post-Operative Joint Replacement Surgery – General 47

MS-16.1: Post-Operative Joint Replacement Surgery – General

- CT without contrast or bone scan (CPT® 78315) or Distribution of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831)* or hybrid SPECT/CT (CPT® 78830)* may be indicated for the evaluation of suspected aseptic loosening of orthopaedic joint replacements when recent plain x-ray is nondiagnostic.
 - ◆ CT shoulder without contrast (CPT® 73200) can be performed as additional imaging following plain x-rays regardless of plain x-ray findings. See: **MS-19: Shoulder**
- CT without contrast is appropriate with a high suspicion for a periprosthetic fracture and a negative plain x-ray.
 - ◆ CT shoulder without contrast (CPT® 73200) can be performed as additional imaging following plain x-rays regardless of plain x-ray findings. See: **MS-19: Shoulder**
- Joint aspiration is the initial evaluation after plain x-ray for a painful joint replacement when periprosthetic infection is suspected.
 - ◆ For suspected infection with negative or inconclusive joint aspiration culture see: **MS-28: Nuclear Medicine**
- MRI hip without contrast (CPT® 73721) or ultrasound (CPT® 76881 or 76882) are both appropriate for **EITHER** of the following:
 - ◆ Diagnosis of ALVAL (aseptic lymphocytic-dominated vasculitis-associated lesion) pseudotumors surrounding metal-on-metal (MoM) hip prostheses. One of these two imaging modalities can be approved but not both. See: **MS-10.1: Soft Tissue Mass or Lesion of Bone**
 - ◆ Metal-On-Metal (MoM) Hip Prostheses that are considered high risk for implant performance issues from THA cup-neck impingement and subsequent ALTR (adverse local tissue reaction) with Co and Cr ion levels greater than 10 ppb.
- CT hip without contrast (CPT® 73700) or MRI hip without contrast (CPT® 73721) is appropriate to evaluate suspected particle disease (aggressive granulomatous disease) of the hip when infection has been excluded.
- For specific joints post-operative from replacement surgery:
 - ◆ See: **MS-19: Shoulder**
 - ◆ See: **MS-20: Elbow**
 - ◆ See: **MS-24: Hip**
 - ◆ See: **MS-25: Knee**
 - ◆ See: **MS-26: Ankle**

Practice Notes

- Complications following joint replacement surgery include (not limited to) periprosthetic fracture, infection, aseptic loosening, failure of fixation/component malposition, and wear.
- *The usefulness of bone scan for the evaluation of suspected aseptic loosening of a shoulder replacement may be limited as bone remodeling–related increased uptake can be seen at the site of joint replacement for up to 1 year following surgery.

References

1. Mintz DN, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
2. Hochman MG, Melenevsky YV, Metter DF, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Total Knee Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69430/Narrative/>.
3. Gyftopoulos S, Rosenberg ZS, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Shoulder Arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/3097049/Narrative/>.
4. Toms AD, Davidson D, Masri BA, et al. Management of peri-prosthetic infection in total joint arthroplasty. *J Bone Joint Surg Br.* 2006;88(2):149-155. doi:10.1302/0301-620X.88B2.17058.
5. Love C, Marwin SE, Tomas MB, et al. Diagnosing infection in the failed joint replacement: A comparison of coincidence detection 18F-FDG and 111In-labeled leukocyte/99mTc-sulfur colloid marrow imaging. *J Nucl Med.* 2004;45(11):1864-1871.
6. Nawabi DH, Gold S, Lyman SL, et al. MRI predicts ALVAL and tissue damage in metal-on-metal hip arthroplasty. *Clin Orthop Relat Res.* 2014;472(2):471-481. doi:10.1007/s11999-013-2788-y.
7. Verberne SJ, Raijmakers PG, and Temmerman OP. The accuracy of imaging techniques in the assessment of periprosthetic hip infection: a systematic review and meta-analysis. *J Bone Joint Surg Am.* 2016;98(19):1638-45. doi:10.2106/JBJS.15.00898.

MS-17: Limb Length Discrepancy

MS-17.1: Limb Length Discrepancy

50

MS-17.1: Limb Length Discrepancy

- Either plain radiographic or “CT scanogram,” both reported with CPT® 77073, is appropriate to radiographically evaluate limb length discrepancy due to congenital anomalies, acquired deformities, growth plate (physeal injuries or surgery), or inborn errors of metabolism.

Reference

1. Leitzes A, Potter HG, Amaral T, et al. Reliability and accuracy of MRI scanogram in the evaluation of limb length discrepancy. *J Pediatr Orthop*. 2005;25(6):747-749.

MS-18: Anatomical Area Tables – General Information

The imaging guidelines for each anatomical area are presented in table format. The table below includes a description of how each column header should be utilized for each guideline **MS-19: Shoulder** through **MS-27: Foot**.

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.

MS-19: Shoulder

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Shoulder Pain	Yes	Yes	➤ MRI shoulder without contrast (CPT® 73221) OR ➤ US shoulder (CPT® 76881 or 76882) OR ➤ CT shoulder with contrast (arthrogram) (CPT® 73201) if MRI contraindicated	
Symptomatic Loose Bodies	Yes	No	➤ MRI shoulder without contrast (CPT® 73221)	
Impingement	Yes	Yes	➤ MRI shoulder without contrast (CPT® 73221) OR ➤ MRI shoulder with contrast (arthrogram) (CPT® 73222) OR ➤ US shoulder (CPT® 76881 or 76882) ➤ CT shoulder with contrast (CPT® 73201) if MRI is contraindicated	
Tendonitis/ Bursitis	Yes	Yes	➤ MRI shoulder without contrast (CPT® 73221) OR ➤ US shoulder (CPT® 76881 or 76882)	
Tendon Rupture (Biceps Long Head)	Yes	No	➤ When clinical exam is inconclusive due to inability to visualize a "Popeye" sign clinically, or for preoperative planning: ◆ MRI shoulder without contrast (CPT® 73221) OR ◆ US shoulder (CPT® 76881 or 76882)	
Tendon Rupture (Pectoralis Major/Minor)	Yes	No	➤ When clinical exam is inconclusive, or for preoperative planning: ◆ MRI Shoulder without contrast (CPT® 73221) OR ◆ MRI Chest without contrast (CPT® 71550) OR ◆ US shoulder (CPT® 76881 or 76882)	
Shoulder Rotator Cuff Tear (Complete and Partial)	Yes	Yes*	➤ MRI shoulder without contrast (CPT® 73221) OR ➤ MRI shoulder with contrast (arthrogram) (CPT® 73222) OR ➤ US shoulder (CPT® 76881 or 76882) ➤ CT shoulder with contrast	*Conservative treatment is not required with an acute shoulder injury prior to the onset of symptoms and consideration

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
			(arthrogram) (CPT® 73201) if MRI is contraindicated	of surgery. If surgery is being considered, MRI or CT arthrogram are required per <u>CMM-315: Shoulder Surgery- Arthroscopic and Open Procedures.</u>
Partial Tendon Rupture (Excluding Partial Rotator Cuff Tears)	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI shoulder without contrast (CPT® 73221) OR ◆ US shoulder (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/ muscle tears.
Shoulder Labral Tear (e.g., SLAP, ALPSA, HAGL)	Yes	Yes	➤ MRI shoulder with contrast (arthrogram) (CPT® 73222) OR ➤ MRI shoulder without contrast (CPT® 73221) OR ➤ CT shoulder with contrast (arthrogram) (CPT® 73201)	For surgery criteria, see: <u>CMM-315: Shoulder Surgery- Arthroscopic and Open Procedures.</u>
Shoulder Dislocation/ Subluxation/ Instability, or Bankart/ Hill-Sachs Lesions	Yes	Yes*	➤ Patients 40 years of age or younger with a first time dislocation, and in patients with recurrent dislocations, conservative treatment not required: ◆ MRI shoulder with contrast (arthrogram)	*Conservative treatment is required in patients over age 40 with a first time dislocation. For surgery

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
			(CPT® 73222) or ♦ MRI shoulder without contrast (CPT® 73221) ➤ CT shoulder with contrast (arthrogram) (CPT® 73201) or CT shoulder without contrast (CPT® 73200) if MRI is contraindicated	criteria, see: <u>CMM-315: Shoulder Surgery-Arthroscopic and Open Procedures.</u>
Frozen Shoulder/ Adhesive Capsulitis	Yes	Yes	➤ Advanced imaging is rarely indicated – in those rare situations, MRI shoulder without contrast (CPT® 73221)	For surgery criteria, see: <u>CMM-310: Manipulation Under Anesthesia</u> and <u>CMM-315: Shoulder Surgery-Arthroscopic</u>
Avascular Necrosis (AVN) of the Humeral Head	Yes	No	➤ MRI shoulder without contrast (CPT® 73221) when suspected and plain x-ray is negative or equivocal ➤ CT shoulder without contrast (CPT® 73200) and/or MRI shoulder without contrast (CPT® 73221) for preoperative planning prior to shoulder replacement	See also: <u>MS-4.1: AVN</u>
Acromio-clavicular (AC) Separation	Yes	No	➤ MRI shoulder without contrast (CPT® 73221) to rule out possible rotator cuff tear following AC separation	
Sterno-clavicular (SC) Dislocation	Yes	No	➤ CT Chest without contrast (CPT® 71250) if posterior SC dislocation is evident or suspected	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative Shoulder Surgery for Impingement, Rotator Cuff Tear, and/or Labral Tear	Yes	Yes	➤ In symptomatic individuals: ◆ MRI shoulder without contrast (CPT® 73221) OR ◆ MRI shoulder with contrast (arthrogram) (CPT® 73222) ➤ US shoulder (CPT® 76881 or 76882) is also appropriate in symptomatic individuals following rotator cuff repair ➤ CT shoulder with contrast (arthrogram) (CPT® 73201) if MRI contraindicated	
Preoperative Shoulder (Glenohumeral) Replacement Surgery	Yes	Yes	➤ CT shoulder without contrast (CPT® 73200) and/or MRI shoulder without contrast (CPT® 73221) for preoperative planning prior to shoulder replacement	See also: <u>MS-12: Osteoarthritis</u> For joint surgery criteria, see: <u>CMM-318: Shoulder Arthroplasty/Arthrodesis</u>
Post-Operative Shoulder (Glenohumeral) Replacement Surgery	Yes	No	➤ For suspected aseptic loosening or fracture as additional imaging following plain x-rays: ◆ CT shoulder without contrast (CPT® 73200) OR ◆ Bone scan (CPT® 78315) OR ◆ Distribution Of Radiopharmaceutical	See also: <u>MS-16: Post-Operative Joint Replacement</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
			Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ For suspected infection with negative or inconclusive joint aspiration culture: ◆ In-111 WBC (CPT® 78800, 78801, 78802, 78803, or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830, 78832) and Tc-99m sulfur colloid scan shoulder (CPT® 78102, 78103, or 78104) OR ◆ Bone scan (CPT® 78315, 78300, 78305, or 78306) ◆ See also: MS-28: Nuclear Medicine ➤ For possible rotator cuff tear: ◆ CT shoulder with contrast (arthrogram) (CPT® 73201) OR ◆ US shoulder (CPT® 76881 or 76882) ➤ For possible nerve injury: ◆ MRI shoulder without contrast (CPT® 73221) OR ◆ US shoulder (CPT® 76881 or 76882)	

References

1. Amini B, Beckmann NM, Beaman FD, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Traumatic. *Am Coll Radiol (ACR)*; Revised 2017. <https://acsearch.acr.org/docs/69433/Narrative/>.
2. Neviaser RJ, Neviaser TJ. Recurrent instability of the shoulder after age 40. *J Shoulder Elbow Surg.* 1995;4(6):416-418.
3. Bradley M, Tung G, Green A. Overutilization of shoulder magnetic resonance imaging as a diagnostic screening tool in patients with chronic shoulder pain. *J Shoulder Elbow Surgery.* 2005;14(3):233-237. doi:10.1016/j.jse.2004.08.002.
4. Fongemie AE, Buss DD, and Rolnick SJ. Management of shoulder impingement syndrome and rotator cuff tears. *Am Fam Physician.* 1998;57(4):667-674. <https://www.aafp.org/afp/1998/0215/p667.html>.
5. Griffin LY. *Essentials of Musculoskeletal Care.* 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:212.
6. Gyftopoulos S, Rosenberg ZS, Roberts CC, ET. Al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Shoulder Arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/3097049/Narrative/>.
7. Hovelius L, Olofsson A, Sandstrom B, et al. Nonoperative treatment of primary anterior shoulder dislocation in patients forty years of age and younger: a prospective twenty-five year follow-up. *J Bone Joint Surg.* 2008;90:945-52. doi:10.2106/JBJS.G.00070.
8. Lin A, Gasbarro G, Sakr M. Clinical Applications of Ultrasonography in the Shoulder and Elbow. *J Am Acad Orthop Surg.* 2018;26:303-312.
9. Magee T. 3-T MRI of the shoulder: is MR arthrography necessary? *AJR J Am Roentgenol.* 2009;192:86-92. doi:10.2214/AJR.08.1097.
10. Major NM, Browne J, Domzalski T, Cothran RL, Helms CA. Evaluation of the glenoid labrum with 3-T MRI: is intraarticular contrast necessary. *AJR Am J Roentgenol.* 2011;196:1139-1144. doi:10.2214/AJR.08.1734.
11. McDonald LS, Dewing CB, Shupe PG, et al. Disorders of the proximal and distal aspects of the biceps muscle. *J Bone Joint Surg.* 2013;95:1235-1245. doi:10.2106/JBJS.L.00221.
12. Petersen SA, Murphy TP. The timing of rotator cuff repair for the restoration of function. *Journal of Shoulder and Elbow Surgery.* 2011;20(1):62-68. doi:10.1016/j.jse.2010.04.045.
13. Rehman A, Robinson P. Sonographic evaluation of injuries of the pectoralis muscles. *AJR Am J Roentgenol.* 2005;184:1205-1211. doi:10.2214/ajr.184.4.01841205.
14. Small KM, Adler RS, Shah SH, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Shoulder Pain - Atraumatic. *Am Coll Radiol (ACR)*; New 2018. <https://acsearch.acr.org/docs/3101482/Narrative/>.
15. Steinbach LS, Chung CB, Yoshioka H. Technical Considerations for MRI of Upper Extremity Joints. In: Chung CB, Steinbach LS, eds. *MRI of the Upper Extremity Shoulder, Elbow, Wrist and Hand* Philadelphia, PA: Lippincott Williams & Wilkins 2010:211.
16. Streubel PN, Krych AJ, Simone JP, et al. Anterior glenohumeral instability: a pathology-based surgical treatment strategy. *J Am Acad Orthop Surg.* 2014;22:283-294. doi:10.5435/JAAOS-22-05-283.
17. Werner BC, Brockmeier SF, and Miller MD. Etiology, diagnosis, and management of failed SLAP repair. *J Am Acad Orthop Surg.* 2014;22(9):554-565. doi:10.5435/JAAOS-22-09-554.
18. Woodward TW and Best TM. The painful shoulder: Part II. Acute and chronic disorders. *Am Fam Physician.* 2000;61(11):3291-3300. <https://www.aafp.org/afp/2000/0601/p3291.html>.
19. Zappia M, Di Pietto F, Aliprandi A, et al. Multi-modal imaging of adhesive capsulitis of the shoulder. *Insights Imaging.* 2016;7:365-371.
20. Frankle MA, Teramoto A, Luo Z-P, Levy JC, Pupello D. Glenoid morphology in reverse shoulder arthroplasty: Classification and surgical implications. *Journal of Shoulder and Elbow Surgery.* 2009;18(6):874-885. doi:10.1016/j.jse.2009.02.013.

MS-20: Elbow				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Elbow Pain	Yes	Yes	➤ MRI elbow without contrast (CPT® 73221) OR ➤ US elbow (CPT® 76881 or 76882)	
Symptomatic Loose Bodies	Yes	No	➤ MRI elbow without contrast (CPT® 73221) if effusion is present OR ➤ MRI elbow with contrast (arthrogram) (CPT® 73222) if no effusion is present	
Tendonitis	Yes	Yes	➤ MRI elbow without contrast (CPT® 73221) or US elbow (CPT® 76881 or 76882)	
Bursitis	Yes	Yes	➤ MRI elbow without and with contrast (CPT® 73223) or MRI elbow without contrast (CPT® 73221) or US elbow (CPT® 76881 or 76882)	
Lateral (tennis elbow) or Medial (golfer's elbow) Epicondylitis	Yes	Yes	➤ To confirm clinical diagnosis of epicondylitis if symptoms persist for longer than 6 months despite at least 6 weeks conservative treatment in the last 3 months: ◆ MRI elbow without contrast (CPT® 73221) OR ◆ US elbow (CPT® 76881 or 76882)	Epicondylitis, caused by tendon degeneration and tear of the common extensor tendon laterally or of the common flexor tendon medially, is a common clinical diagnosis for which imaging is not medically necessary except as noted.

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Suspected Osteochondral Injury	Yes	No	➤ If plain x-rays are negative and an osteochondral fracture is still suspected: ◆ MRI elbow without contrast (CPT® 73221) OR ◆ MRI elbow with contrast (arthrogram) (CPT® 73222) OR ◆ CT elbow with contrast (arthrogram) (CPT® 73201)	See: <u>MS-13: Chondral/Osteochondral Lesions</u>
Ruptured Biceps Insertion at Elbow	Yes	No	➤ When clinical exam is inconclusive or for preoperative planning: ◆ MRI elbow without contrast (CPT® 73221) OR ◆ US elbow (CPT® 76881 or 76882)	
Ruptured Triceps Insertion at Elbow	Yes	No	➤ When clinical exam is inconclusive or for preoperative planning: ◆ MRI elbow without contrast (CPT® 73221) OR ◆ US elbow (CPT® 76881 or 76882)	
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI elbow without contrast (CPT® 73221) OR ◆ US elbow (CPT® 76881 or 76882)	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
Trauma	Yes	No	➤ When surgery is being considered: ◆ MRI elbow without contrast (CPT® 73221) OR ◆ CT elbow without contrast (CPT® 73200)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Ulnar Collateral Ligament (UCL) Tear	Yes	No	➤ Following acute or repetitive (including overhead throwing athletes) elbow trauma: ◆ MRI elbow with contrast (arthrogram) (CPT® 73222) OR ◆ MRI elbow without contrast (CPT® 73221) OR ◆ US elbow (CPT® 76881 or 76882)	
Suspected Nerve Abnormality	NA	NA	➤ See: <u>PN-2: Focal Neuropathy</u>	
Post-Operative	Yes	Yes	➤ CT elbow without contrast (CPT® 73200) in symptomatic post-operative patients following surgical treatment of complex fractures; or ➤ MRI elbow without contrast (CPT® 73221) in symptomatic post-operative patients following soft-tissue surgery	
Preoperative Elbow Replacement Surgery	Yes	Yes	➤ CT elbow without contrast (CPT® 73200) for preoperative planning prior to elbow replacement when congenital or post-traumatic deformities exist	See: <u>MS-12: Osteoarthritis</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative Elbow Replacement Surgery	Yes	No	➤ For suspected aseptic loosening or periprosthetic fracture when recent plain x-ray is nondiagnostic: ◆ CT elbow without contrast (CPT® 73200) OR ◆ Bone scan (CPT® 78315) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ For suspected infection with negative or inconclusive joint aspiration culture: ◆ In-111 WBC (CPT® 78800, 78801, 78802, 78803, or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830, or 78832) and Tc-99m sulfur colloid scan elbow (CPT® 78102, 78103, or 78104) OR ◆ Bone scan (CPT® 78315, 78300, 78305, or 78306) ◆ See: <u>MS-28: Nuclear Medicine</u>	

References

1. McDonald LS, Dewing CB, Shupe PG, et al. Disorders of the proximal and distal aspects of the biceps muscle. *J Bone Joint Surg*. 2013;95:1235-1245. doi:10.2106/JBJS.L.00221.
2. Torp-Pedersen TE, Torp-Pedersen ST, Qvistgaard E, et al. Effect of glucocorticosteroid injections in tennis elbow verified on colour Doppler ultrasonography: evidence of inflammation. *Br J Sports Med*. 2008;42(12):978-982. doi:10.1136/bjsm.2007.041285.
3. Johnson GW, Cadwallader K, Scheffel SB, et al. Treatment of lateral epicondylitis. *Am Fam Physician*. 2007;76(6):843-848. <https://www.aafp.org/afp/2007/0915/p843.html>.
4. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:279-280.
5. Hayes CW, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Elbow Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2015. <https://acsearch.acr.org/docs/69423/Narrative/>.
6. Bruce JR, Andrews JR. Ulnar collateral ligament injuries in the throwing athlete. *J Am Acad Orthop Surg*. 2014;22:315-325.
7. Beltran J, Rosenberg ZS. Diagnosis of compressive and entrapment neuropathies of the upper extremity: value of MR imaging. *AJR Am J Roentgenol*. 1994;163(3):525-531. doi:10.2214/ajr.163.3.8079837.
8. Lin A, Gasbarro G, Sakr M. Clinical Applications of Ultrasonography in the Shoulder and Elbow. *J Am Acad Orthop Surg*. 2018;26:303-312.

MS-21: Wrist				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Wrist Pain	Yes	Yes	➤ MRI wrist without contrast (CPT® 73221) OR ➤ US wrist (CPT® 76881 or 76882)	
Tendonitis	Yes	Yes	➤ MRI wrist without contrast (CPT® 73221) OR ➤ US wrist (CPT® 76881 or 76882)	
Kienbock's Disease (Avascular Necrosis (AVN) of the Lunate)/ Preiser's Disease (Avascular Necrosis (AVN) of the Scaphoid)	Yes	No	➤ MRI wrist without contrast (CPT® 73221) when suspected and plain x-ray is negative or equivocal ➤ If diagnosed on plain x-ray, CT wrist without contrast (CPT® 73200) or MRI wrist without contrast (CPT® 73221)	See also: <u>MS-4.1: AVN</u>
Suspected Navicular/ Scaphoid Fracture	Yes	No	➤ When suspected based on history and physical exam: ◆ MRI wrist without contrast (CPT® 73221) OR ◆ CT wrist without contrast (CPT® 73200)	See also: <u>MS-5.2: Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints</u>
Distal Radioulnar Joint (DRUJ) Instability	Yes	No	➤ CT of both wrists without contrast (CPT® 73200) (should include wrists in supination and pronation)	
Complex Distal Radius/ Ulna Fracture	Yes	No	➤ CT wrist without contrast (CPT® 73200)	
Carpal Tunnel Syndrome/ Ulnar Tunnel Syndrome	NA	NA	➤ See: <u>PN-2: Focal Neuropathy</u>	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Intrinsic Ligament (e.g. scapholunate)/ Triangular Fibrocartilage Complex (TFCC) Injuries	Yes	Yes	➤ MRI wrist with contrast (arthrogram) (CPT® 73222) OR ➤ CT wrist with contrast (arthrogram) (CPT® 73201)	
Complete Rupture of a Specific Named Tendon Not Otherwise Specified	Yes	No	➤ For preoperative planning: ◆ MRI wrist without contrast (CPT® 73221) OR ◆ US wrist (CPT® 76881 or 76882)	
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI wrist without contrast (CPT® 73221) OR ◆ US wrist (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/muscle tears.
Post-Operative	Yes	Yes	➤ CT wrist without contrast (CPT® 73200) in symptomatic patients following surgery for navicular/scaphoid fractures and complex distal radius/ulna fractures; or ➤ MRI wrist with contrast (arthrogram) (CPT® 73222) in symptomatic patients following DRUJ or TFCC surgery	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Preoperative Wrist Replacement Surgery	Yes	Yes	➤ CT wrist without contrast (CPT® 73200) for preoperative planning prior to wrist replacement when congenital or post-traumatic deformities exist	See: <u>MS-12: Osteoarthritis</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative Wrist Replacement Surgery	Yes	No	➤ For suspected aseptic loosening or periprosthetic fracture when recent plain x-ray is nondiagnostic: ◆ CT wrist without contrast (CPT® 73200) OR ◆ Bone scan (CPT® 78315, 78300, or 78306) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ For suspected infection with negative or inconclusive joint aspiration culture: ◆ In-111 WBC (CPT® 78800, 78801, 78802, 78803, or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830 or 78832) and Tc-99m sulfur colloid scan wrist (CPT® 78102, 78103, or 78104) OR ◆ Bone scan (CPT® 78315, 78300, 78305, or 78306) ◆ See: <u>MS-28:</u> <u>Nuclear Medicine</u>	

One Study/Area Only

In hand and wrist advanced imaging, studies are frequently ordered of both areas. This is unnecessary since wrist MRI will image from above the wrist to the mid-metacarpal area. **Only one CPT® code should be reported.**

References

1. Bruno MA, Weissman BN, Kransdorf MJ, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hand and Wrist Trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
2. Rubin DA, Roberts CC, Bencardino JT, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Wrist Pain. *Am Coll Radiol (ACR)*; Revised: 2017. <https://acsearch.acr.org/docs/69427/Narrative/>.
3. Hayter CL, Gold SL, Potter HG. Magnetic resonance imaging of the wrist: bone and cartilage injury. *J Magn Reson Imaging*. 2013;37(5):1005-19. doi:10.1002/jmri.23845.
4. Pruitt DL, Gilula LA, Manske PR, et al. Computed tomography scanning with image reconstruction in evaluation of distal radius fractures. *J Hand Surg Am*. 1994;19(5):720-727. doi:10.1016/0363-5023(94)90174-0.
5. Magee T. Comparison of 3-T MRI and arthroscopy of intrinsic wrist ligament and TFCC tears. *AJR Am J Roentgenol*. 2009;192:80-85. doi:10.2214/AJR.08.1089.
6. Lee RK, Ng AW, Tong CS, et al. Intrinsic ligament and triangular fibrocartilage complex tears of the wrist: comparison of MDCT arthrography, conventional 3-T MRI, and MR arthrography. *Skeletal Radiol*. 2013;42:1277-85. doi:10.1007/s00256-013-1666-8.
7. Pahwa S, Srivastava DN, Sharma R, et al. Comparison of conventional MRI and MR arthrography in the evaluation wrist ligament tears: A preliminary experience. *Indian J Radiol Imaging*. 2014;3:259-67. doi:10.4103/0971-3026.137038.

MS-22: Hand				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Hand Pain	Yes	Yes	➤ MRI hand or finger without contrast (CPT® 73218) OR ➤ US hand (CPT® 76881 or 76882)	
Tendonitis	Yes	Yes	➤ MRI hand or finger without contrast (CPT® 73218) OR ➤ US hand or finger (CPT® 76881 or 76882)	
Occult Fracture	Yes	No	➤ Advanced imaging guided by: <u>MS-5.2: Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints</u>	
Complex Fracture	Yes	No	➤ CT hand or finger without contrast (CPT® 73200) when plain x-ray shows a complex fracture	
Ulnar Collateral Ligament (UCL) Thumb Injury	Yes	No	➤ If rule out for Stener lesion or complete tear of UCL of the thumb MCP joint: ◆ MRI thumb without contrast (CPT® 73218) OR ◆ US thumb (CPT® 76881 or 76882)	Also called "Gamekeeper's Thumb" or "Skier's Thumb"
Complete Rupture of a Specific Named Tendon not Otherwise Specified	Yes	No	➤ For preoperative planning: ◆ MRI hand or finger without contrast (CPT® 73218) OR ◆ US hand or finger (CPT® 76881 or 76882)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI hand or finger without contrast (CPT® 73218) OR ◆ US hand or finger (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/muscle tears.
Post-Operative	Yes	Yes	➤ In symptomatic post-operative patients following surgical treatment for complex hand or finger fractures or following soft-tissue surgery: ◆ CT hand or finger without contrast (CPT® 73200) OR ◆ MRI hand or finger without contrast (CPT® 73218)	

One Study/Area Only

In hand and wrist advanced imaging, studies are frequently ordered of both areas. This is unnecessary since wrist MRI will image from above the wrist to the mid-metacarpal area. **Only one CPT® code should be reported.**

References

1. Bruno MA, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hand and Wrist Trauma. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69418/Narrative/>.
2. Hayter CL, Gold SL, Potter HG. Magnetic resonance imaging of the wrist: Bone and cartilage injury. *J Magn Reson Imaging*. 2013;37(5):1005-19. doi:10.1002/jmri.23845.

MS-23: Pelvis

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Pain-Pelvis	Yes	Yes	➤ MRI pelvis without contrast (CPT® 72195) OR ➤ MRI RT and/or LT hip without contrast (CPT® 73721)	
Tendonitis	Yes	Yes	➤ MRI pelvis without contrast (CPT® 72195) OR ➤ MRI RT and/or LT hip without contrast (CPT® 73721)	
Occult/ Insufficiency Fracture	Yes	No	➤ MRI pelvis without contrast (CPT® 72195) OR ➤ CT pelvis without contrast (CPT® 72192)	See also: <u>MS-5.2: Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints</u> for occult and stress fractures of the pelvis
Complex Fracture/ Dislocation - Pelvis, Sacrum and Acetabulum	Yes	No	➤ CT pelvis without contrast (CPT® 72192)	Additionally, 3D rendering may be appropriate for preoperative planning. See also: <u>MS-3: 3D Rendering</u>
Sacro-iliac (SI) Joint Pain, Sacroiliitis, Coccydynia	Yes	Yes	➤ Advanced imaging guided by: ◆ <u>SP-10.1: Sacroiliac (SI) Joint Pain/ Sacroiliitis</u> ◆ <u>SP-5.2: Coccydynia without Neurological Features</u>	
Complete Rupture of a Specific Named Tendon	Yes	No	➤ MRI pelvis without contrast (CPT® 72195) for preoperative planning	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Partial Tendon Rupture	Yes	No	➤ MRI Pelvis without contrast (CPT® 72195) for a suspected partial tendon rupture of a specific named tendon not otherwise specified	MRI is <i>NOT</i> needed for muscle belly strains/muscle tears.
Osteitis Pubis/ Symphysis Pubis Diastasis	Yes	Yes	➤ MRI Pelvis without contrast (CPT® 72195)	
Athletic Pubalgia (Sports Hernia)	Yes	Yes	➤ To evaluate for the cause of suspected athletic pubalgia: ♦ MRI pelvis without contrast (athletic pubalgia protocol) (CPT® 72195) OR ♦ Dynamic pelvic ultrasound (CPT® 76857)	
Post-Operative	Yes	Yes	➤ CT pelvis without contrast (CPT® 72192) in symptomatic patients following surgery for complex pelvic ring/acetabular fractures	

References

1. Bencardino JT, Stone TJ, Roberts CC, et al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
2. Mehta S, Auerbach JD, Born CT, et al. Sacral fractures. *J Am Acad Orthop Surg*. 2006;14:656-665.
3. Omar IM, Zoga AC, Kavanagh EC, et al. Athletic Pubalgia and "Sports Hernia": Optimal MR Imaging Technique and Findings. *RadioGraphics*. 2008;28:1415-1438. doi:10.1148/rg.285075217.
4. Khan W, Zoga AC, Meyers WC. Magnetic Resonance Imaging of Athletic Pubalgia and the Sports Hernia - Current Understanding and Practice. *Magn Reson Imaging Clin N Am*. 2013;21:97-110. doi:10.1016/j.mric.2012.09.008.
5. Morley N, Grant T, Blount K, et al. Sonographic evaluation of athletic pubalgia. *Skeletal Radiol*. 2016 May;45(5):689-99. doi:10.1007/s00256-016-2340-8.
6. Caudill P, Nyland J, Smith C, et al. Sports hernias: a systematic literature review. *British Journal of Sports Medicine*. 2008;42(12):954-964. doi:10.1136/bjsm.2008.047373.
7. Suarez JC, Ely EE, Mutnal AB, et al. Comprehensive approach to the evaluation of groin pain. *Journal of the American Academy of Orthopaedic Surgeons*. 2013;21:558-570. doi:10.5435/JAAOS-21-09-558.
8. Heer ST, Callander JW, Kraeutler MJ, Mei-Dan O, Mulcahey MK. Hamstring Injuries. *The Journal of Bone and Joint Surgery*. 2019;101(9):843-853. doi:10.2106/jbjs.18.00261.

MS-24: Hip				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Hip Pain	Yes	Yes	➤ MRI hip without contrast (CPT® 73721) OR ➤ US hip (CPT® 76881 or 76882)	
Symptomatic Loose Bodies	Yes	No	➤ MRI hip without contrast (CPT® 73721)	
Tendonitis/ Bursitis	Yes	Yes	➤ MRI hip without contrast (CPT® 73721) OR ➤ US hip (CPT® 76881 or 76882)	
Hip Abductor Tendon Tear/ Avulsion	Yes	No	➤ MRI hip without contrast (CPT® 73721) OR ➤ US hip (CPT® 76881 or 76882)	
Complete Rupture of a Specific Named Tendon	Yes	No	➤ For preoperative planning: ◆ MRI hip without contrast (CPT® 73721) OR ◆ US hip (CPT® 76881 or 76882)	
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI hip without contrast (CPT® 73721) OR ◆ US hip (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/ muscle tears.
Occult/ Insufficiency Fracture	Yes	No	➤ MRI hip without contrast (CPT® 73721) OR ➤ CT hip without contrast (CPT® 73700)	See also: <u>MS-5.2: Suspected Occult/ Stress/ Insufficiency Fracture/ Stress Reaction and Shin Splints</u> for occult and stress fractures of the hip

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Avascular Necrosis (AVN) of the Femoral Head	Yes	No	➤ MRI hip without contrast (CPT® 73721) when suspected and plain x-ray is negative or equivocal ➤ MRI hip without contrast (CPT® 73721) or CT hip without contrast (CPT® 73700) with femoral head collapse for preoperative planning	See also: <u>MS-4.1: AVN</u>
Labral Tear	Yes	Yes	➤ MRI hip with contrast (arthrogram) (CPT® 73722) OR ➤ CT hip with contrast (arthrogram) (CPT® 73701) OR ➤ MRI hip without contrast (CPT® 73721)	For surgery criteria, see: <u>CMM-314: Hip Surgery-Arthroscopic and Open Procedures</u>
Femoroacetabular Impingement	Yes	Yes	➤ For preoperative planning for femoroacetabular impingement: ◆ MRI hip without contrast (CPT® 73721) OR ◆ MRI hip with contrast (arthrogram) (CPT® 73722) - IN ADDITION TO: ◆ CT hip without contrast (CPT® 73700) OR ◆ CT pelvis without contrast (CPT® 72192)	For surgery criteria, see: <u>CMM-314: Hip Surgery-Arthroscopic and Open Procedures</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Piriformis Syndrome	Yes	Yes	➤ For preoperative planning: ◆ MRI pelvis without contrast (CPT® 72195) OR ◆ CT pelvis without contrast (CPT® 72192)	EMG/NCV may confirm the diagnosis. Refer to <u>PN-2: Focal Neuropathy</u>
Post-Operative	Yes	Yes	➤ Symptomatic patients following surgery for labral tears and femoroacetabular impingement: ◆ MRI hip with contrast (arthrogram) (CPT® 73722) ➤ Symptomatic patients following surgery for hip fracture and/or hip avascular necrosis: ◆ CT hip without contrast (CPT® 73700) OR ◆ MRI hip without contrast (CPT® 73721)	
Preoperative Hip Replacement Surgery	Yes	Yes	➤ CT hip without contrast (CPT® 73700) for preoperative planning prior to hip replacement when congenital or post-traumatic deformities exist	See also: <u>MS-12: Osteoarthritis</u> For surgery criteria, see: <u>CMM-313: Hip Arthroplasty- Total and Partial</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative Hip Replacement Surgery	Yes	No*	➤ For suspected aseptic loosening of hip replacement when recent plain x-ray is nondiagnostic: ◆ CT hip without contrast (CPT® 73700) OR ◆ Bone scan (CPT® 78315) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ For suspected infection with negative or inconclusive joint aspiration culture: ◆ In-111 WBC (CPT® 78800, 78801, 78802, 78803, or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830 or 78832) and Tc-99m sulfur colloid scan hip (CPT® 78102, 78103, or 78104) OR ◆ Bone scan (CPT® 78315, 78300, 78305, or 78306) ◆ See <u>MS-28: Nuclear Medicine</u> ➤ For suspicion of a periprosthetic fracture when recent plain x-ray is nondiagnostic:	See also: <u>MS-16: Post-Operative Joint Replacement</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative Hip Replacement Surgery (continued)	Yes	No*	◆ CT hip without contrast (CPT® 73700) OR ◆ Bone scan (CPT® 78315) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ To evaluate component malposition or heterotopic bone after plain x-ray: ◆ CT hip without contrast (CPT® 73700) ➤ For possible nerve injury: ◆ MRI hip without contrast (CPT® 73721) ➤ For suspected for suspected tendinitis/bursitis (*requires conservative treatment): ◆ MRI hip without contrast (CPT® 73721) OR ◆ US hip (CPT® 76881 or 76882)	

References

1. Greene WB (Ed.). *Essentials of Musculoskeletal Care*. 2nd Ed. Rosemont, IL, American Academy of Orthopaedic Surgeons, 2001.
2. Manek NJ and Lane NE. Osteoarthritis: Current concepts in diagnosis and management. *Am Fam Physician*. 2000;61(6):1795-1804. <https://www.aafp.org/afp/2000/0315/p1795.html>.
3. Papadopoulos EC and Kahn SN. Piriformis syndrome and low back pain: a new classification and review of the literature. *Orthop Clin North Am*. 2004;35(1):65-71. doi:10.1016/S0030-5898(03)00105-6.
4. Reurink G, Sebastian, Bisselink JM, et al. Reliability and Validity of Diagnostic Acetabular Labral Lesions with Magnetic Resonance Arthrography. *J Bone Joint Surg A*. 2012;94(181):1643-1648. doi:10.2106/JBJS.K.01342.
5. Steinbach LS, Palmer WE, and Schweitzer ME. Special Focus Session MR Arthrography1. *RadioGraphics*. 2002;22(5):1223-1246.
6. Redmond JM, Chen AW, and Domb BG. Greater Trochanteric Pain Syndrome. *J Am Acad Orthop Surg*. 2016;24(4):231-240. doi:10.5435/JAAOS-D-14-00406.
7. Center for Devices and Radiological Health. Metal-on-Metal Hip Implants - Information for Orthopaedic Surgeons. U S Food and Drug Administration Home Page.
8. Ward RJ, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Hip Pain-Suspected Fracture. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Last Review: 2018. <https://acsearch.acr.org/docs/3082587/Narrative/>.
9. Weissman BN, Palestro CJ, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Total Hip Arthroplasty. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2015. <https://acsearch.acr.org/docs/3094200/Narrative/>.
10. Mintz DN, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Hip Pain. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69425/Narrative/>.
11. Murphey MD, Roberts CC, Bencardino JT, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Osteonecrosis of the Hip. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2015. <https://acsearch.acr.org/docs/69420/Narrative/>.
12. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
13. Verberne SJ, Rajmakers PG, and Temmerman OP. The Accuracy of Imaging Techniques in the Assessment of Periprosthetic Hip Infection. A Systematic Review and Meta-Analysis. *J Bone Joint Surg Am*. 2016;98(19):1638-45. doi:10.2106/JBJS.15.00898.
14. Shin AY, Morin WD, Gorman JD, et al. The superiority of magnetic resonance imaging in differentiating the cause of hip pain in endurance athletes. *Am J Sports Med*. 1996;24:168-76. doi:10.1177/036354659602400209.
15. Slocum KA, Gorman JD, Puckett ML, et al. Resolution of abnormal MR signal intensity in patients with stress fractures of the femoral neck. *AJR Am J Roentgenol*. 1997;168:1295-9. doi:10.2214/ajr.168.5.9129429.
16. Lee EY, Margherita AJ, Gierada DS, et al. MRI of Piriformis Syndrome. *American Journal of Roentgenology*. 2004;183:63-64. doi:10.2214/ajr.183.1.1830063.
17. Jankiewicz JJ, Hennrikus WL, and Houkom JA. The appearance of the piriformis muscle syndrome in computed tomography and magnetic resonance imaging: a case report and review of the literature. *Clin Orthop*. 1991;262:205-09.
18. Rossi P, Cardinali P, Serrao M, et al.. Magnetic resonance imaging findings in piriformis syndrome: a case report. *Arch Phys Med Rehabil*. 2001;82(4):519-21. doi:10.1053/apmr.2001.21971.
19. Heer ST, Callander JW, Kraeutler MJ, Mei-Dan O, Mulcahey MK. Hamstring Injuries. *The Journal of Bone and Joint Surgery*. 2019;101(9):843-853. doi:10.2106/jbjs.18.00261.

MS-25: Knee				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Knee Pain	Yes	Yes	➤ MRI knee without contrast (CPT® 73721) - OR ➤ US knee (CPT® 76881 or 76882)	
Symptomatic Loose Bodies	Yes	No	➤ MRI knee without contrast (CPT® 73721) ➤ CT knee with contrast (arthrogram) (CPT® 73701) if MRI cannot be performed	
Tendonitis	Yes	Yes	➤ MRI knee without contrast (CPT® 73721) - OR ➤ US knee (CPT® 76881 or 76882)	
Complex Knee Fracture	Yes	No	➤ MRI knee without contrast (CPT® 73721) - OR ➤ CT knee without contrast (CPT® 73700)	See: <u>MS-5: Fractures</u>
Meniscus Tear	Yes	Yes*	➤ MRI knee without contrast (CPT® 73721) *Conservative treatment is not required if at least 2 of following 4 criteria are met: 1) Positive McMurray's, positive Thessaly, or positive Apley's Compression Test 2) twisting or acute injury of the knee 3) locked knee/inability to fully extend the knee 4) knee effusion ➤ MRI knee without contrast (CPT® 73721) for clinical suspicion of a symptomatic degenerative meniscus tear in a patient with osteoarthritis following conservative treatment	For surgery criteria, see: <u>CMM-312: Knee Surgery-Arthroscopic and Open Procedures</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved ? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Ligament Tear	Yes	Yes*	➤ MRI knee without contrast (CPT® 73721) *Conservative treatment is not required if any of the following signs are positive in comparison to the normal knee: ◆ Anterior drawer ◆ Lachman ◆ Pivot shift ◆ Posterior drawer ◆ Posterior sag ◆ Valgus stress ◆ Varus stress	For surgery criteria, see: <u>CMM-312: Knee Surgery-Arthroscopic and Open Procedures</u>
Knee Joint Dislocation	Yes	No	➤ Following significant trauma to evaluate for ligament and vascular injury: ◆ MRI knee without contrast (CPT® 73721) AND EITHER ◆ MR Angiography lower extremity without and with contrast (CPT® 73725) OR ◆ CT Angiography lower extremity without and with contrast (CPT® 73706)	
Patellar Dislocation/Subluxation	Yes	No	➤ MRI knee without contrast (CPT® 73721) with acute knee injury, consideration of surgery and concern for osteochondral fracture or loose osteochondral fracture fragment	For surgery criteria, see: <u>CMM-312: Knee Surgery-Arthroscopic and Open Procedures</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Recurrent Patellar Instability	Yes	Yes	➤ MRI knee without contrast (CPT® 73721) if consideration for surgery	For surgery criteria, see: <u>CMM-312: Knee Surgery-Arthroscopic and Open Procedures</u>
Patellofemoral Pain Syndrome/ Anterior Knee Pain/ Tracking Disorder	Yes	Yes	➤ MRI knee without contrast (CPT® 73721) if consideration for surgery	
Suspected Osteochondral Injury	Yes	No	➤ If plain x-rays are negative and an osteochondral fracture is still suspected: ◆ MRI knee without contrast (CPT® 73721) OR ◆ MRI knee with contrast (arthrogram) (CPT® 73722) OR ◆ CT knee with contrast (arthrogram) (CPT® 73701)	See: <u>MS-13: Chondral Osteochondral Lesions</u> for other osteochondral injury scenarios. For surgery criteria, see: <u>CMM-312: Knee Surgery-Arthroscopic and Open Procedures</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Avascular Necrosis (AVN) of the Distal Femur	Yes	No	➤ MRI knee without contrast (CPT® 73721) when suspected and plain x-ray is negative or equivocal or with AVN confirmed by plain x-ray if needed for treatment planning	See: <u>MS-4.1: Avascular Necrosis</u>
Baker's Cyst (Popliteal Cyst)	Yes	Yes	➤ US knee (CPT® 76881 or 76882) is the initial imaging study ➤ MRI knee without contrast (CPT® 73721) for preoperative planning	See also: <u>PVD-12: Acute Limb Swelling</u>
Plica (Symptomatic Synovial Plica/ Medial Synovial Shelf)	Yes	Yes	➤ MRI knee without contrast (CPT® 73721)	
Hemarthrosis	Yes	No	➤ MRI knee without contrast (CPT® 73721) for clinical suspicion of cruciate ligament tear (requires a positive objective sign for ACL/PCL tear) or patellar dislocation (requires a positive apprehension sign) ➤ CT knee without contrast (CPT® 73700) for clinical suspicion of non-displaced intra-articular fracture	
Complete Rupture of the Distal Quadriceps Tendon or Patellar Ligament/ Tendon	Yes	No	➤ For preoperative planning: ◆ MRI knee without contrast (CPT® 73721) OR ◆ US knee (CPT® 76881 or 76882)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI knee without contrast (CPT® 73721) OR ◆ US knee (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/ muscle tears.
Post-Operative	Yes	Yes	➤ In symptomatic patients following surgery for meniscus tears and reconstruction of the anterior cruciate ligament: ◆ MRI knee with contrast (arthrogram) (CPT® 73722) OR ◆ MRI knee without contrast (CPT® 73721) ➤ In symptomatic patients following surgery for fracture/dislocation: ◆ CT knee without contrast (CPT® 73700)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Preoperative Knee Replacement Surgery	Yes	Yes	➤ CT knee without contrast (CPT® 73700) for preoperative planning prior to knee replacement when congenital or post-traumatic deformities exist of the patella, distal femur and/or proximal tibia	See also: <u>MS-12: Osteoarthritis</u> For surgery criteria, see: <u>CMM-311: Knee Arthroplasty- Total and Partial</u>
Post-Operative Knee Replacement Surgery	Yes	No*	➤ For suspected aseptic loosening when recent plain x-ray is nondiagnostic: ◆ CT knee without contrast (CPT® 73700) OR ◆ Bone scan (CPT® 78315) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ For suspected infection with negative or inconclusive joint aspiration culture: ◆ Tc-99m 3-phase bone scan (CPT® 78315) and In-111 WBC scan knee (CPT® 78800, 78801, 78802, 78803, or 78831) OR	See also: <u>MS-16: Post-Operative Joint Replacement Surgery</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative Knee Replacement Surgery (continued)	Yes	No*	◆ In-111 WBC (CPT® 78800-78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830, or 78832) and Tc-99m sulfur colloid scan knee (CPT® 78102, 78103, or 78104) OR ◆ Bone scan (CPT® 78315, 78300, 78305, or 78306) ◆ See <u>MS-28: Nuclear Medicine</u> ➤ Following plain x-ray for suspected periprosthetic fracture: ◆ CT knee without contrast (CPT® 73700) OR ◆ Bone scan (CPT® 78315) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830) ➤ For suspected osteolysis or component instability, rotation, or wear: ◆ CT knee without contrast (CPT® 73700) OR ◆ MRI knee without contrast (CPT® 73721)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
			➤ For suspected periprosthetic soft tissue abnormality unrelated to infection (e.g., tendinopathy, arthrofibrosis, patellar clunk syndrome, impingement of nerves or other soft tissue) *requires conservative treatment: ◆ MRI knee without contrast (CPT® 73721) OR ◆ US knee (CPT® 76881 or 76882)	

References

1. Harrison BK, Abell BE, Gibson TW. The Thessaly test for detection of meniscal tears: validation of a new physical examination technique for primary care medicine. *Clin J Sport Med*. 2009;19:9-12. doi:10.1097/JSM.0b013e31818f1689.
2. Landewé RBM, Günther KP, Lukas C, et al. EULAR/EFFORT recommendations for the diagnosis and initial management of patients with acute or recent onset swelling of the knee. *Ann Rheum Dis*. 2010;69:12-19. doi:10.1136/ard.2008.104406.
3. Johnson MW. Acute knee effusions: a systematic approach to diagnosis. *Am Fam Physician*. 2000;61(8):2391-2400. <https://www.aafp.org/afp/2000/0415/p2391.html>.
4. ACR Appropriateness Criteria, Nontraumatic knee pain, 2008.
5. Sung-Jae Kim, Byoung-Yoon Hwang, Choi DH, et al. *J Bone Joint Surg A*. 2012;94(16):e118 1-7.
6. Kannus P, Järvinen M. Nonoperative treatment of acute knee ligament injuries. A review with special reference to indications and methods. *Sports Med*. 1990;9(4):244-260. doi:10.2165/00007256-199009040-00005.
7. Manek NJ and Lane NE. Osteoarthritis: Current concepts in diagnosis and management. *Am Fam Physician*. 2000;61(6):1795-1804. <https://www.aafp.org/afp/2000/0315/p1795.html>.
8. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:84, 541-545.
9. Lee IS, Choi JA, Kim TK, et al. Reliability analysis of 16-MDCT in preoperative evaluation of total knee arthroplasty and comparison with intraoperative measurements. *Am J Roentgenol*. 2006;186(6):1778-1782. doi:10.2214/AJR.05.1191.
10. Morrissey RT, Weinstein SL (Eds.). *Lovell and Winter's Pediatric Orthopaedics*. 6th Ed. Philadelphia, PA: Lippincott Williams and Wilkins; 2005:1413.
11. Woolson ST, Harris AHS, Wagner DW, et al; Component alignment during total knee arthroplasty with use of standard or custom instrumentation: A Randomized Clinical Trial Using Computed Tomography for Postoperative Alignment Measurement. *Journal of Bone and Joint Surgery*. 2014;96:366-372. doi:10.2106/JBJS.L.01722.
12. Vance K, Meredick R, Schweitzer ME, et al. Magnetic resonance imaging of the postoperative meniscus. *Arthroscopy*. 2009;25:522-30. doi:10.1016/j.arthro.2008.08.013.
13. Magee T, Shapiro M, and Williams D. Prevalence of meniscal radial tears of the knee revealed by MRI after surgery. *Am J Roentgenol*. 2004;184:931-936. doi:10.2214/ajr.182.4.1820931.
14. Meyers AB, Haims AH, Menn K, et al. Imaging of anterior cruciate ligament repair and its complications. *Am J Roentgenol*. 2010;194:476-484. doi:10.2214/AJR.09.3200.
15. Tuite, MJ, Kransdorf MJ, Beaman FD, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Knee. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2019. <https://acsearch.acr.org/docs/69419/Narrative/>.
16. Bennett DL, Nelson JW, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Nontraumatic Knee Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2018. <https://acsearch.acr.org/docs/69432/Narrative/>.
17. Zoga AC, Weissman BN, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Soft-Tissue Masses. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2017. <https://acsearch.acr.org/docs/69434/Narrative/>.
18. Hochman MG, Melenevsky YV, Metter DF, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Imaging After Total Knee Arthroplasty. *Am Coll Radiol (ACR)*; Revised: 2017. Available at: <https://acsearch.acr.org/docs/69430/Narrative/>.
19. Kurosaka M, Yagi M, Yoshiya S, Muratsu H, Mizuno K. Efficacy of the axially loaded pivot shift test for the diagnosis of a meniscal tear. *Int Orthop*. 1999;23:271-274. doi:10.1007/s002640050369.
20. Fowler PJ, Lubliner JA. The predictive value of five clinical signs in the evaluation of meniscal pathology. *Arthroscopy*. 1989;5(3):184-186. doi:10.1016/0749-8063(89)90168-0.

MS-26: Ankle				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Ankle Pain	Yes	Yes	➤ MRI ankle without contrast (CPT® 73721) OR ➤ US ankle (CPT® 76881 or 76882)	
Symptomatic Loose Bodies	Yes	No	➤ MRI ankle without contrast (CPT® 73721)	
Complex Fracture	Yes	No	➤ MRI ankle without contrast (CPT® 73721) OR ➤ CT ankle without contrast (CPT® 73700)	
Ankle Sprain, Including Avulsion Fracture	Yes	Yes	➤ MRI ankle without contrast (CPT® 73721) OR ➤ CT without contrast (CPT® 73700)	
High Ankle Sprain (Syndesmosis Injury)	Yes	No	➤ MRI ankle without contrast (CPT® 73721) OR ➤ CT ankle without contrast (CPT® 73700)	
Suspected Osteochondral Injury	Yes	No	➤ If plain x-rays are negative and an osteochondral fracture is still suspected, ONE of the following: ◆ MRI ankle without contrast (CPT® 73721) ◆ CT ankle without contrast (CPT® 73700)	See: <u>MS-13: Chondral/Osteochondral Lesions</u> for other osteochondral injury scenarios
Avascular Necrosis (AVN) of the Talus	Yes	No	➤ MRI ankle without contrast (CPT® 73721) when suspected and plain x-ray is negative or equivocal or with plain x-ray-confirmed AVN if needed for treatment planning	See: <u>MS-4.1: AVN</u>
Anterior Impingement	Yes	Yes	➤ MRI ankle with contrast (arthrogram) (CPT® 73722) OR	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Anterior-Lateral Impingement Posterior Impingement (e.g., Os Trigonum Syndrome)			➤ CT ankle with contrast (arthrogram) (CPT® 73701) OR ➤ MRI ankle without contrast (CPT® 73721)	
Tendonitis	Yes	Yes	➤ For suspected posterior tibial dysfunction, peroneal tendon or subluxation, Achilles tendonitis: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ US ankle (CPT® 76881 or 76882)	
Ruptured Achilles Tendon	Yes	No	➤ For preoperative evaluation: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ US ankle (CPT® 76881 or 76882)	
Complete Rupture -Tear of a Specific Named Tendon	Yes	No	➤ For preoperative planning: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ US ankle (CPT® 76881 or 76882)	
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ US ankle (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/ muscle tears.
Instability	Yes	Yes	➤ For preoperative evaluation: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ MRI ankle with contrast (arthrogram) (CPT® 73722)	
Charcot Ankle	Yes	Yes	➤ MRI ankle without contrast (CPT® 73721)	
Post-Operative	Yes	Yes	➤ In symptomatic patients following surgery for	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
			ligament/tendon injuries, one of the following: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ US ankle (CPT® 76881 or 76882) ➤ For symptomatic patients following surgery for complex fractures: ◆ CT ankle without contrast (CPT® 73700)	
Preoperative Ankle Replacement Surgery	Yes	Yes	➤ CT ankle without contrast (CPT® 73700) for preoperative planning prior to ankle replacement when congenital or post-traumatic deformities exist	See also <u>MS-12: Osteoarthritis</u>
Post-Operative Ankle Replacement Surgery	Yes	No	➤ For suspected aseptic loosening or periprosthetic fracture when recent plain x-ray is nondiagnostic: ◆ CT ankle without contrast (CPT® 73700) OR ◆ Bone scan (CPT® 78315, 78300, or 78306) OR ◆ Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830)	See also: <u>MS-16: Post-Operative Joint Replacement Surgery</u>

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
			➤ For suspected infection with negative or inconclusive joint aspiration culture: ◆ In-111 WBC (CPT® 78800, 78801, 78802, 78803, or 78831) and Tc-99m 3-phase bone scan (CPT® 78315) OR ◆ In-111 WBC (CPT® 78800-78803 or 78831) OR ◆ Hybrid SPECT/CT (CPT® 78830 or 78832) and Tc-99 sulfur colloid scan ankle (CPT® 78102, 78103, or 78104) OR ◆ Bone scan (CPT® 78315, 78300, 78305, or 78306) ➤ See: <u>MS-28: Nuclear Medicine</u>	

One Study/Area Only

In foot and ankle advanced imaging, studies are frequently ordered of both areas. This is unnecessary since ankle MRI will image from above the ankle to the mid-metatarsal area. **Only one CPT® code should be reported.**

References

1. Donovan A, Rosenberg ZS. MRI of ankle and lateral hindfoot impingement syndromes. *AJR*. 2010;195: 595-604.
2. Wolfe MW, Uhl TL, and McClusky LC. Management of ankle sprains. *Am Fam Physician* 2001;63(1):93-104. <https://www.aafp.org/afp/2001/0101/p93.html>.
3. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:593-596; 606-609; 683.
4. Bergkvist D, Astrom I, Josefsson PO, et al. Acute Achilles Tendon Rupture: A Questionnaire Follow-up of 487 Patients. *J Bone Joint Surg Am*. 2012;94(13):1229-1233. doi:10.2106/JBJS.J.01601.
5. Hartgerink P, Fessell DP, Jacobson JA, et al. Full- versus partial-thickness Achilles tendon tears: sonographic accuracy and characterization in 26 cases with surgical correlation. *Radiology* 2001;220:406-412. doi: 10.1148/radiology.220.2.r01au41406.
6. Jones MP, Riaz JK, Smith RLC. Surgical Interventions for Treating Acute Achilles Tendon Rupture: Key Findings from a Recent Cochrane Review. *J Bone Joint Surg Am*. 2012;94(12):e88 1-6. doi:10.2106/jbjs.j.01829.
7. Vaseenon T, Amendola A. Update on anterior ankle impingement. *Current Reviews in Musculoskeletal Medicine*. 2012;5:140-150. doi:10.1007/s12178-012-9117-z.
8. Talusan PG, Toy J, Perez J, Milewski MD, et al. Anterior ankle impingement: diagnosis and treatment. *J Am Acad Orthop Surg*. 2014;22:333-339. doi:10.5435/JAAOS-22-05-333.
9. Nault ML, Kocher MS, Micheli LJ. Os Trigonum Syndrome. *J Am Acad Orthop Surg*. 2014;22:545-553. doi:10.5435/JAAOS-22-09-545.
10. Peace KAL, Jillier JC, Hulme A, et al. MRI features of posterior ankle impingement syndrome in ballet dancers: a review 25 cases. *Clinical Radiology*. 2004;59:1025-1033. doi:10.1016/j.crad.2004.02.010.
11. J Kane and R Zell. Achilles Tendon Rupture. Physician Resource Center. *American Orthopaedic Foot & Ankle Society*. Last reviewed July 2015.
12. Garras DN, et al. MRI is Unnecessary for Diagnosing Acute Achilles Tendon Ruptures. *Clinical Orthopaedics and Related Research*. 2012;470:2268–2273 Retrospective Analysis with finding. doi:10.1007/s11999-012-2355-y.
13. Mosher TJ, Kransdorf MJ, Adler R, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Ankle. *Am Coll Radiol (ACR)*; Date of Origin: 2013. Last review: 2020. <https://acsearch.acr.org/docs/69436/Narrative/>.
14. Luchs JS, Flug JA, Weissman BN, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Ankle Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2017. <https://acsearch.acr.org/docs/69422/Narrative/>.
15. Dodd A, Daniels TR. Charcot Neuroarthropathy of the Foot and Ankle. *J Bone Joint Surg Am*. 2018;100:696-711. doi:10.2106/JBJS.17.00785.

MS-27: Foot				
Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
General Foot Pain	Yes	Yes	➤ MRI foot without contrast (CPT® 73718)	
Complex Fractures	Yes	No	➤ CT foot without contrast (CPT® 73700)	
Plantar Plate Disorders, Including Turf Toe Injuries	Yes	Yes	➤ MRI foot without contrast (CPT® 73718)	
Sesamoid Disorders	Yes	Yes	➤ MRI foot without contrast (CPT® 73718) OR ➤ CT foot without contrast (CPT® 73700)	
Lisfranc Tarsometatarsal Fracture or Dislocation	Yes	No	➤ MRI foot without contrast (CPT® 73718) OR ➤ CT foot without contrast (CPT® 73700)	
Tarsal Navicular Stress/Occult Fracture	Yes	No	➤ MRI foot without contrast (CPT® 73718) ➤ Tc-99m bone scan foot (CPT® 78315) if MRI cannot be performed ➤ CT foot without contrast (CPT® 73700) for follow-up of healing fractures	See also: <u>MS-5.2: Suspected Occult/ Stress/ In-sufficiency Fracture/ Stress Reaction and Shin Splints</u>
Avascular Necrosis (AVN) of the Tarsal Navicular (Kohler Disease) or Metatarsal Head (Frieberg's Infraction)	Yes	No	➤ MRI foot without contrast (CPT® 73718) when suspected and plain x-ray is negative or equivocal or with AVN confirmed by plain x-ray if needed for treatment planning	See: <u>MS-4.1: AVN</u>
Tendonitis	Yes	Yes	➤ MRI foot without contrast (CPT® 73718) OR ➤ US foot (CPT® 76881 or 76882)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Complete rupture/tear of a specific named tendon	Yes	No	➤ For preoperative planning: ◆ MRI foot without contrast (CPT® 73718) OR ◆ US foot (CPT® 76881 or 76882)	
Partial Tendon Rupture	Yes	No	➤ For a suspected partial tendon rupture of a specific named tendon not otherwise specified: ◆ MRI foot without contrast (CPT® 73718) OR ◆ US foot (CPT® 76881 or 76882)	MRI is NOT needed for muscle belly strains/muscle tears.
Morton's Neuroma	Yes	Yes	➤ For preoperative planning: ◆ MRI foot without contrast (CPT® 73718) OR ◆ US foot (CPT® 76881 or 76882)	
Plantar Fasciitis	Yes	Yes*	➤ For preoperative planning: ◆ MRI foot without contrast (CPT® 73718) OR ◆ US foot (CPT® 76881 or 76882)	*Provider-directed conservative treatment must be for 6 months or more.
Suspected Plantar Fascia Rupture or Tear	Yes	Yes	➤ MRI foot without contrast (CPT® 73718) OR ➤ US foot (CPT® 76881 or 76882)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Diabetic Foot Infection	Yes*	No	➤ For suspected osteomyelitis or soft tissue infection as a complement to plain x-ray (both plain x-ray and MRI are indicated): ◆ MRI foot without and with contrast (CPT® 73720) OR ◆ MRI foot without contrast (CPT® 73718)	* Plain x-ray results do not preclude the necessity for advanced imaging studies. See also <u>MS 9.1: Infection-General</u>
Tarsal Tunnel Syndrome including Baxter's Neuropathy	Yes	Yes	➤ For preoperative planning if mass/lesion is suspected as etiology of entrapment: ◆ MRI foot without contrast (CPT® 73718) OR ◆ US foot (CPT® 76881 or 76882)	
Tarsal Coalition	Yes	Yes	➤ For preoperative planning: ◆ MRI ankle without contrast (CPT® 73721) OR ◆ CT without contrast (CPT® 73700)	
Sinus Tarsi Syndrome	Yes	Yes	➤ MRI ankle without contrast (CPT® 73721) if diagnosis is unclear or for preoperative evaluation	
Charcot Foot	Yes	Yes	➤ MRI foot without contrast (CPT® 73718) OR ➤ MRI foot without and with contrast (CPT® 73720)	
CRPS Type I	Yes	Yes	➤ Triple phase bone scan (CPT® 78315) OR ➤ MRI foot without contrast (CPT® 73718)	

Condition	Plain X-Ray?	Conservative Treatment	Advanced Imaging	Comments
Patient's condition	Are the results of an initial plain x-ray required before advanced imaging can be approved? (Yes or No)	Is failure of 6 weeks of provider-directed conservative treatment within the past 12 weeks with clinical re-evaluation required? (Yes or No)	The appropriate advanced imaging indicated for this condition. In some scenarios, advanced imaging may not be indicated.	Additional comments related to the condition.
Post-Operative	Yes	Yes	➤ In symptomatic patients following surgery for conditions including the tendons, ligaments and plantar plate, ONE of the following: ◆ MRI foot without contrast (CPT® 73718) ◆ US foot (CPT® 76881 or 76882) ➤ In symptomatic patients following surgery for complex fractures, sesamoid fractures and subtalar arthrodesis: ◆ CT foot without contrast (CPT® 73700)	

One Study/Area Only

In foot and ankle advanced imaging, studies are frequently ordered of both areas. This is unnecessary since ankle MRI will image from above the ankle to the mid- metatarsal area. **Only one CPT® code should be reported.**

References

1. Griffin LY. *Essentials of Musculoskeletal Care*. 3rd edition. Rosemont, IL: American Academy of Orthopaedic Surgeons; 2005:619-622;667-671;681-684;697-699;700-702.
2. Needell S, Cutler J. Morton neuroma imaging. eMedicine, April 11, 2011
3. Morton's Neuroma. MDGuidelines™.
4. Berquist TH. *Radiology of the Foot and Ankle*. 2nd Ed. Philadelphia, Lippincott, 2000, pp.155-156.
5. Bouche R. Sinus Tarsi Syndrome. What is Sinus Tarsi Syndrome, Testing and Treatment. http://www.aapsm.org/sinus_tarsi_syndrome.html.
6. D Resnick. *Internal Derangements of Joints 2006: Imaging-Arthroscopic Correlation*. Washington, DC.Oct.31-Nov. 4, 2006.
7. Doty JF, Coughlin MJ. Metatarsophalangeal joint instability of the lesser toes and plantar plate deficiency. *J Am Acad Orthop Surg*. 2014;22(4):235-245. doi:10.5435/JAAOS-22-04-235.
8. Lareau CR, Sawyer GA, Wang JH, et al. Plantar and medial heel pain: diagnosis and management. *J Am Acad Orthop Surg*. 2014;22:372-380. doi:10.5435/JAAOS-22-06-372.
9. Sung, W, Weil L Jr, Weill LS Sr, et al. Diagnosis of plantar plate injury by magnetic resonance imaging with reference to Intraoperative findings. *Journal of Foot Ankle Surgery*. 2012;51(5):570-574. doi:10.1053/j.jfas.2012.05.009.
10. Bancroft LW, Kransdorf MJ, Adler R, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Acute Trauma to the Foot. *Am Coll Radiol (ACR)*; Date of Origin: 2010. Last Review: 2019. <https://acsearch.acr.org/docs/70546/Narrative/>.
11. Wise JN, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
12. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.
13. Kransdorf MJ, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2019. <https://acsearch.acr.org/docs/69340/Narrative/>.
14. Thomas JL, Christensen JC, Kravitz SR, et al. The Diagnosis and Treatment of Heel Pain: A Clinical Practice Guideline - Revision 2010. *J Foot Ankle Surg*. 2010;49:S1-S19. doi: 10.1053/j.jfas.2010.01.001.
15. Goff JD, Crawford R. Diagnosis and Treatment of Plantar Fasciitis. *Am Fam Physician*. 2011 Sep;84(6):676-682. <https://www.aafp.org/afp/2011/0915/p676.html>.
16. Baxter D, Pfeffer G. Treatment of chronic heel pain by surgical release of the first branch of the lateral plantar nerve. *Clin Orthop*. 1992;279:229–236.
17. Schepsis A, Leach R, Gorzyca J. Plantar fasciitis: etiology, treatment, surgical results, and review of the literature. *Clin Orthop* 1991;266:185–196.
18. Neufeld SK, Cerato R. Plantar fasciitis: evaluation and treatment. *J Am Acad Orthop Surg*. 2008;16:338-46. doi:10.5435/00124635-200806000-00006.
19. Dodd A, Daniels TR. Charcot Neuroarthropathy of the Foot and Ankle. *J Bone Joint Surg Am*. 2018;100:696-711. doi:10.2106/JBJS.17.00785.

MS-28: Nuclear Medicine

- SPECT scan may be approved for any of the indications for which a bone scan can be approved.
 - ◆ If the request is for CPT® 78300 and CPT® 78803, then only CPT® 78803 is to be approved if medical necessity is established.
 - ◆ If the request is for CPT® 78305 or CPT® 78306 and CPT® 78803, then two CPT® codes may be approved if medical necessity is established.
- Nuclear Medicine may be used in the evaluation of some musculoskeletal disorders, and other rare indications exist as well.
 - ◆ Evaluation of suspected aseptic loosening of orthopedic prostheses when recent plain x-ray is nondiagnostic:
 - Bone scan (CPT® 78315) OR
 - Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803, or 78831) OR
 - Hybrid SPECT/CT (CPT® 78830)
 - See: **MS-16: Post-Operative Joint Replacement Surgery** and anatomic tables
 - ◆ For detection of ischemic or infarcted regions in sickle cell disease:
 - Nuclear medicine bone marrow imaging (CPT® 78102, 78103, 78104) OR
 - SPECT (CPT® 78803) OR
 - Hybrid SPECT/CT (CPT® 78830)
 - See also: **PEDMS-1.3: Modality General Considerations**
 - ◆ Evaluation of complex regional pain syndrome or reflex sympathetic dystrophy
 - Triple phase bone scan (CPT® 78315)
 - For interventional pain criteria see: **CMM-209: Regional Sympathetic Blocks** and **CMM-211: Spinal Cord Stimulators**
 - ◆ Evaluation of Paget's disease
 - Bone scan (CPT® codes: 78300, 78305, 78306) OR
 - Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803) OR
 - Hybrid SPECT/CT (CPT® 78830)
 - See: **MS-10: Soft Tissue Mass or Lesion of Bone**
 - ◆ Suspected fractures, if MRI cannot be performed
 - Tc-99m bone scan whole body (CPT® 78306) with SPECT of the area of interest (CPT® 78803) OR
 - Hybrid SPECT/CT (CPT® 78830) OR
 - Bone scan (CPT® 78315, 78305, or 78300)
 - See: **MS-5.2: Suspected Occult/Stress/Insufficiency Fracture/Stress Reaction and Shin Splints.**
 - ◆ Evaluation of suspected bone infection if MRI cannot be done and when infection is multifocal, or when the infection is associated with orthopedic hardware or chronic bone alterations from trauma or surgery
 - Bone scan (CPT® 78315, 78300, 78305, or 78306) OR
 - Distribution Of Radiopharmaceutical Agent SPECT (CPT® 78803 or 78831) OR
 - Hybrid SPECT/CT (CPT® 78830 or 78832)

- Combining bone scintigraphy with a labeled leukocyte scan enhances sensitivity. A labeled leukocyte scan (radiopharmaceutical inflammatory imaging - one of CPT® codes: 78800, 78801, 78802, or 78803) in concert with Tc-99m sulfur colloid marrow imaging (one of CPT® codes: 78102, 78103, or 78104) or Hybrid SPECT/CT (CPT® 78830) is particularly useful in cases with altered bone marrow distribution, such as joint prosthesis.
- See: **MS-16: Post-Operative Joint Replacement Surgery**
- For specific joints post-operative from replacement surgery:
 - See: **MS-19: Shoulder**
 - See: **MS-20: Elbow**
 - See: **MS-24: Hip**
 - See: **MS-25: Knee**
 - See: **MS-26: Ankle**

References

1. Beaman FD, von Herrmann PF, Kransdorf MJ, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *Am Coll Radiol (ACR)*; Date of Origin: 2016. <https://acsearch.acr.org/docs/3094201/Narrative/>.
2. Kransdorf MJ, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Suspected Osteomyelitis of the Foot in Patients with Diabetes Mellitus. *Am Coll Radiol (ACR)*; Date of Origin: 1995. Last Review: 2019. <https://acsearch.acr.org/docs/69340/Narrative/>.
3. Wise JN, Weissman BN, Appel M, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Chronic Foot Pain. *Am Coll Radiol (ACR)*; Date of Origin: 1998. Last Review: 2020. <https://acsearch.acr.org/docs/69424/Narrative/>.
4. Bencardino JT, Stone TJ, Roberts CC, et. al. Expert Panel on Musculoskeletal Imaging. ACR Appropriateness Criteria® Stress (Fatigue/Insufficiency) Fracture, Including Sacrum, Excluding Other Vertebrae. *Am Coll Radiol (ACR)*; Revised: 2016. <https://acsearch.acr.org/docs/69435/Narrative/>.