

Embarc Benefit Protection Policy

POLICY: Spinal Muscular Atrophy – Zolgensma® (onasemnogene abeparvovec-xioi intravenous infusion – Novartis)

EFFECTIVE DATE: November 17, 2023

VERSION: V2.0.2023

Overview

Zolgensma, an adeno-associated virus vector-based gene therapy, is indicated for the treatment of pediatric patients less than 2 years of age with spinal muscular atrophy with bi-allelic mutations in the survival motor neuron 1 (SMN1) gene.¹

Limitations of use are that the safety and effectiveness of repeat administration of Zolgensma have not been evaluated.¹ The use of Zolgensma in patients with advanced spinal muscular atrophy (e.g., complete paralysis of limbs, permanent ventilator dependence) has not been assessed. Use of Zolgensma in premature neonates before reaching full-term gestational age is not recommended because concomitant treatment with corticosteroids may adversely affect neurological development. Zolgensma therapy should be delayed until full-term gestational age is achieved. The definition of full-term pregnancy commences at 39 weeks and 0 days gestation.²

Disease Overview

Spinal muscular atrophy is a genetic, autosomal recessive muscular disorder caused by deletion or loss of function mutation in the SMN1 gene.³⁻⁶ The reduced levels of survival motor neuron (SMN) protein causes degeneration of lower motor neurons.⁶ Although the condition is a multisystem disorder, it is clinically characterized by progressive muscle weakness and atrophy. Patients have difficulties with ambulation, head control, feeding and respiration. Cognitive development is not impacted. In the US, spinal muscular atrophy affects approximately one in 11,000 infants and has an average carrier frequency of one in 54 individuals; as many as 10,000 to 20,000 children and adults in the US may be impacted.⁶ Although the condition can be present in individuals of any age, it is more frequently diagnosed in infants and children, as it is more severe in this population.³⁻⁶ The phenotypic expression of the disease is impacted by the presence of the survival motor neuron 2 (SMN2) gene copy number. SMN1 is responsible for producing most of the effective SMN protein, although some SMN protein can be made by the SMN2 gene. Therefore, patients with a deletion of the SMN1 gene may have the potential for making some SMN protein through the SMN2 gene copy, although in most cases the resulting protein made by this gene is truncated and is not as effective or functional. Data have shown that patients with a higher number of SMN2 gene copies generally have a more mild phenotypic disease expression. Gene deletion testing for spinal muscular atrophy can be performed at many diagnostic laboratories. Table 1 describes disease types. A different manner of categorization classifies the three most common types as follows: Type 1 patients are “non-sitters”, Type 2 patients are “sitters”, and Type 3 patients are “walkers”.^{4,6}

Table 1. Types of Spinal Muscular Atrophy.³⁻⁶

SMA Type	Age at Onset	Features/Clinical Presentation	Lifespan	SMN2 Gene Copy Number
0	Prenatal	Severe hypotonia and weakness; respiratory failure at birth. There is no achievement of motor milestones.	A few weeks to < 6 months	0 to 1
1	< 6 months	Poor muscle tone, lack of movement, and respiratory assistance needed at birth. Patients are never able to sit.	< 2 years	1 to 2
2	Before 18 months	Patients are able to sit. However, patients are unable to walk or stand without assistance.	75% of patients are alive at 25 years of age	2 to 3
3	> 18 months	Walks independently, but may lose this ability as the disease progresses.	Normal	3 to 4
4	Adulthood	Walk until adulthood.	Normal	≥ 4

SMA – Spinal muscular atrophy; SMN2 – Survival motor neuron 2.

Besides Zolgensma, other therapies are available. **Spinraza**® (nusinersen intrathecal injection), a SMN2-directed antisense oligonucleotide, is indicated for the treatment of spinal muscular atrophy in pediatric and adult patients.⁷ Spinraza is given by intrathecal injection. Although studies and experience continue, the primary pivotal data include infantile-onset (Type 1) and later-onset (Type 2 and Type 3) spinal muscular atrophy primarily in children. Data are also available with Spinraza in presymptomatic infants who were genetically diagnosed with spinal muscular atrophy. There is an accumulation of data with Spinraza in adults as well.

Evrysdi® (risdiplam oral solution), a SMN2 splicing modifier, is indicated for the treatment of spinal muscular atrophy in pediatric and adult patients.⁸ The primary pivotal data include infantile-onset (Type 1) and later-onset (Type 2 and Type 3) spinal muscular atrophy primarily in children and adults up to 25 years of age. Trials are ongoing in older adults, as well as in presymptomatic infants who were genetically diagnosed with spinal muscular atrophy.

Clinical Efficacy

The efficacy of Zolgensma was evaluated in patients less than 2 years of age with spinal muscular atrophy who had bi-allelic mutations in the SMN1 gene.^{1,9-14} One trial was an open-label, single-arm study which is ongoing (STR1VE [n = 21])¹¹ and the other was an open-label, single-arm, ascending-dose clinical trial (START [n = 15] {12 patients received a therapeutic dose}).^{1,9-10} Symptoms onset occurred before patients were 6 months of age. All patients had genetically confirmed bi-allelic SMN1 gene deletions and two SMN2 gene copies. In both trials, Zolgensma was given as a single-dose intravenous infusion. Efficacy was assessed on parameters such as survival and achievement of developmental motor milestones (e.g., sitting without support). The definition of survival was the time from birth to either death or permanent ventilation. Other efficacy parameters were evaluated (e.g., assessment of Children's Hospital of Philadelphia Infant Test of Neuromuscular Disorders [CHOP-INTEND] scores, evaluation of ventilator use). The ongoing clinical trial involved 21 patients with infantile-onset spinal muscular atrophy. The mean CHOP-INTEND score was 31.0 (range 18 to 47). The mean patient age at the time of treatment was 3.9 months (range 0.5 to 5.9 months). As of the March 2019 cutoff date, 19 patients were alive without permanent ventilation. Compared with natural history data, Zolgensma is effective as more patients attained the ability to sit without support.¹

The completed clinical trial involved 15 patients with infantile-onset spinal muscular atrophy.^{1,9} Three patients were in a low-dose cohort and 12 patients were in a high-dose cohort.¹ At the time of treatment, the mean age of patients in the low-dose cohort was 6.3 months (range 5.9 to 7.2 months) and 3.4 months (range 0.9 to 7.9 months) in the high-dose group. The dose in the low-dose cohort was approximately one-third of the dosage received by patients in the high-dose cohort. At 24 months following Zolgensma infusion, one patient in the low-dose cohort met the endpoint of permanent ventilation; all 12 patients in the high-dose cohort were alive without permanent ventilation. In the high-dose cohort, 9 of 12 patients (75%) were able to stand and walk without assistance.^{1,9} At longer-term follow-up from the START trial, all 10 patients followed in the high-dose group were alive without permanent ventilation at the dataset on June 11, 2020. In STRIVE, at the March 2019 data cutoff, 19 patients were alive without permanent ventilation.¹ Up until November 2019, data revealed that 13 of 22 patients achieved the coprimary endpoint of functional independent sitting for 30 seconds or longer at the 18 months of age study visit.¹¹ Other data are also available.¹²⁻¹⁵

Guidelines

The Spinal Muscular Atrophy Newborn Screening Multidisciplinary Working Group is comprised of clinicians and geneticists with expertise in spinal muscular atrophy who developed a treatment algorithm in 2018 for infants who have positive results from a newborn screening test for spinal muscular atrophy.¹⁶ Spinal muscular atrophy Types 1 and 2 comprise a large majority of cases and account for many patients who screen positively for spinal muscular atrophy with three or fewer SMN2 gene copies. Immediate treatment is recommended in patients with two or three SMN2 gene copies. Treatment recommendations for patients who screen positive for spinal muscular atrophy and have only one SMN2 gene copy are more complicated. It is likely that patients with only one SMN2 gene copy will likely be symptomatic at birth and the physician should determine if treatment is warranted.¹⁶ In 2020, the Working Group updated recommendations that infants diagnosed with spinal muscular atrophy via newborn screening with four SMN2 gene copies should receive immediate treatment.¹⁷ Also, patients with five (or more) SMN2 gene copies should be observed and screened for symptoms.

Dosing

The recommended dose of Zolgensma is 1.1×10^{14} vector genomes (vg) per kg of body weight.¹ Administer Zolgensma as an intravenous infusion over 60 minutes. Starting 1 day prior to Zolgensma infusion, give systemic corticosteroids equivalent to oral prednisolone 1 mg/kg of body weight for a total of 30 days. Examine liver function after this juncture and follow recommended guidelines.

Safety

Zolgensma has a Boxed Warning regarding acute serious liver injury and acute liver failure.¹ Elevated aminotransferases can occur with Zolgensma. Patients with preexisting liver impairment may be at higher risk. Prior to infusion, evaluate liver function in all patients by clinical examination and laboratory testing. One day before Zolgensma infusion, commence administration of systemic corticosteroids equivalent to oral prednisolone at 1 mg/kg of body weight per day for a total of 30 days. Prior to administration of Zolgensma, evaluate creatinine and complete blood counts. Perform baseline anti-AAV9 antibody testing prior to Zolgensma infusion. Patients in the Zolgensma trials were required to have baseline anti-AAV9 antibody titers of $\leq 1:50$.

POLICY STATEMENT

Prior Authorization is recommended for benefit coverage of Zolgensma. Approval is recommended for those who meet the **Criteria** and **Dosing** for the listed indication. Because of the specialized skills required for evaluation and diagnosis of patients treated with Zolgensma as well as the specialized training required for administration of Zolgensma, approval requires Zolgensma to be prescribed by a physician who has consulted with or who specializes in the condition. All approvals are provided for one dose per lifetime. The approval duration is 1 month to allow for an adequate timeframe to prepare and administer one dose of therapy. For certain criteria, verification is required as noted by **[verification in claims history required]**. All reviews (approvals and denials) will be forwarded to the Medical Director for evaluation.

DOCUMENTATION

Documentation is required where noted in the criteria as **[documentation required]**. Documentation may include, but is not limited to, chart notes, laboratory tests, claims records, and/or other information.

Automation

None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Zolgensma is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Spinal Muscular Atrophy - Treatment.** Approve for a one-time per lifetime dose if the patient meets the following (A, B, C, D, E, F, G, H, I, J, K, L, M, and N):
 - A) Patient is less than 2 years of age; AND
 - B) If the patient is a premature neonate, full-term gestation age of 39 weeks and 0 days has been met; AND
Note: Full-term gestational age can be defined as the postmenstrual age (gestational age plus chronological age) being equal to ≥ 39 weeks and 0 days.
 - C) Patient has had a genetic test confirming the diagnosis of spinal muscular atrophy with bi-allelic pathogenic variants in the survival motor neuron 1 (SMN1) gene **[documentation required]**; AND
Note: Pathogenic variants may include homozygous deletion, compound heterozygous mutation, or a variety of other rare mutations.
 - D) Patient meets one of the following (i or ii):
 - i. Patient has three or fewer survival motor neuron 2 (SMN2) gene copies **[documentation required]**; OR
 - ii. Patient meets both of the following (a and b):
 - a) Patient has four survival motor neuron 2 (SMN2) gene copies **[documentation required]**;
AND
 - b) The number of survival motor neuron 2 (SMN2) gene copies has been determined by a quantitative assay capable of distinguishing between four SMN2 gene copies and five or greater SMN2 gene copies; AND
 - E) According to the prescribing physician, patient has started or will receive systemic corticosteroids equivalent to oral prednisolone at a dose of 1 mg/kg per day commencing 1 day prior to Zolgensma infusion and for a total of 30 days; AND
 - F) Baseline anti-AAV9 antibody titers are $\leq 1:50$ **[documentation required]**; AND

- G)** Patient has undergone a liver function assessment within the last 30 days and meets all of the following(i, ii, iii, and iv):
- i. Alanine aminotransferase levels are ≤ 2 times the upper limit of normal **[documentation required]**; AND
 - ii. Aspartate aminotransferase levels are ≤ 2 times the upper limit of normal **[documentation required]**; AND
 - iii. Total bilirubin levels are ≤ 2 times the upper limit of normal **[documentation required]**; AND
Note: Patient with elevated bilirubin levels due to neonatal jaundice are acceptable.
 - iv. Prothrombin time results are ≤ 2 times the upper limit of normal **[documentation required]**; AND
- H)** Patient has undergone a renal function assessment within the last 30 days and has a creatinine level < 1.0 mg/dL **[documentation required]**; AND
- I)** A complete blood count has been obtained within the last 30 days and the patient meets both of the following (i and ii):
- i. White blood cell count is $\leq 20,000$ cells per mm^3 **[documentation required]**; AND
 - ii. Hemoglobin levels are between 8 g/dL and 18 g/dL **[documentation required]**; AND
- J)** Patient has not received Zolgensma in the past **[verification in claims history required]**; AND
Note: Verify through claims history that the patient has not previously received Zolgensma. If no claim for Zolgensma is present, the prescribing physician confirms that the patient has not previously received Zolgensma.
- K)** For a patient currently receiving or who has received prior treatment with Spinraza (nusinersen intrathecal injection), the prescribing physician confirms that further therapy with Spinraza will be discontinued; AND
- L)** For a patient currently receiving or who has received prior treatment with Evrysdi (risdiplam oral solution), the prescribing physician confirms that further therapy with Evrysdi will be discontinued; AND
- M)** Medication is prescribed by a physician who has consulted with or who specializes in the management of patients with spinal muscular atrophy and/or neuromuscular disorders; AND
- N)** If criteria A through M are met, approve one single intravenous infusion of Zolgensma at a dose of 1.1×10^{14} vector genomes per kg (vg/kg) based on the current patient weight in kg (within the last 14 days) **[documentation required]**. Zolgensma is provided as a customized kit to meet dosing requirements for each patient per their documented weight (in kilograms). Configuration of the dose kit is based on weight (per the cited NDC) as in Table 2 below.

Dosing. The recommended dose of Zolgensma for single-dose intravenous infusion is 1.1×10^{14} vector genomes (vg)/kg based on the current patient weight in kg (within the last 14 days). Zolgensma is provided as a customized kit to meet dosing requirements for each patient per their documented weight (in kilograms). Refer to the appropriate NDC number below for approval.

Table 2. Dose of Zolgensma Based on Availability.¹

Patient Weight Range (kg)	Dose Volume (mL)*	Zolgensma Kit Configuration			NDC Number
		5.5 mL vial	8.3 mL vial	Total Vials per Kit	
2.6 to 3.0	16.5	0	2	2	71894-120-02
3.1 to 3.5	19.3	2	1	3	71894-121-03
3.6 to 4.0	22.0	1	2	3	71894-122-03
4.1 to 4.5	24.8	0	3	3	71894-123-03
4.6 to 5.0	27.5	2	2	4	71894-124-04
5.1 to 5.5	30.3	1	3	4	71894-125-04
5.6 to 6.0	33.0	0	4	4	71894-126-04
6.1 to 6.5	35.8	2	3	5	71894-127-05
6.6 to 7.0	38.5	1	4	5	71894-128-05
7.1 to 7.5	41.3	0	5	5	71894-129-05
7.6 to 8.0	44.0	2	4	6	71894-130-06
8.1 to 8.5	46.8	1	5	6	71894-131-06
8.6 to 9.0	49.5	0	6	6	71894-132-06
9.1 to 9.5	52.3	2	5	7	71894-133-07
9.6 to 10.0	55.0	1	6	7	71894-134-07
10.1 to 10.5	57.8	0	7	7	71894-135-07
10.6 to 11.0	60.5	2	6	8	71894-136-08
11.1 to 11.5	63.3	1	7	8	71894-137-08
11.6 to 12.0	66.0	0	8	8	71894-138-08
12.1 to 12.5	68.8	2	7	9	71894-139-09
12.6 to 13.0	71.5	1	8	9	71894-140-09
13.1 to 13.5	74.3	0	9	9	71894-141-09
13.6 to 14.0	77.0	2	8	10	71894-142-10
14.1 to 14.5	79.8	1	9	10	71894-143-10
14.6 to 15.0	82.5	0	10	10	71894-144-10
15.1 to 15.5	85.3	2	9	11	71894-145-11
15.6 to 16.0	88.0	1	10	11	71894-146-11
16.1 to 16.5	90.8	0	11	11	71894-147-11
16.6 to 17.0	93.5	2	10	12	71894-148-12
17.1 to 17.5	96.3	1	11	12	71894-149-12
17.6 to 18.0	99.0	0	12	12	71894-150-12
18.1 to 18.5	101.8	2	11	13	71894-151-13
18.6 to 19.0	104.5	1	12	13	71894-152-13
19.1 to 19.5	107.3	0	13	13	71894-153-13
19.6 to 20.0	110.0	2	12	14	71894-154-14
20.1 to 20.5	112.8	1	13	14	71894-155-14
20.6 to 21.0	115.5	0	14	14	71894-156-14

* Dose volume is calculated using the upper limit of the patient weight range for pediatric patients less than 2 years of age between 2.6 kg and 21.0 kg.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Zolgensma is not recommended in the following situations:

- 1. Patient has Complete Paralysis of All Limbs.** This is cited as a limitation of use in the Zolgensma prescribing information.¹ Data are needed to determine if this patient population would derive benefits from Zolgensma.
- 2. Patient has Permanent Ventilator Dependence.** This is cited as a limitation of use in the Zolgensma prescribing information.¹ Data are needed to determine if this patient population would derive benefits from Zolgensma.
- 3. Administration in Individuals in Utero.** Zolgensma is not approved for in utero administration per the prescribing information.
- 4.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Zolgensma® intravenous infusion [prescribing information]. Bannockburn, IL: Novartis; October 2023.
2. ACOG Committee Opinion No 579: Definition of term pregnancy. *Obstet Gynecol.* 2013;122(5):1139-1140.
3. Arnold ES, Fischbeck KH. Spinal muscular atrophy. *Handb Clin Neurol.* 2018;148:591-601.
4. Prior TW, Leach ME, Finanger E. Spinal Muscular Atrophy. 2000 Feb 24 [Updated 2020 December 3]. In: Adam MP, Ardinger, HH, Pagon RA, et al, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2020. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK1352/>. Accessed on November 1, 2023.
5. Nicolau S, Waldrop MA, Connolly AM, Mendell JR. Spinal muscular atrophy. *Semin Pediatr Neurol.* 2021;37:100878.
6. Yeo CJJ, Darras BT. Overturning the paradigm of spinal muscular atrophy as just a motor neuron disease. *Pediatr Neurol.* 2020;109:12-19.
7. Spinraza® intrathecal injection [prescribing information]. Cambridge, MA: Biogen; February 2023.
8. Evrysdi® oral solution [prescribing information]. South San Francisco, CA: Genentech/Roche; October 2023.
9. Mendell JR, Al-Zaidy S, Shell R, et al. Single-dose gene replacement therapy for spinal muscular atrophy. *N Engl J Med.* 2017;377(18):1713-1722.
10. Mendell JR, Al-Zaidy SA, Lehman KJ, et al. Five-year extension results of the Phase I START trial of onasemnogene abeparvovec in spinal muscular atrophy. *JAMA Neurol.* 2021;78(7):834-841.
11. Day JW, Finkel RS, Chiriboga CA, et al. Onasemnogene abeparvovec gene therapy for symptomatic infantile-onset muscular atrophy in patients with two copies of SMN2 (STR1VE): an open-label, single-arm, multicenter, phase 3 trial. *Lancet Neurol.* 2021;20:284-293.
12. Strauss KA, Farrar MS, Muntoni F, et al. Onasemnogene abeparvovec for presymptomatic infants with three copies of SMN2 at risk for spinal muscular atrophy: the Phase III SPR1NT trial. *Nat Med.* 2022;28:1390-1397.
13. Strauss KA, Farrar MS, Muntoni F, et al. Onasemnogene abeparvovec for presymptomatic infants with two copies of SMN2 at risk for spinal muscular atrophy type 1: the Phase III SPR1NT trial. *Nat Med.* 2022;28:1381-1389.
14. Mercuri E, Muntoni F, Baranello G, et al. Onasemnogene abeparvovec gene therapy for symptomatic infantile-onset spinal muscular atrophy type 2 (STR1VE-EU): an open-label, single-arm, multicenter, phase 3 trial. *Lancet Neurol.* 2021;20:832-841.
15. Blair HA. Onasemnogene abeparvovec: a review of spinal muscular atrophy. *CNS Drugs.* 2022;36:995-1005.
16. Glascock J, Sampson J, Haidet-Phillips A, et al. Treatment algorithm for infants diagnosed with spinal muscular atrophy through newborn screening. *J Neuromuscul Dis.* 2018;5:145-158.
17. Glascock J, Sampson J, Connolly AM, et al. Revised recommendations for the treatment of infants diagnosed with spinal muscular atrophy via newborn screening who have 4 copies of SMN2. *J Neuromuscul Dis.* 2020;7(2):97-100.