

Evidence Discussion

(AB-10.1) **EviCore**
By EVERNORTH

- Intra-abdominal injury is an indication for ultrasound (US) and/or advanced imaging. Advanced imaging in the acute situation is generally with CT of the Abdomen and/or Pelvis with contrast. Both US and CT can be completed rapidly. CT with contrast can be more diagnostic for bleeding and/or soft tissue inflammation/injury.

References

1. ACR Appropriateness Criteria® blunt abdominal trauma Clinical Practice Guidelines. Guideline Central.
2. Soto JA and Anderson SW. Multidetector CT of Blunt Abdominal Trauma. Radiology. 2012;265(3):678-693.
3. Nishijima DK, Simel DL, Wisner DH, et al. Does this adult patient have a blunt intra-abdominal injury? JAMA 2012; 307:1517.
4. Washington State Department of Health Office of Community Health Systems: Trauma Clinical Guideline. May 2017. <https://www.doh.wa.gov/Portals/1/Documents/Pubs/530168.pdf>.
5. Jansen JO, Yule SR, Loudon MA. Investigation of blunt abdominal trauma. Bmj. 2008;336(7650):938-942. doi:10.1136/bmj.39534.686192.80.
6. Diercks DB, Mehrotra A, Nazarian DJ, Promes SB, Decker WW, Fesmire FM. Clinical Policy: Critical Issues in the Evaluation of Adult Patients Presenting to the Emergency Department With Acute Blunt Abdominal Trauma. Annals of Emergency Medicine. 2011;57(4):387-404. doi:10.1016/j.annemergmed.2011.01.013.