

Apokyn® (apomorphine)

When requesting Apokyn® (apomorphine), the individual requiring treatment must be diagnosed with an FDA-approved indication and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-Approved Indication

Apokyn® (apomorphine) is indicated for the acute, intermittent treatment of hypomobility, “off” episodes (“end-of-dose wearing off” and unpredictable “on/off” episodes) associated with advanced Parkinson’s disease.

Coverage Guidelines

Parkinson’s Disease

The individual must meet **all** of the following criteria for approval:

- Is experiencing “off” episodes such as muscle stiffness, slow movements, or difficulty starting movements;
- Is currently receiving carbidopa/levodopa therapy;
- Has had significant intolerance OR inadequate efficacy to one other treatment for “off” episodes (e.g., entacapone, rasagiline, pramipexole, ropinirole, tolcapone, cabergoline, selegiline, Kynmobi, Ongentys, Xadago);
- Apokyn is prescribed by or in consultation with a neurologist.

Approval duration: 12 months

Dosing Recommendation

Apokyn is indicated for subcutaneous administration only.

The recommended starting dose of APOKYN is 0.1 mL (1 mg) to 0.2 mL (2 mg). Titrate on the basis of effectiveness and tolerance, up to a maximum recommended dose of 0.6 mL (6 mg)

References

1. Apokyn® subcutaneous injection [prescribing information] Louisville, KY: US WorldMeds; June 2022.
2. Pahwa R, Factor SA, Lyons KE, et al. Practice parameter: Treatment of Parkinson disease with motor fluctuations and dyskinesia (an evidence-based review). Report of the quality standards subcommittee of the American Academy of Neurology. Neurology. 2006;66:983-995.