

Evidence Discussion (CD 5.2, 5.4-5.7)

- Guidelines and appropriate use criteria support the use of transthoracic echocardiogram (TTE) as the initial study for a broad range of cardiac conditions. TTE can be used to evaluate cardiac morphology and function, and provides information necessary to diagnose and guide treatment in conditions including heart failure and cardiomyopathy, ischemic heart disease, and valvular dysfunction. TTE is also used to initially assess intracardiac and extracardiac structures, including suspected cardiac masses, sources of emboli, and pericardial conditions. TEE visualizes portions of the great vessels and provides initial imaging of these structures.
- TTE has significant advantages over other imaging modalities: it is easily accessible and even portable, provides no exposure to radiation, is inexpensive, and has minimal, if any risks.
- Like TTE, MRI assesses cardiac morphology and function; additionally, MRI provides information regarding myocardial perfusion, metabolism, and tissue composition.
- When questions remain unanswered after TTE, MRI can provide additional information due to its ability to assess myocardial composition and mass; these characteristics make cardiac MRI especially useful in the settings of cardiomyopathy, non-compaction, infiltrative heart disease, post cardiac transplant, cardiac masses, pericardial disease, and myocarditis.
- Beyond imaging with TTE, MRI can provide improved visualization of valvular pathology, because images are not limited by body habitus, is of high spatial resolution, can more accurately quantify the magnitude of valve insufficiency and stenosis, and can assess the effects of valve dysfunction on ventricular mass or volume. Qualities of MRI that augment evaluation of valvular disease may also improve visualization of intracardiac shunts over TTE alone.
- Advantages of MRI assessment of the great vessels include avoidance of ionizing radiation, provision of excellent spatial resolution of structures, and is not limited by body habitus.
- Conditions that are better evaluated initially by MRI, include coronary artery anomalies, and diseases of abnormal myocardial composition: Arrhythmogenic right ventricular dysplasia, Fabry disease, cardiac sarcoidosis, and Duchenne Muscular Dystrophy.
- When TTE and other less advanced imaging provide the information necessary to diagnose and treat cardiovascular conditions, MRI is not indicated.
- Disadvantages of MRI include being less accessible, and possibly being incompatible for use in those with implanted devices. The gadolinium contrast agents used in MRI require precautions in those with advanced renal disease and severe liver disease.
- The decision to use MR imaging is made in the context of other testing; duplication of information should be avoided, and more readily available modalities with lower risk, initially considered. An assessment should be made as to whether additional testing will provide complementary diagnostic, therapeutic, or prognostic information that will optimize care.

REFERENCES

1. Hendel RC, Kramer CM, Patel MR, et al. ACCF/ACR/SCCT/SCMR/ ASNC/NASCI/SCAI/SIR 2006 Appropriateness Criteria for computed tomography and cardiac magnetic resonance imaging. *J Am Coll Cardiol*,2006; 48(7):1475-1497. Accessed November 30, 2017. doi:10.1016/j.jacc.2006.07.003
2. Hundley WG, Bluemke DA, Finn JP, et al. ACCF/ACR/AHA/NASCI/SCMR 2010 Expert Consensus Document on Cardiovascular Magnetic Resonance. *J Am Coll Cardiol*. 2010;55(23):2614-2662. doi:10.1016/j.jacc.2009.11.011.
3. Doherty JU, Kort S, Mehran R, et al. ACC/AATS/AHA/ASE/ASNC/HRS/SCAI/SCCT/SCMR/STS 2019 Appropriate Use Criteria for Multimodality Imaging in the Assessment of Cardiac Structure and Function in Nonvalvular Heart Disease. *J Am Coll Cardiol*. 2019;73(4):488-516. doi:10.1016/j.jacc.2018.10.038.
4. Pennell DJ, Sechtem UP, Higgins CB, et al. Clinical indications for cardiovascular magnetic resonance (CMR): Consensus Panel report. *Eur Heart J* 2004;25:1940–65.
5. Douglas PS, Garcia MJ, Haines DE, et al. ACCF/ASE/AHA/ASNC/ HFSA/HRS/SCAI/SCCM/SCCT/SCMR 2011 appropriate use criteria for echocardiography: a report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, and Society for Cardiovascular Magnetic Resonance. *J Am Coll Cardiol* 2011;57:1126-66.
6. Otto CM, Nishimura RA, Bonow RO, Carabello BA, Erwin JP 3rd, Gentile F, Jneid H, Krieger EV, Mack M, McLeod C, O’Gara PT, Rigolin VH, Sundt TM 3rd, Thompson A, Toly C. 2020 ACC/AHA guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2021;143:e72-e227. doi: 10.1161/CIR.0000000000000923
7. Doherty JU, Kort S, Mehran R, et al. ACC/AATS/AHA/ASE/ASNC/ HRS/SCAI/SCCT/SCMR/STS 2017 appropriate use criteria for multimodality imaging in valvular heart disease: a report of the American College of Cardiology Appropriate Use Criteria Task Force, American Association for Thoracic Surgery, American Heart Association, American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Cardiovascular Computed Tomography, Society for Cardiovascular Magnetic Resonance, and Society of Thoracic Surgeons. *J Am Coll Cardiol* 2017;70:1647-72.
8. Nagueh SF, Phelan D, Abraham T, Armour A, Desai MY, Dragulescu A, Gilliland Y, Lester SJ, Maldonado Y, Mohiddin S, Nieman K, Sperry BW, Woo A. Recommendations for Multimodality Cardiovascular Imaging of Patients with Hypertrophic Cardiomyopathy: An Update from the American Society of Echocardiography, in Collaboration with the American Society of Nuclear Cardiology, the Society for Cardiovascular Magnetic Resonance, and the Society of Cardiovascular Computed Tomography. *J Am Soc Echocardiogr*. 2022 Jun;35(6):533-569. doi: 10.1016/j.echo.2022.03.012. PMID: 35659037.
9. Ommen SR, Mital S, Burke MA, Day SM, Deswal A, Elliott P, Evanovich LL, Hung J, Joglar JA, Kantor P, Kimmelstiel C, Kittleson M, Link MS, Maron MS, Martinez MW, Miyake CY, Schaff HV, Semsarian C, Sorajja P. 2020 AHA/ACC Guideline for the Diagnosis and Treatment of Patients With Hypertrophic Cardiomyopathy: Executive Summary: A Report of

the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. J Am Coll Cardiol. 2020 Dec 22;76(25):3022-3055. doi: 10.1016/j.jacc.2020.08.044. Epub 2020 Nov 20.

10. Gersh BJ, Maron BJ, Bonow RO, et al. 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2011;58:e212-60.
11. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol 2013;62:e147-239.
12. Silvestry FE, Cohen MS, Armsby LB, et al. Guidelines for the echocardiographic assessment of atrial septal defect and patent foramen ovale: from the American Society of Echocardiography and Society for Cardiac Angiography and Interventions. J Am Soc Echocardiogr 2015;28:910-58.
13. Power LC, O'Grady GL, Hornung TS, Jefferies C, Gusso S, Hofman PL. Imaging the heart to detect cardiomyopathy in Duchenne muscular dystrophy: A review. Neuromuscular Disorders. 2018;28(9):717- 730. doi:10.1016/j.nmd.2018.05.011.
14. Riele AST, Tandri H, Sanborn DM, Bluemke DA. Noninvasive Multimodality Imaging in ARVD/C. JACC: Cardiovasc Imaging. 2015;8(5):597-611. doi:10.1016/j.jcmg.2015.02.007.
15. Ammirati E, Frigerio M, Adler ED, et al. Management of Acute Myocarditis and Chronic Inflammatory Cardiomyopathy: An Expert Consensus Document. Circ Heart Fail. 2020;13(11):e007405. doi:10.1161/CIRCHEARTFAILURE.120.007405.
16. Friedrich MG, Sechtem U, Schulz-Menger J, et al. Cardiovascular Magnetic Resonance in Myocarditis: A JACC White Paper. J Am Coll Cardiol. 2009;53(17):1475-1487. doi:10.1016/j.jacc.2009.02.007.
17. Nadel J, Lancefield T, Voskoboinik A, Taylor AJ. Late gadolinium enhancement identified with cardiac magnetic resonance imaging in sarcoidosis patients is associated with long-term ventricular arrhythmia and sudden cardiac death. Eur Heart J Cardiovasc Imaging 2015;16:634-41.
18. Fontana M, Pica S, Reant P, et al. Prognostic value of late gadolinium enhancement cardiovascular magnetic resonance in cardiac amyloidosis. Circulation 2015;132:1570-9.
19. Hiratzka LF, Bakris GL, Beckman JA, et al. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with thoracic aortic disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, American Association for Thoracic Surgery, American College of Radiology, American Stroke Association, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society of Interventional Radiology, Society of Thoracic Surgeons, and Society for Vascular Medicine. J Am Coll Cardiol 2010;55:e27-129.