

# Evidence Discussion Renal Cell Cancer (ONC-17)

---

## **Initial Staging**

American Urological Association (AUA) and the National Comprehensive Cancer Network (NCCN) guidelines state that pre- and post-contrast enhanced axial imaging, either CT or MRI, is the ideal imaging technique for the diagnosis and staging of clinically localized renal masses. Contrast-enhanced abdominal imaging best characterizes the mass, provides information regarding both the affected and unaffected renal unit, can assess extrarenal tumor spread (venous invasion or regional lymph nodes), and evaluates the adrenal glands and other abdominal organs for visceral metastasis.

Masses initially diagnosed by ultrasound should be confirmed with pre- and post-contrast enhanced imaging.

In patients with RCC or suspicion of RCC, complete staging is typically completed with chest radiography (CXR) or chest CT. Chest CT scan should be obtained selectively, in patients with pulmonary symptoms or abnormal CXR, or for patients with high-risk disease.(5,6) Bone scans should be reserved primarily for patients with bone pain or elevated alkaline phosphatase. Brain imaging is reserved for patients with neurologic symptoms.(8-10)

## **Restaging/Recurrence**

Add supporting statements for use of CT imaging for restaging

## **Surveillance**

### Active surveillance Stage T1a

Active surveillance entails serial abdominal imaging in order to study the growth rate of the tumor and it is recommended that abdominal imaging (CT or MRI with contrast) within 6 months from initiation of active surveillance; subsequent imaging (with CT, MRI, or ultrasound [US]) may be performed annually thereafter. CT and MRI have both been found to accurately predict pathologic tumor size in a retrospective analysis.(13) Therefore, best clinical judgment should be used in choosing the imaging modality. A chest x-ray or chest CT at baseline and annually as clinically indicated to assess for pulmonary metastases. Repeat chest imaging can be considered if intervention is being contemplated.

### Follow up after ablative therapy for Stage T1a

An abdominal imaging either CT or MRI with and without IV contrast (unless otherwise contraindicated) at 1 through 6 months to assess treatment response, followed by annual abdominal CT or MRI (preferred) for 5 years or longer as clinically indicated. If the patient cannot receive IV contrast, MRI is preferred

#### Follow up after partial or radical nephrectomy for Stages 1-2

Stage I RCC, it is recommended that abdominal CT or MRI (preferred) within 3 to 12 months following renal surgery, then annually for up to 5 years or longer as clinically indicated. For patients with stage II RCC, it is recommended to increase in abdominal imaging frequency, with baseline abdominal CT or MRI (preferred) every 6 months for 2 years, then annually for up to 5 years or longer, as clinically indicated.

It is also recommended that yearly chest x-ray or CT for at least 5 years and as clinically indicated thereafter.

#### Follow up for patients with Stage 3 RCC

It is recommended to obtain a baseline abdominal CT or MRI within 3 to 6 months following surgery, followed by CT, MRI (preferred), or US every 3 to 6 months for at least 3 years, and annually thereafter for up to 5 years.

A baseline chest CT within 3 to 6 months following surgery, is also recommended as well as continued imaging (CT preferred) every 3 to 6 months for at least 3 years, and annually thereafter for up to 5 years.

CT is the preferred modality for those with a high risk of recurrence.

#### Follow up for patient with relapse or unresectable disease or Stage 4 RCC

It is recommended to obtain chest, abdominal and pelvic imaging at baseline and then as clinically indicated based on clinical status, and therapeutic schedule.

#### **References:**

1. Kutikov A, Fossett LK, Ramchandani P et al: Incidence of benign pathologic findings at partial nephrectomy for solitary renal mass presumed to be renal cell carcinoma on preoperative imaging. Urology 2006; 68: 737.
2. Lane BR, Babineau D, Kattan MW et al: A preoperative prognostic nomogram for solid enhancing renal tumors 7 cm or less amenable to partial nephrectomy. J Urol 2007; 178: 429.
3. Johnson DC, Vukina J, Smith AB et al: Preoperatively misclassified, surgically removed benign renal masses: a systematic review of surgical series and United States population level burden estimate. J Urol 2015; 193: 30.
4. Davenport MS, Perazella MA, Yee J et al: Use of intravenous iodinated contrast media in patients with kidney disease: Consensus statements from the american college of radiology and the national kidney foundation. Radiology 2020; 294: 660.
5. Lim DJ, Carter MF: Computerized tomography in the preoperative staging for pulmonary metastases in patients with renal cell carcinoma. J Urol 1993; 150: 1112.

6. Mano R, Vertosick E, Sankin AI et al: Subcentimeter pulmonary nodules are not associated with disease progression in patients with renal cell carcinoma. *J Urol* 2015; 193: 776.
7. Weinreb JC, Rodby RA, Yee J et al: Use of intravenous gadolinium-based contrast media in patients with kidney disease: Consensus statements from the American College of Radiology and the national kidney foundation. *Radiology* 2021; 298: 28.
8. Seaman E, Goluboff ET, Ross S et al: Association of radionuclide bone scan and serum alkaline phosphatase in patients with metastatic renal cell carcinoma. *Urology* 1996; 48: 692.
9. Marshall ME, Pearson T, Simpson W et al: Low incidence of asymptomatic brain metastases in patients with renal cell carcinoma. *Urology* 1990; 36: 300.
10. Koga S, Tsuda S, Nishikido M et al: The diagnostic value of bone scan in patients with renal cell carcinoma. *J Urol* 2001; 166: 2126.
11. Campbell SC, Clark PE, Chang SS et al: Renal Mass and Localized Renal Cancer: Evaluation, Management, and Follow-Up: AUA Guideline Part I. *J Urol* 2021; **206**: 199.  
Campbell SC, Uzzo RG, Karam JA, et al: Renal Mass and Localized Renal Cancer: Evaluation, Management, and Follow-up: AUA Guideline: Part II. *J Urol* 2021; **206**: 209
12. Park JW, Jo MK, Lee HM. Significance of 18F-fluorodeoxyglucose positron-emission tomography/computed tomography for the postoperative surveillance of advanced renal cell carcinoma. *BJU Int* 2009;103:615-619
13. Dabestani S, Beisland C, Stewart GD, et al. Long-term outcomes of follow-up for initially localised clear cell renal cell carcinoma: RECUR database analysis. *Eur Urol Focus* 2019;5:857-866.
14. Motzer RJ, Jonasch E, et al. National Comprehensive Cancer Network (NCCN) Guidelines Version 3.2024 – March 11, 2024. Kidney Cancer, available at: [https://www.nccn.org/professionals/physician\\_gls/pdf/kidney.pdf](https://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf) Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines™) for Kidney Cancer 3.2024 – March 11, 2024. ©2024 National Comprehensive Cancer Network, Inc. All rights reserved. The NCCN Guidelines™ and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most recent and complete version of the NCCN Guidelines™, go online to NCCN.org.