

Evidence Discussion (HD-14.1, HD-14.2)

- A head CT is recommended for quickly assessing intracranial infections in urgent cases. However, MRI is more effective for examining conditions that affect the cranial nerves, brain tissue, and meninges.¹ Although MRI is superior in detecting minor changes in the brain associated with infections, CT scans can promptly identify pathophysiological changes that may influence the patient's prognosis.²
- Acute bacterial meningitis often presents with a normal CT scan. However, due to the high mortality rate of up to 50% when left untreated, a CT scan is necessary to rule out other causes of encephalopathy or neurologic deficits. In certain clinical situations, CT is also required to exclude increased intracranial pressure before performing a lumbar puncture. Venous thromboses increase the risk of hemorrhage and are linked to high mortality. Consequently, both CT and MRI scans, including T1-weighted sequences, are recommended. These scans help detect high signal changes in several venous sinuses and can be complemented with CT- or MR-venography.
- Similar considerations apply when diagnosing non-bacterial central nervous system (CNS) infections. CT scans are advantageous due to quick access to care, faster diagnosis, and earlier treatment initiation, which can reduce morbidity. However, MRI is superior in detecting patterns of vasogenic versus cytotoxic edema, contrast enhancement, and the distribution of involvement, whether multifocal or unifocal/unihemispheric. These distinctions are crucial in differentiating between differential diagnoses, such as systemic infections with hematogenous spread versus head/neck infections with a direct spread pattern.⁷
- Many patients present with neurologic signs and symptoms that are indicative of either a cortical or subcortical syndrome. Often, these cannot be fully characterized by clinical presentation alone. As a result, radiologic evaluation becomes essential in diagnosing the etiology of the underlying process. The potential causes are varied and include meningoencephalitis, acute cerebrovascular disease, hemorrhagic necrotizing encephalopathy, immune-mediated (Bickerstaff) encephalitis, and demyelinating diseases such as acute disseminated encephalomyelitis (ADEM) and multiple sclerosis (MS).¹³ Therefore, prompt and accurate diagnosis is critical to select the most appropriate imaging method (MRI vs. CT) for each clinical scenario.

References

1. ACR–ASNR–SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF COMPUTED TOMOGRAPHY (CT) OF THE HEAD <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CTHead.pdf>
2. Abdalkader M, Xie J, Cervantes-Arslanian A, Takahashi C, Mian AZ. Imaging of Intracranial Infections. *Seminars in Neurology*. 2019;39(03):322-333. doi:10.1055/s-0039-1693161
3. Allan R. Tunkel, Barry J. Hartman, Sheldon L. Kaplan, Bruce A. Kaufman, Karen L. Roos, W. Michael Scheld, Richard J. Whitley, Practice Guidelines for the Management of Bacterial Meningitis, *Clinical Infectious Diseases*, Volume 39, Issue 9, 1 November 2004, Pages 1267–1284, <https://doi.org/10.1086/425368>
4. Abdelbaker M et al. Imaging of Intracranial Infections. *Seminars in Neurology*. 2019.
5. Young N, Thomas M. Meningitis in adults: diagnosis and management. *Intern Med J*. 2018 Nov;48(11):1294-1307. doi: 10.1111/imj.14102. PMID: 30387309.
6. N. Proulx, D. Fréchette, B. Toye, J. Chan, S. Kravcik, Delays in the administration of antibiotics are associated with mortality from adult acute bacterial meningitis, *QJM: An*

7. Nguyen I, Urbanczyk K, Mtui E, Li S. Intracranial CNS Infections: A Literature Review and Radiology Case Studies. *Semin Ultrasound CT MR*. 2020 Feb;41(1):106-120. doi: 10.1053/j.sult.2019.09.003. Epub 2019 Oct 2. PMID: 31964490.
8. Sklinda K, Dorobek M, Wasilewski PG, Dreżewski K, Dębicka M, Walecki J, Mruk B. Radiological Manifestation of Neurological Complications in the Course of SARS-CoV-2 Infection. *Front Neurol*. 2021 Oct 20;12:711026. doi: 10.3389/fneur.2021.711026. PMID: 34744963; PMCID: PMC8563625.
9. Paterson RW, Brown RL, Benjamin L, Nortley R, Wiethoff S, Bharucha T, et al.. The emerging spectrum of COVID-19 neurology: clinical, radiological and laboratory findings. *Brain*. (2020) 143:3104–20. 10.1093/brain/awaa240
10. Kremer S, Lersy F, de Sèze J, Ferré JC, Maamar A, Carsin-Nicol B, et al.. Brain MRI findings in severe COVID-19: a retrospective observational study. *Radiology*. (2020) 297:E242–51. 10.1148/radiol.2020202222
11. Radmanesh A, Raz E, Zan E, Derman A, Kaminetzky M. Brain imaging use and findings in COVID-19: a single academic center experience in the epicenter of disease in the United States. *Am J Neuroradiol*. (2020) 41:1179–83. 10.3174/ajnr.A6610
12. Tu TM, Yi SJ, Koh JS, et al. Incidence of Cerebral Venous Thrombosis Following SARS-CoV-2 Infection vs mRNA SARS-CoV-2 Vaccination in Singapore. *JAMA Netw Open*. 2022;5(3):e222940. doi:10.1001/jamanetworkopen.2022.2940
13. Sklinda K et al. Radiological Manifestation of Neurological Complications in the Course of SARS-CoV-2 Infection. *Front Neurol*. 2021.