

Evidence Discussion – IBD (Crohn's Disease or Ulcerative Colitis) (AB-23.1)

In individuals with suspected inflammatory bowel disease, cross-sectional imaging can be performed after initial endoscopy is suggestive of inflammatory changes or if abnormal inflammatory markers concerning for IBD, or positive family history of IBD. Cross-sectional imaging methods such as computed tomography and magnetic resonance imaging are complementary to endoscopy, which allows diagnosis of disease when endoscopy is negative and diagnosis is still in doubt.

References

1. Lichtenstein GR, Loftus EV, Isaacs KL, Regueiro MD, Gerson LB, Sands BE. ACG Clinical Guideline: Management of Crohn's Disease in Adults. American Journal of Gastroenterology. 2018; 113(4):481-517. doi:10.1038/ajg.2018.27.
2. Hara AK, Leighton JA, Heigh RI, et al. Crohn Disease of the Small Bowel: Preliminary Comparison among CT Enterography, Capsule Endoscopy, Small-Bowel Follow-through, and Ileoscopy | Radiology.
3. Lin MF and Narra V. Developing role of magnetic resonance imaging in Crohn's disease. Current Opinion in Gastroenterology. 2008, 24(2):135-140.
4. Expert Panel on Gastrointestinal Imaging. ACR Appropriateness Criteria® Crohn's disease. American College of Radiology (ACR); Reviewed 2021.
5. Rubin DT, Ananthakrishnan AN, Siegel CA, Sauer BG, Long MD. ACG Clinical Guideline: Ulcerative Colitis in Adults. American Journal of Gastroenterology. 2019 Mar; 114(3):384-413.