

Evidence Discussion (ONC 31.5)

Bone Metastases

Bone scan supplemented by plain Xray is the generally the first-line modality for patients with current or history of malignancy who have new bone pain, rising tumor markers or elevated alkaline phosphatase. Bone scan is 79-86% sensitive and 81-88% specific for metastatic lesions (Yang 2011, Qu 2012). Bone scan allows rapid whole-skeletal evaluation, to ensure additional bony disease is not missed by focusing on a single site of cross sectional imaging. MRI is a useful problem-solving tool if there is continued suspicion for bony metastatic disease with a negative bone scan (DiPrimio 2023, Yang 2011). MRI is supported when a soft tissue component is suspect to avoid understaging and undertreatment (ACR 2024, DiPrimio 2023). MRI is also the most specific study to supplement plain xray to differentiate Paget disease of bone from neoplastic disease (Lombardi 2022).

MRI is more accurate than CT or bone scan for the evaluation of malignant vertebral compression fractures and additionally can assess for cord compression, edema or leptomeningeal disease (Liu 2017). Patients with known stage IV cancer with new back pain or any signs of neurologic compromise may be immediately evaluated by MRI of the whole spine without or without and with contrast (ACR 2024, Liu 2017). MRI is also indicated for suspected leptomeningeal disease (ACR 2024, Liu 2017). CT has the lower accuracy than MRI or bone scan in this setting, and is only supported when MRI is contraindicated (Liu 2017). New leptomeningeal disease should prompt an MRI of the brain for complete neuroaxis imaging..

Where imaging is suspicious for bony metastatic disease and a workup for multiple myeloma is negative, CT chest, abdomen and pelvis with contrast are supported to look for a primary malignancy (Karsell 1982, Piccoli 2015).

The sensitivity and specificity of FDG PET-CT for bony metastatic disease varies with the malignancy. For example, for breast cancer, PET may be more sensitive (96% vs 76% for bone scan), but may be less specific (92% vs 95% for bone scan). However for some cancers sensitivity of PET is as low as 56 percent (Liu 2017, Qu 201). Given this variability, PET-CT is supported as a problem-solving tool when both bone scan and MRI or CT are inconclusive. NaF PET is considered investigational due to varying sensitivity and specificity (Zhang-Yin 2023, Ahmed 2022).

References

1. Yang HL, Liu T, Wang XM, Xu Y, Deng SM. Diagnosis of bone metastases: a meta-analysis comparing ^{18}F FDG PET, CT, MRI and bone scintigraphy. *Eur Radiol*. 2011 Dec;21(12):2604-17. doi: 10.1007/s00330-011-2221-4. Epub 2011 Sep 2. PMID: 21887484.
2. Qu X, Huang X, Yan W, Wu L, Dai K. A meta-analysis of ^{18}F FDG-PET-CT, ^{18}F FDG-PET, MRI and bone scintigraphy for diagnosis of bone metastases in patients with lung cancer. *Eur J Radiol*. 2012 May;81(5):1007-15. doi: 10.1016/j.ejrad.2011.01.126. Epub 2011 Feb 26. PMID: 21354739
3. Di Primio G, Boyd GJ, Fung CI, Hurrell C, Brahm GL, Bird JR, Co SJ, Kirkpatrick IDC. Recommendations for the Management of Incidental Musculoskeletal Findings on MRI and CT. *Can Assoc Radiol J*. 2023 Aug;74(3):514-525. doi: 10.1177/08465371231152151. Epub 2023 Jan 29. PMID: 36710521.
4. Liu T, Wang S, Liu H, Meng B, Zhou F, He F, Shi X, Yang H. Detection of vertebral metastases: a meta-analysis comparing MRI, CT, PET, BS and BS with SPECT. *J Cancer Res Clin Oncol*. 2017 Mar;143(3):457-465. doi: 10.1007/s00432-016-2288-z. Epub 2016 Oct 17. PMID: 27752772.
5. Lombardi AF, Aihara AY, Fernandes ADRC, Cardoso FN. Imaging of Paget's Disease of Bone. *Radiol Clin North Am*. 2022 Jul;60(4):561-573. doi: 10.1016/j.rcl.2022.02.005. PMID: 35672089.
6. Zhang-Yin J, Panagiotidis E. Role of ^{18}F -NaF PET/CT in bone metastases. *Q J Nucl Med Mol Imaging*. 2023 Dec;67(4):249-258. doi: 10.23736/S1824-4785.23.03534-3. Epub 2023 Sep 26. PMID: 37750848.
7. American College of Radiology. ACR Appropriateness Criteria®. Available at <https://acsearch.acr.org/list> . Accessed 6/11/2024
8. Karsell PR, Sheedy PF 2nd, O'Connell MJ. Computed tomography in search of cancer of unknown origin. *JAMA*. 1982 Jul 16;248(3):340-3. PMID: 7087129.
9. Piccioli A, Maccauro G, Spinelli MS, Biagini R, Rossi B. Bone metastases of unknown origin: epidemiology and principles of management. *J Orthop Traumatol*. 2015 Jun;16(2):81-6. doi: 10.1007/s10195-015-0344-0. Epub 2015 Mar 1. PMID: 25726410; PMCID: PMC4441638.
10. Ahmed N, Sadeq A, Marafi F, Gnanasegaran G, Usmani S. Therapy-induced bone changes in oncology imaging with ^{18}F -sodium fluoride (NaF) PET-CT. *Ann Nucl Med*. 2022 Apr;36(4):329-339. doi: 10.1007/s12149-022-01730-y. Epub 2022 Feb 26. PMID: 35218508.