

Evidence Discussion (PVD 3.2)

Surveillance Imaging with NO History of Carotid Surgery or Intervention

DU is established as the primary diagnostic test for carotid surveillance imaging. Consensus statements have established the appropriate time intervals for repeat imaging. If however, there are new symptoms or physical findings, repeat duplex imaging is supported, regardless of time interval. If at any time there is $\geq 70\%$ re-stenosis, CTA or MRA is indicated.³⁻⁵

1. Grant, EG, Benson, CB, et al. Carotid artery stenosis: gray-scale and Doppler US diagnosis--Society of Radiologists in Ultrasound Consensus Conference Radiology. 2003 Nov;229(2):340-6.
2. Wardlaw JM, Chappell FM, Stevenson M, et al. Accurate, practical and cost-effective assessment of carotid stenosis in the UK. Health Technol Assess. 2006;10(30).
3. Ali F. AbuRahma, MD,a Efthymios D., et al. Society for Vascular Surgery clinical practice guidelines for management of extracranial cerebrovascular disease. J Vasc Surg. 2022 Jan;75(1S):4S-22S.
4. Ricotta J, AbuRahma A, Ascher E, et al. Updated Society for Vascular Surgery guidelines for management of extracranial carotid disease. J Vasc Surg. 2011;54(3):e1–e31.
5. AbuRahma A.F, Srivastava M, Stone P.A, et al. Critical appraisal of the Carotid Duplex Consensus criteria in the diagnosis of carotid artery stenosis. J Vasc Surg. 2011;53:53–60.