

Cigna Medical Coverage Policies – Musculoskeletal Thoracic and Thoracolumbar Fusion (Arthrodesis) Guidelines

Effective November 1, 2024



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by eviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2024 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

©Copyright 2024 eviCore healthcare

CMM-614: Thoracic / Thoracolumbar Fusion (Arthrodesis)

CMM-614.1: General Guidelines

CMM-614.2: Osteotomy

CMM-614.3 Pediatric Spinal Deformity

CMM-614.4: Thoracic or Thoracolumbar Fusion (Arthrodesis) with Decompression

CMM-614.5: Thoracic or Thoracolumbar Fusion (Arthrodesis) without Decompression

CMM-614.6: Adjacent Segment Disease/Proximal Junctional Failure

CMM-614.7: Repeat Thoracic Fusion (Arthrodesis) at the Same Level

CMM-614.8: Non-Indications

Procedure (CPT®) Codes (CMM-614)

References (CMM-614)

CMM-614.1: General Guidelines

Application of Guideline

- The determination of medical necessity for the performance of thoracic or thoracolumbar fusion (with or without osteotomy) is always made on a case-by-case basis.
- For additional timing and documentation requirements, see **CMM-600.1: Prior Authorization Requirements**.

Urgent/Emergent Indications/Conditions

- The presence of urgent/emergent indications/conditions warrants definitive surgical treatment. **Imaging findings noted in the applicable procedure section(s) are required.**
 - ◆ The following criteria are **NOT** required for confirmed urgent/emergent conditions:
 - Provider-directed non-surgical management
 - Proof of smoking cessation
 - Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)
 - Timeframe for repeat procedure
- Urgent/emergent conditions for thoracic or thoracolumbar fusion and/or osteotomy include **ANY** of the following:
 - ◆ Traumatic spinal fractures or dislocations with or without neural compression when instability is present or decompression of the spinal canal is anticipated to result in iatrogenic instability
 - ◆ Infection (e.g., discitis, epidural abscess, osteomyelitis) when instability is present or debridement and/or decompression is anticipated to result in iatrogenic instability
 - ◆ Primary or metastatic neoplastic disease causing pathologic fracture, cord compression, when instability is present or resection and/or decompression is anticipated to result in iatrogenic instability
 - ◆ A condition otherwise meeting criteria listed in the applicable procedure section(s) with documentation of severe debilitating pain and/or dysfunction to the point of being incapacitated

CMM-614.2: Osteotomy

Posterior Column Osteotomy (PCO)

Thoracic or thoracolumbar posterior column osteotomy (PCO) (i.e., Smith-Peterson osteotomy [SPO] or Ponte osteotomy) is considered **medically necessary** (in addition to fusion) when **ALL** of the following criteria have been met:

- Correction of non-fixed deformity requiring 5° to 10° of correction (SPO) per spinal segment for **EITHER** of the following:
 - ◆ Thoracic or thoracolumbar sagittal plane deformities (i.e., <10° or >40°) when the deformity is non-fixed
 - ◆ Larger coronal deformities where there is limited flexibility (i.e., >40° on dynamic imaging)
- Posterior column osteotomy is limited to a maximum of four (4) posterior column osteotomies performed in the apex of the deformity per correction surgery.
 - ◆ **Criteria exception:** There is no limit to posterior column osteotomies for correction of Scheuermann's Kyphosis as this deformity is long, gradual, rounded, and is amenable to more than four (4) posterior column osteotomies.
- **ALL** of the criteria for thoracic or thoracolumbar fusion have been met per the applicable procedure-specific section (pediatric deformity; fusion with decompression; fusion without decompression; adjacent segment disease; or, repeat fusion)

Three-Column Osteotomy

Thoracic or thoracolumbar three-column osteotomy (i.e., pedicle subtraction osteotomy [PSO] or vertebral column resection [VCR]) is considered **medically necessary** (in addition to fusion) when **ALL** of the following criteria have been met:

- Performed for **EITHER** of the following:
 - ◆ Correction of fixed sagittal plane deformity requiring more than 30° of correction (PSO)
 - ◆ Large fixed coronal deformities greater than 60° that are amenable to asymmetric osteotomy
- **ALL** of the criteria for thoracic or thoracolumbar fusion have been met per the applicable procedure-specific section (pediatric spinal deformity; fusion with decompression; fusion without decompression; adjacent segment disease; or, repeat fusion)

CMM-614.3: Pediatric Spinal Deformity

Pediatric Thoracic or Thoracolumbar Fusion

Thoracic or thoracolumbar fusion (arthrodesis) is considered **medically necessary** when performed for **ANY** of the following conditions when **BOTH** of the following criteria have been met:

- Confirmatory imaging studies (advanced or plain X-rays) are required
- The presence of **ANY** of the following pediatric spinal deformities warrants definitive surgical treatment:
 - ◆ Adolescent idiopathic scoliosis with over 50° curve
 - ◆ Congenital scoliosis
 - ◆ Neuromuscular scoliosis
 - ◆ Infantile/juvenile scoliosis

Pediatric Osteotomy

Thoracic or thoracolumbar osteotomy is considered **medically necessary** (in addition to a fusion) when **ALL** of the following criteria have been met per **CMM-614.2: Osteotomy**.

CMM-614.4: Initial Thoracic or Thoracolumbar Fusion (Arthrodesis) with Decompression

Thoracic or thoracolumbar fusion (arthrodesis) with decompression is considered **medically necessary** when performed for **ANY** of the following conditions when **ALL** of the associated criteria have been met:

Actual Instability

- The individual is a candidate for thoracic decompression per **CMM-613: Thoracic Decompression-Discectomy**.
- Imaging shows **ANY** of the following:
 - ◆ Degenerative spondylolisthesis with **EITHER** of the following:
 - Dynamic segmental instability on flexion-extension plain X-rays **OR** comparison of a supine and upright image, with a difference in translational alignment between vertebrae greater than 3 mm between views
 - Meyerding Grade II or higher spondylolisthesis
 - ◆ Imaging documenting post-operative instability created by the disruption of the posterior elements due to facet joint excision that exceeds 50% bilaterally or 75% or more of a single facet
 - ◆ Pars fracture
 - ◆ Previous thoracic or thoracolumbar spinal decompression that resulted in iatrogenic spondylolisthesis
 - ◆ **Criteria exception:** When instability is created and/or identified intra-operatively, the above imaging criteria are **NOT** required.
 - See **Anticipated Iatrogenic Instability**

- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤ 10 ng/mL

Anticipated Iatrogenic Instability

- The individual is a candidate for thoracic decompression or corpectomy per **CMM-613: Thoracic Decompression-Discectomy**.
- Anticipated iatrogenic instability with **ANY** of the following:
 - ◆ Created by disruption of the posterior elements due to facet joint excision that exceeds 50% bilaterally or 75% or more of a single facet during spinal decompression
 - ◆ Created by excessive disc removal that could result in anterior instability
 - ◆ Created by removal of the pars interarticularis that requires fusion to stabilize
 - ◆ Created by decompression for Meyerding Grade I or higher spondylolisthesis with foraminal stenosis
 - ◆ Created by complete or partial thoracic/thoracolumbar corpectomy (i.e., removal of at least one-third of the vertebral body)
 - For thoracic or thoracolumbar corpectomy, see **CMM-613.1: General Guidelines**.
- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤ 10 ng/mL

Adult Symptomatic Degenerative Spinal Deformity

- The individual is a candidate for thoracic decompression per **CMM-613: Thoracic Decompression-Discectomy**.
- Imaging findings include **EITHER** of the following:
 - ◆ Coronal plane deformity which includes **ANY** of the following:
 - Cobb angle greater than 30°
 - Asymmetric disk collapse causing foraminal narrowing
 - Coronal imbalance causing head and trunk shift off the midline
 - ◆ Sagittal imbalance which includes **ANY** of the following:
 - Sagittal vertebral axis measurement greater than 8 cm
 - Pelvic incidence-lumbar lordosis greater than 15°
- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤ 10 ng/mL

Adult Residual Idiopathic Scoliotic Deformity

- The individual is a candidate for thoracic decompression or corpectomy per **CMM-613: Thoracic Decompression-Discectomy**.
- Imaging findings include **ANY** of the following:
 - ◆ Cobb angle of greater than 50°
 - ◆ Asymmetric disk collapse causing foraminal narrowing
 - ◆ Coronal imbalance causing head and trunk shift off the midline
- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤10 ng/mL

Adult Symptomatic Residual Scheuermann's Kyphosis

- The individual is a candidate for thoracic decompression or corpectomy per **CMM-613: Thoracic Decompression-Discectomy**.
- Imaging findings include **ANY** of the following:
 - ◆ Sagittal Vertebral Axis measurement greater than 8 cm
 - ◆ Pelvic incidence-lumbar lordosis greater than 15°
 - ◆ Sagittal Cobb angle greater than 75°
- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤10 ng/mL

CMM-614.5: Initial Thoracic or Thoracolumbar Fusion (Arthrodesis) without Decompression

Thoracic or thoracolumbar fusion without decompression is considered **medically necessary** when performed for **ANY** of the following conditions when **ALL** of the associated criteria have been met:

Adult Symptomatic Degenerative Spinal Deformity

- Imaging shows **EITHER** of the following:
 - ◆ Coronal plane deformity which includes **ANY** of the following:
 - Cobb angle of greater than 30°
 - Asymmetric disk collapse causing foraminal narrowing
 - Coronal imbalance causing head and trunk shift off the midline
 - ◆ Sagittal imbalance which includes **ANY** of the following:
 - Sagittal vertebral axis measurement greater than 8 cm
 - Pelvic incidence-lumbar lordosis greater than 15°
- Less than clinically meaningful improvement with **EITHER** of the following for at least 3 consecutive months (unless contraindicated):
 - ◆ Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs
 - ◆ Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)
- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤10 ng/mL

Adult Residual Idiopathic Scoliotic Deformity

- Imaging shows **ANY** of the following:
 - ◆ Cobb angle greater than 50°
 - ◆ Asymmetric disk collapse causing foraminal narrowing
 - ◆ Coronal imbalance causing head and trunk shift off the midline
- For those cases with a less than 50° curve or global coronal imbalance, less than clinically meaningful improvement with **EITHER** of the following for at least 3 consecutive months (unless contraindicated):
 - ◆ Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs
 - ◆ Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)

- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤ 10 ng/mL

Adult Symptomatic Residual Scheuermann's Kyphosis

- Imaging shows **ANY** of the following:
 - ◆ Sagittal Vertebral Axis measurement greater than 8 cm
 - ◆ Pelvic incidence-lumbar lordosis greater than 15°
 - ◆ Sagittal Cobb angle greater than 75°
- Less than clinically meaningful improvement with **EITHER** of the following for at least 3 consecutive months (unless contraindicated):
 - ◆ Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs
 - ◆ Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)
- Documentation of nicotine-free status with **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤ 10 ng/mL

CMM-614.6: Adjacent Segment Disease/Proximal Junctional Failure

Thoracic or thoracolumbar fusion for adjacent segment disease is considered **medically necessary** when **ALL** of the following criteria have been met:

- The individual meets criteria for thoracic or thoracolumbar fusion per the applicable procedure-specific section below:
 - ◆ **CMM-614.4: Initial Thoracic or Thoracolumbar Fusion (Arthrodesis) with Decompression**
 - ◆ **CMM-614.5: Initial Thoracic or Thoracolumbar Fusion (Arthrodesis) without Decompression**
- The prior thoracic fusion procedure at an adjacent level was performed at least 6 months prior
- Plain X-rays show **EITHER** of the following:
 - ◆ Evidence of anterolisthesis on plain X-rays resulting in **ANY** of the following:
 - Dynamic segmental instability on flexion-extension plain X-rays **OR** comparison of a supine and upright image, with a difference in translational alignment between vertebra greater than 3 mm between views
 - Meyerding Grade II or higher spondylolisthesis
 - ◆ Evidence of kyphosis on plain X-rays

- Any spinal level showing 15° change of the sagittal Cobb angle (proximal junctional angle) with progressive pain or neurologic symptoms caused by either bony or posterior ligamentous complex failure

CMM-614.7: Repeat Thoracic or Thoracolumbar Fusion (Arthrodesis) at the Same Level

Repeat thoracic or thoracolumbar fusion (with or without thoracic or thoracolumbar decompression) at the same level is considered **medically necessary** for **EITHER** of the following conditions when **ALL** of the associated criteria have been met:

Malposition or Failure of Implant/Instrumentation or Structural Bone Graft

- Post-operative imaging shows evidence of malposition or failure of the implant/instrumentation or structural bone graft (e.g., migration, pedicle screw breakage, pedicle screw loosening, dislodged hooks, rod breakage, rod bending, rod loosening, loss of curve correction, decompensation, etc.)

Symptomatic Pseudoarthrosis

- Greater than 6 months since the last thoracic or thoracolumbar fusion surgery
- Subjective symptoms include significant level of pain on a daily basis defined as clinically significant functional impairment (e.g., inability to perform household chores, prolonged standing, etc.)
- Post-operative physical examination findings are concordant with the individual's symptoms
- Less than clinically meaningful improvement with 6 weeks of non-surgical treatment with **BOTH** of the following (unless contraindicated):
 - ◆ Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs
 - ◆ Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician
- Post-operative imaging (performed at no less than 6 months after the prior thoracic or thoracolumbar fusion) shows pseudoarthrosis at the requested level(s)
- Post-operative MRI/CT findings are concordant with the individual's symptoms
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)
- Documentation of nicotine-free status including **EITHER** of the following:
 - ◆ Individual is a never-smoker
 - ◆ Individual has refrained from smoking, use of smokeless tobacco products, and/or nicotine replacement therapy for at least 6 weeks prior to planned surgery as evidenced by blood cotinine lab results of ≤ 10 ng/mL

CMM-614.8: Non-Indications

Not Medically Necessary

- **Thoracic or thoracolumbar fusion** performed without meeting the criteria in the **General Guidelines** (when applicable for urgent/emergent conditions) **and** the criteria in the applicable procedure-specific section(s) (pediatric spinal deformity; fusion with decompression; fusion without decompression; adjacent segment disease; or, repeat fusion) is considered **not medically necessary**
- **Thoracic or thoracolumbar osteotomy** performed without meeting the criteria in the **General Guidelines** (when applicable for urgent/emergent conditions) **and** the criteria in the applicable procedure-specific section (osteotomy; pediatric spinal deformity; fusion with decompression; fusion without decompression; adjacent segment disease; or, repeat fusion) is considered **not medically necessary**.
- **Thoracic or thoracolumbar fusion** and/or **osteotomy** performed for **ANY** of the following sole indications is considered **not medically necessary**:
 - ◆ Multi-level degenerative disc disease without instability or significant adult deformity
 - ◆ Facet joint disorders without instability
 - ◆ Initial discectomy/laminectomy without instability
 - ◆ An adjunct to primary decompression of central and/or lateral recess stenosis in the absence of instability, spondylolisthesis, adult deformity, or an actual or anticipated bony resection that will result in iatrogenic instability

Experimental, Investigational, or Unproven (EIU)

Thoracic or Thoracolumbar Fusion

- Thoracic or thoracolumbar fusion is considered **experimental, investigational, or unproven (EIU)** for **ALL** of the following devices/procedures (not an all-inclusive list):
 - ◆ Minimally invasive thoracic or thoracolumbar fusion using direct visualization via endoscopy (endoscopic fusion) or indirect visualization (e.g., percutaneous fusion)
 - ◆ Endoscopic thoracic decompression with interbody fusion
 - ◆ Isolated facet fusion, with or without instrumentation, including allograft bone graft substitutes used exclusively as stand-alone stabilization devices (e.g., TruFuse [any level], NuFix® [any level])
 - ◆ Total facet arthroplasty
 - ◆ Device/implant not FDA approved (in thoracic spine)

Procedure (CPT®) Codes (CMM-614)

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
22206	Osteotomy of spine, posterior or posterolateral approach, 3 columns, 1 vertebral segment (e.g., Pedicle/vertebral body subtraction); thoracic
+22208	Osteotomy of spine, posterior or posterolateral approach, 3 columns, 1 vertebral segment (e.g., Pedicle/vertebral body subtraction); each additional vertebral segment (List separately in addition to code for primary procedure)
22212	Osteotomy of spine, posterior or posterolateral approach, 1 vertebral segment; thoracic
+22216	Osteotomy of spine, posterior or posterolateral approach, 1 vertebral segment; each additional vertebral segment (List separately in addition to code for primary procedure)
22222	Osteotomy of spine, including discectomy, anterior approach, single vertebral segment; thoracic
+22226	Osteotomy of spine, including discectomy, anterior approach, single vertebral segment, each additional vertebral segment (List separately in addition to code for primary procedure)
22532	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic
+22534	Arthrodesis, lateral extracavitary technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic or lumbar, each additional vertebral segment (List separately in addition to code for primary procedure)
22556	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); thoracic
+22585	Arthrodesis, anterior interbody technique, including minimal discectomy to prepare interspace (other than for decompression); each additional interspace (List separately in addition to code for primary procedure)
22610	Arthrodesis, posterior or posterolateral technique, single level; thoracic (with lateral transverse technique, when performed)
+22614	Arthrodesis, posterior or posterolateral technique, single level; each additional vertebral segment (List separately in addition to code for primary procedure)
22800	Arthrodesis, posterior, for spinal deformity, with or without cast; up to 6 vertebral segments

CMM-614: Thoracic / Thoracolumbar Fusion (Arthrodesis)

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
22802	Arthrodesis, posterior, for spinal deformity, with or without cast; 7 to 12 vertebral segments
22804	Arthrodesis, posterior, for spinal deformity, with or without cast; 13 or more vertebral segments
22808	Arthrodesis, anterior, for spinal deformity, with or without cast; 2 to 3 vertebral segments
22810	Arthrodesis, anterior, for spinal deformity, with or without cast; 4 to 7 vertebral segments
22812	Arthrodesis, anterior, for spinal deformity, with or without cast; 8 or more vertebral segments
63003	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (eg, spinal stenosis), 1 or 2 vertebral segments; thoracic
63016	Laminectomy with exploration and/or decompression of spinal cord and/or cauda equina, without facetectomy, foraminotomy or discectomy (eg, spinal stenosis), more than 2 vertebral segments; thoracic
63046	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [eg, spinal or lateral recess stenosis]), single vertebral segment; thoracic
+63048	Laminectomy, facetectomy and foraminotomy (unilateral or bilateral with decompression of spinal cord, cauda equina and/or nerve root[s], [eg, spinal or lateral recess stenosis]), single vertebral segment; each additional segment, cervical, thoracic, or lumbar (List separately in addition to code for primary procedure)
63055	Transpedicular approach with decompression of spinal cord, equina and /or nerve root(s) (e.g., herniated intervertebral disc), single segment; thoracic
+ 63057	Transpedicular approach with decompression of spinal cord, equina and/or nerve(s) (e.g., herniated intervertebral disc), single segment; each additional segment, thoracic or lumbar (List separately in addition to code for primary procedure)
63077	Discectomy, anterior, with decompression of spinal cord and/or nerve root(s), including osteophytectomy; thoracic, single interspace
+63078	Discectomy, anterior, with decompression of spinal cord and/or nerve root(s), including osteophytectomy; thoracic, each additional interspace (List separately in addition to code for primary procedure)

CMM-614: Thoracic / Thoracolumbar Fusion (Arthrodesis)

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code’s inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definition
63266	Laminectomy for excision or evacuation of intraspinal lesion other than neoplasm, extradural; thoracic
0220T	Placement of a posterior intrafacet implant(s), unilateral or bilateral, including imaging and placement of bone graft(s) or synthetic device(s), single level; thoracic

This list may not be all-inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual’s policy or benefit entitlement structure as well as claims processing rules.

References (CMM-614)

1. American Academy of Orthopedic Surgeons (AAOS). *Surgery and smoking*. July 2007.
2. American Medical Association. Code 20660 as an Independent or Unrelated Procedure- Coding Tip. *CPT® Assistant Newsletter*. April 2012;11. Accessed October 5, 2023. Available at: <https://ocm.ama-assn.org/OCM/CPTAA/Newsletters.do?articleType=IssueArticle&filename=20120411&hitTerms=corpectomy>.
3. Bridwell KH, Lewis SJ, Lenke LG, Baldus C, Blanke K. Pedicle subtraction osteotomy for the treatment of fixed sagittal imbalance. *JBJS*. 2003;85(3):454-463. doi: 10.2106/00004623-200303000-00009.
4. Bridwell KH, Lewis SJ, Rinella A, Lenke LG, Baldus C, Blanke K. Pedicle subtraction osteotomy for the treatment of fixed sagittal imbalance. Surgical technique. *JBJS*. 2004;86(suppl_1):44-50. doi: 10.2106/00004623-200403001-00007.
5. Chatley A, Kumar R, Jain V, Behari S, Sahu R. Effect of spinal cord signal intensity changes on clinical outcome after surgery for cervical spondylotic myelopathy. *J Neurosurg Spine*. 2009;11(5):562-567. doi: 10.3171/2009.6.spine091.
6. Cheung JPY. The importance of sagittal balance in adult scoliosis surgery. *Ann Transl Med*. 2020;8(2):35. doi: 10.21037/atm.2019.10.19.
7. Cohen SP, Hanling S, Bicket MC, et al. Epidural steroid injections compared with gabapentin for lumbosacral radicular pain: multicenter randomized double blind comparative efficacy study. *BMJ*. 2015;350(apr16 26):h1748-h1748. doi: 10.1136/bmj.h1748.
8. Danielsson, A. Natural history of adolescent idiopathic scoliosis: a tool for guidance in decision of surgery of curves above 50°. *J Child Orthop*. 2013;7:37-41. doi: 10.1007/s11832-012-0462-7.
9. Diebo BG, Varghese JJ, Lafage R, Schwab FJ, Lafage V. Sagittal alignment of the spine: What do you need to know? *Clin Neurol Neurosurg*. 2015;139:295-301. doi: 10.1016/j.clineuro.2015.10.024.
10. Dorward IG, Lenke LG. Osteotomies in the posterior-only treatment of complex adult spinal deformity: a comparative review. *Neurosurg Focus*. 2010;28(3):E4. doi: 10.3171/2009.12.focus09259.
11. Glassman SD, Anagnost SC, Parker A, et al. The effect of cigarette smoking and smoking cessation on spinal fusion. *Spine (Phila Pa 1976)*. 2000;25(20):2608-2615.
12. Graham RB, Sugrue PA, Koski TR. Adult degenerative scoliosis. *Clin Spine Surg*. 2016;29(3):95-107. doi: 10.1097/BSD.0000000000000367.
13. Halanski MA, Cassidy JA. Do multilevel Ponte osteotomies in thoracic idiopathic scoliosis surgery improve curve correction and restore thoracic kyphosis?. *J Spinal Disord Tech*. 2013;26(5):252-255. doi: 10.1097/BSD.0b013e318241e3cf.
14. Jackson KL, Devine JG. The Effects of Smoking and Smoking Cessation on Spine Surgery: A Systemic Review of the Literature. *Global Spine J*. 2016; 6(7):695-701. doi: 10.1055/s-0036-1571285.
15. Kelly DM, McCarthy RE, McCullough FL, Kelly HR. Long-term outcomes of anterior spinal fusion with instrumentation for thoracolumbar and lumbar curves in adolescent idiopathic scoliosis. *Spine*. 2010;35(2):194-198. doi: 10.1097/BRS.0b013e3181bc948e.
16. La Marca F, Brumblay H. Smith-Petersen Osteotomy in Thoracolumbar Deformity Surgery. *Neurosurgery*. 2008;63(3):A163-A170. doi: 10.1227/01.neu.0000320428.67620.4f.
17. Lee BS, Nault R, Grabowski M, et al. Utility of repeat magnetic resonance imaging in surgical patients with lumbar stenosis without disc herniation. *Spine J*. 2019;19:191-198.
18. Lindstrom D, Azodi OS, Wladis A, et al. Effects of a perioperative smoking cessation intervention on postoperative complications: a randomized trial. *Ann Surg*. 2008;248:739-745.
19. McCullen G, Vaccaro AR, Garfin SR. Thoracic and lumbar trauma: Rationale for selecting the appropriate fusion technique. *Orthop Clin North Am*. 1998;29(4):813-828.
20. Miller DJ, Cahill PJ, Vitale MG, Shah SA. Posterior Correction Techniques for Adolescent Idiopathic Scoliosis. *JAAOS*. Published online October 2019;1. doi: 10.5435/jaaos-d-18-00399.
21. Mills E, Eyawo O, Lockhart I, et al. Smoking cessation reduces postoperative complications: a systematic review and meta-analysis. *Am J Med*. 2011;124(2):144-154.
22. Moller AM, Villebro N, Pedersen T, et al. Effect of preoperative smoking intervention on postoperative complications: a randomized clinical trial. *Lancet*. 2002;359:114-117.
23. Myers K, Hajek P, Hinds C, et al. Stopping smoking shortly before surgery and postoperative complications: a systematic review and meta-analysis. *Arch Intern Med*. 2011;171: 983-989.
24. Naqvi U, Sherman A. *Muscle Strength Grading*. Ncbi.nlm.nih.gov. In: *Statpearls [Internet]*. Sep 2021. StatPearls Publishing. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK436008/>. Published 2021.
25. Osman NS, Cheung ZB, Hussain AK, et al. Outcomes and complications following laminectomy alone for thoracic myelopathy due to ossified ligamentum flavum: A systematic review and meta-analysis. *Spine (Phila Pa 1976)*. 2018;43(14):E842-E848.
26. Panagopoulos J, Hush J, Steffens D, Hancock MJ. Do MRI Findings Change Over a Period of Up to 1 Year in Patients With Low Back Pain and/or Sciatica? *Spine*. 2017;42(7):504-512. doi: 10.1097/brs.0000000000001790.
27. Raizman NM, O'Brien JR, Poehling-Monaghan KL, Yu WD. Pseudarthrosis of the Spine. *JAAOS*. 2009;17(8):494-503. doi: 10.5435/00124635-200908000-00003.

28. Ries ZG, Glassman SD, Vasilyev I, Metcalfe L, Carreon LY. Updated imaging does not affect revision rates in adults undergoing spine surgery for lumbar degenerative disease. *J Neurosurg Spine*. Published online Nov 2018. 2019;30(2):228-223. doi: 10.3171/2018.8.spine18586.
29. Saifi C, Laratta JL, Petridis P, Shillingford JN, Lehman RA, Lenke LG. Vertebral Column Resection for Rigid Spinal Deformity. *Global Spine J*. 2017;7(3):280-290. doi: 10.1177/2192568217699203.
30. Sardar ZM, Ames RJ, Lenke L. Scheuermann's Kyphosis: Diagnosis, Management, and Selecting Fusion Levels. *J Am Acad Orthop Surg*. 2019;27(10):e462-e472.
31. Siecean A, Seicean S, Alan N, et al. Effect of smoking on the perioperative outcomes of patients who undergo elective spine surgery. *Spine*. 2013;38:1294-1302. doi: 10.1097/BRS.0b013e31828e2747.
32. Sivaganesan A, Kim HJ. A Review of Indications, Surgical Technique, and Outcomes for the Cervical Pedicle Subtraction Osteotomy. *JAAOS*. 2022;30(3):e295-e300. doi: 10.5435/jaaos-d-21-00177.
33. Smith JS, Shaffrey CI, Glassman SD, et al. Risk-benefits assessment of surgery for adult scoliosis: An analysis based on patient age. *Spine*. 2011;36(10):817-824. doi: 10.1097/BRS.0b013e3181e21783.
34. Smith JS, Shaffrey CI, Lafage V, et al. Comparison of best versus worst clinical outcomes for adult spinal deformity surgery: A retrospective review of a prospectively collected, multicenter database with 2-year follow-up. *J Neurosurg Spine*. 2015;23(3):349-359. doi: 10.3171/2014.12.SPINE14777.
35. Sorensen LT. Wound healing and infection in surgery: the pathophysiological impact of smoking, smoking cessation, and nicotine replacement therapy: a systemic review. *Ann Surg*. 2012;255:1069-1079. doi: 10.1097/SLA.0b013e31824f632d.
36. Theadom A, Cropley M. Effects of preoperative smoking cessation on the incidence and risk of intraoperative and postoperative complications in adult smokers: a systematic review. *Tobacco Control*. 2006;15:352-358.
37. Tsuji H. Laminoplasty for patients with compressive myelopathy due to so-called spinal canal stenosis in cervical and thoracic regions. *Spine*. 1982;7(1):28-34.
38. Wan SH, Wong DL, To SC, Meng N, Zhang T, Cheung JP. Patient and surgical predictors of 3D correction in posterior spinal fusion: a systematic review. *Eur Spine J*. 2023;32(6):1927-1946. doi: 10.1007/s00586-023-07708-2.
39. Wong J, Lam DP, Abrishami A, et al. Short-term preoperative smoking cessation and postoperative complications: a systematic review and meta-analysis. *Can J Anaesth*. 2012;59(3):268-279.
40. Yaksi A, Özgönenel L, Özgönenel B. The Efficiency of Gabapentin Therapy in Patients With Lumbar Spinal Stenosis. *Spine*. 2007;32(9):939-942. doi: 10.1097/01.brs.0000261029.29170.e6.
41. Yin L, Zhang J, Wu Y, Li J, Yang Q. Increased signal intensity of spinal cord on T2W magnetic resonance imaging for cervical spondylotic myelopathy patients. *Medicine (Baltimore)*. 2020;99(49):e23098. doi: 10.1097/md.00000000000023098.
42. York PJ, Kim HJ. Degenerative Scoliosis. *Curr Rev Musculoskelet Med*. 2017;10(4):547-558. doi: 10.1007/s12178-017-9445-0.
43. York PJ, Kim HJ. Degenerative scoliosis. *Curr Rev Musculoskelet Med*. 2017;10(4):547-558.