



CLINICAL GUIDELINES

CMM-605: Cervical Microdiscectomy Guidelines

Effective November 1, 2024

VERSION 2.0.2024

EviCore
By **EVERNORTH**

EviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

EviCore's Clinical Review Criteria (CRC) and related content is made available for the limited uses of: reference; and individual use, only limited to facilitating the determination of medically necessary and appropriate clinical treatment by clinicians for specific delegated patients under their care. The CRC and related content is proprietary information of EviCore, and copyrighted to the full extent of the law. Except as expressly permitted, you may not modify, copy, reproduce, republish, upload, post, transmit, hyperlink to or from, or distribute in any way the CRC, nor may you sell, transfer, distribute, assign, lease, reproduce, or otherwise use the CRC in commerce, in a manner that competes with us or infringes upon our rights, or for any public or commercial endeavor without our prior and express written consent.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2024 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

©2024 EVICORE HEALTHCARE. ALL RIGHTS RESERVED

CMM-605: Cervical Microdiscectomy

CMM-605.1: General Guidelines**CMM-605.2: Initial Primary Cervical Microdiscectomy****CMM-605.3: Repeat Cervical Microdiscectomy at the Same Level****CMM-605.4: Non-Indications****Procedure (CPT®) Codes (CMM-605)****References (CMM-605)**

CMM-605.1: General Guidelines

Application of Guideline

- The determination of medical necessity for the performance of cervical microdiscectomy is always made on a case-by-case basis.
- For additional timing and documentation requirements, see **CMM-600.1: Prior Authorization Requirements**.

Urgent/Emergent Conditions/Indications

- The presence of urgent/emergent indications/conditions warrants definitive surgical treatment. **Imaging findings noted in the applicable procedure section(s) are required.**
 - ◆ The following criteria are **NOT** required for confirmed urgent/emergent conditions:
 - ◆ Provider-directed non-surgical management
 - ◆ Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)
 - ◆ Timeframe for repeat procedure
- Urgent/emergent conditions for cervical microdiscectomy include **ANY** of the following:
 - ◆ Central cord syndrome
 - ◆ Myelopathy or Cord signal changes on MRI due to cord compression
 - ◆ Documentation of progressive neurological deficit on two separate physical exams
 - ◆ **ANY** of the following due to a neurocompressive pathology
 - Motor weakness of grade 3/5 or less of specified muscle(s)
 - Rapidly progressive symptoms of motor loss
 - Bowel incontinence
 - Bladder incontinence/retention
 - ◆ A condition otherwise meeting criteria listed in the applicable procedure section(s) with documentation of severe debilitating pain and/or dysfunction to the point of being incapacitated

CMM-605.2: Initial Primary Cervical Microdiscectomy

Initial primary cervical microdiscectomy is considered **medically necessary** when performed for **EITHER** of the following conditions when **ALL** of the associated criteria have been met:

Radiculopathy

- Subjective symptoms include **BOTH** of the following:
 - ◆ Significant level of pain on a daily basis defined as clinically significant functional impairment (e.g., inability to perform household chores, prolonged standing, etc.)
 - ◆ Unremitting radicular pain to shoulder girdle and/or upper extremity resulting in disability
- Objective physical exam findings include **ANY** of the following:
 - ◆ Dermatomal sensory deficit
 - ◆ Motor deficit (e.g., biceps, triceps weakness)
 - ◆ Reflex changes
 - ◆ Shoulder abduction relief sign
 - ◆ Nerve root tension sign (e.g., Spurling's maneuver)
 - ◆ Unremitting radicular pain to shoulder girdle and/or upper extremity without concordant objective exam findings
- Less than clinically meaningful improvement with at least **TWO** of the following (unless contraindicated):
 - ◆ Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs for 6 weeks
 - ◆ Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician for 6 weeks
 - ◆ Epidural steroid injection(s) or selective nerve root block(s) performed at the same level as the requested surgery
- MRI/CT shows neural structure compression at the requested level(s) that is concordant with the individual's symptoms **and** physical exam findings and that is caused by **ANY** of the following:
 - ◆ Herniated disc(s) (retained disc material or a recurrent disc herniation)
 - ◆ Synovial cyst or arachnoid cyst
 - ◆ Central/lateral/foraminal stenosis
 - ◆ Osteophytes
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)

Myelopathy

- Subjective symptoms include **ANY** of the following:
 - ◆ Upper/lower extremity weakness, numbness, or pain
 - ◆ Fine motor dysfunction (buttoning, handwriting, clumsiness of hands)
 - ◆ Gait disturbance
 - ◆ New-onset bowel or bladder dysfunction
 - ◆ Frequent falls
- Objective physical exam findings include **ANY** of the following:
 - ◆ Grip and release test
 - ◆ Ataxic gait
 - ◆ Hyperreflexia
 - ◆ Hoffmann sign
 - ◆ Babinski sign
 - ◆ Tandem walking test demonstrating ataxia
 - ◆ Inverted brachial radial reflex
 - ◆ Increased muscle tone or spasticity
 - ◆ Clonus
 - ◆ Myelopathic hand
- MRI/CT shows findings that are concordant with the individual's symptoms **and** physical exam findings and that are caused by **EITHER** of the following:
 - ◆ Cervical spinal cord compression
 - ◆ Cervical spinal stenosis

CMM-605.3: Repeat Cervical Microdiscectomy at the Same Level

Repeat cervical microdiscectomy at the same level is considered **medically necessary** when performed for **EITHER** of the following conditions when **ALL** of the associated criteria have been met:

Radiculopathy

- Greater than 12 weeks since the initial primary cervical microdiscectomy
- Subjective symptoms include **BOTH** of the following:
 - ◆ Significant level of pain on a daily basis defined as clinically significant functional impairment (e.g., inability to perform household chores, prolonged standing, etc.)
 - ◆ Unremitting radicular pain to shoulder girdle and/or upper extremity resulting in disability

- Objective physical exam findings include **ANY** of the following:
 - ◆ Dermatomal sensory deficit
 - ◆ Motor deficit (e.g., biceps, triceps weakness)
 - ◆ Reflex changes
 - ◆ Shoulder abduction relief sign
 - ◆ Nerve root tension sign (e.g., Spurling's maneuver)
 - ◆ Unrelenting radicular pain to shoulder girdle and/or upper extremity without concordant objective physical exam findings
- Less than clinically meaningful improvement with at least **TWO** of the following (unless contraindicated):
 - ◆ Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs for 6 weeks
 - ◆ Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician for 6 weeks
 - ◆ Epidural steroid injection(s) or selective nerve root block(s) performed at the same level as the requested surgery
- Post-operative MRI /CT shows neural structure compression at the requested level(s) that is concordant with the individual's symptoms **and** physical exam findings and that is caused by **ANY** of the following:
 - ◆ Herniated disc(s) (retained disc material or a recurrent disc herniation)
 - ◆ Synovial cyst or arachnoid cyst
 - ◆ Central/lateral/foraminal stenosis
 - ◆ Osteophytes
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)

Myelopathy

- Subjective symptoms include **ANY** of the following:
 - ◆ Upper/lower extremity weakness, numbness, or pain
 - ◆ Fine motor dysfunction (buttoning, handwriting, clumsiness of hands)
 - ◆ Gait disturbance
 - ◆ New-onset bowel or bladder dysfunction
 - ◆ Frequent falls
- Objective physical exam findings include **ANY** of the following:
 - ◆ Grip and release test
 - ◆ Ataxic gait
 - ◆ Hyperreflexia
 - ◆ Hoffmann sign
 - ◆ Babinski sign
 - ◆ Tandem walking test demonstrating ataxia
 - ◆ Inverted brachial radial reflex
 - ◆ Increased muscle tone or spasticity
 - ◆ Clonus
 - ◆ Myelopathic hand
- Post-operative MRI /CT shows findings that are concordant with the individual's symptoms **and** physical exam findings and that are caused by **EITHER** of the following:
 - ◆ Cervical spinal cord compression
 - ◆ Cervical spinal stenosis

CMM-605.4: Non-Indications

Not Medically Necessary

- Cervical microdiscectomy performed without meeting the criteria in the **General Guidelines** (when applicable for urgent/emergent conditions) **and** the criteria in the applicable procedure-specific section (initial microdiscectomy or repeat microdiscectomy) is considered **not medically necessary**.
- Cervical microdiscectomy performed for **ANY** of the following sole indications is considered **not medically necessary**:
 - ◆ Annular tears
 - ◆ Concordant discography
 - ◆ MR Spectroscopy results
 - ◆ Degenerative disc disease

Experimental, Investigational, or Unproven (EIU)

- Percutaneous cervical discectomy (i.e., cervical discectomy performed with indirect visualization) is considered **experimental, investigational, or unproven (EIU)**.
- Endoscopic cervical discectomy is considered **experimental, investigational, or unproven (EIU)**.

Procedure (CPT®) Codes (CMM-605)

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.	
CPT®	Code Description/Definition
63020	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc; 1 interspace, cervical
+63035	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc; each additional interspace, cervical or lumbar (List separately in addition to code for primary procedure)
63040	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc, reexploration, single interspace; cervical
+63043	Laminotomy (hemilaminectomy), with decompression of nerve root(s), including partial facetectomy, foraminotomy and/or excision of herniated intervertebral disc, reexploration, single interspace; each additional cervical interspace (List separately in addition to code for primary procedure)
This list may not be all-inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.	

References (CMM-605)

1. Barton C, Kalakoti P, Bedard NA, Hendrickson NR, Saifi C, Pugely AJ. What Are the Costs of Cervical Radiculopathy Prior to Surgical Treatment? *Spine*. 2019;44(13):937-942. doi:10.1097/brs.0000000000002983.
2. Bond M, McIntosh G, Fisher C, et al. Treatment of Mild Cervical Myelopathy. *Spine (Phila Pa 1976)*. 2019;44(22):1606-1612. doi:10.1097/brs.0000000000003124.
3. Bono CM, Chiselli G, Gilbert TJ, et al. An evidence-based clinical guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *Spine J*. 2011;11(1):64-72. doi:10.1016/j.spinee.2010.10.023.
4. Boonstra AM, Schiphorst Preuper HR, Balk GA, Stewart RE. Cut-off points for mild, moderate, and severe pain on the visual analogue scale for pain in patients with chronic musculoskeletal pain. *Pain*. 2014;155(12):2545-2550. doi:10.1016/j.pain.2014.09.014.
5. Broekema AEH, Simões de Souza NF, Soer R, et al. Noninferiority of Posterior Cervical Foraminotomy vs Anterior Cervical Discectomy With Fusion for Procedural Success and Reduction in Arm Pain Among Patients With Cervical Radiculopathy at 1 Year. *JAMA Neurol*. 2023;80(1):40-40. doi:10.1001/jamaneurol.2022.4208
6. Celestre PC, Pazmiño PR, Mikhail MM, et al. Minimally invasive approaches to the cervical spine. *Orthop Clin North Am*. 2012;43(1):137-147. doi:10.1016/j.ocl.2011.08.007.
7. Cervical spine surgery. A guide to preoperative and postoperative patient care. AANN Clinical Practice Guideline Series [Internet] American Association of Neuroscience Nurses. 2007.
8. Changoor S, Farshchian J, Patel N, et al. Comparing outcomes between anterior cervical disc replacement (ACDR) and minimally invasive posterior cervical foraminotomy (MI-PCF) in the treatment of cervical radiculopathy. *Spine J*. 2024;24(5):800-806. doi:10.1016/j.spinee.2023.12.010.
9. Chatley A, Kumar R, Jain V, Behari S, Sahu R. Effect of spinal cord signal intensity changes on clinical outcome after surgery for cervical spondylotic myelopathy. *J Neurosurg Spine*. 2009;11(5):562-567. doi:10.3171/2009.6.spine091.
10. Cohen SP, Hanling S, Bicket MC, et al. Epidural steroid injections compared with gabapentin for lumbosacral radicular pain: multicenter randomized double blind comparative efficacy study. *BMJ*. 2015;350:h1748.. Published 2015 Apr 16. doi:10.1136/bmj.h1748.
11. Cunningham MR, Hershman S, Bendo J. Systematic review of cohort studies comparing surgical treatments for cervical spondylotic myelopathy. *Spine*. 2010;35(5):537-543.
12. Decker RC. Surgical treatment and outcomes of cervical radiculopathy. *Phys Med Rehabil Clin N Am*. 2011;22(1):179-191. doi:10.1016/j.pmr.2010.12.001.
13. de Rooij J, Harhangi B, Aukes H, Groeneweg G, Stronks D, Huygen F. The Effect of Percutaneous Nucleoplasty vs Anterior Discectomy in Patients with Cervical Radicular Pain due to a Single-Level Contained Soft-Disc Herniation: A Randomized Controlled Trial. *Pain Physician*. 2020;23(6):553-564.
14. Emami A, Coban D, Changoor S, et al. Comparing Mid-Term Outcomes Between ACDF and Minimally Invasive Posterior Cervical Foraminotomy in the Treatment of Cervical Radiculopathy. *Spine (Phila Pa 1976)*. 2021; Publish Ahead of Print. doi:10.1097/brs.0000000000004140.
15. Farshad M, Burgstaller JM, Held U, et al. Do preoperative corticosteroid injections increase the risk for infections or wound healing problems after spine surgery? *Spine*. 2018;43(15):1089-1094.
16. Fejer R, Jordan A, Hartvigsen J. Categorising the severity of neck pain: Establishment of cut-points for use in clinical and epidemiological research. *Pain*. 2005;119(1-3):176-182. doi:10.1016/j.pain.2005.09.033.
17. Grabowski G, Cornett CA, Kang JD. Esophageal and vertebral artery injuries during complex cervical spine surgery--avoidance and management. *Orthop Clin North Am*. 2012;43(1):63-74, viii.
18. Heary RF, Ryken TC, Matz PG, et al. Cervical laminoforaminotomy for the treatment of cervical degenerative radiculopathy. *J Neurosurg Spine*. 2009;11(2):198-202. doi:10.3171/2009.2.SPINE08722.
19. Hilton B, Tempest-Mitchell J, Davies B, Kotter M. Assessment of degenerative cervical myelopathy differs between specialists and may influence time to diagnosis and clinical outcomes. *PLoS ONE*. 2018;13(12). doi:10.1371/journal.pone.0207709.
20. Hsu WK. Advanced techniques in cervical spine surgery. *JBJS Am*. 2011;93(8):780-788.
21. Jagannathan J, Sherman JH, Szabo T, Shaffrey CI, Jane JA. The posterior cervical foraminotomy in the treatment of cervical disc/osteophyte disease: a single-surgeon experience with a minimum of 5 years' clinical and radiographic follow-up. *J Neurosurg Spine*. 2009;10(4):347-356.
22. Jayaram RH, Joo P, Gouzoulis MJ, Ratnasamy PP, Caruana D, Moore HE. Single-level Anterior Cervical discEctomy and Fusion has Lower Five-year Revisions than Posterior Cervical Foraminotomy in a Large National Cohort. *Spine*. 2023;48(18):1266-1271. doi:10.1097/brs.0000000000004754.
23. Kushchayev SV, Glushko T, Jarraya M, et al. ABCs of the degenerative spine. *Insights Imaging*. 2018;9(2):253-274. doi:10.1007/s13244-017-0584-z.
24. Lee BS, Nault R, Grabowski M, et al. Utility of repeat magnetic resonance imaging in surgical patients with lumbar stenosis without disc herniation. *Spine J*. 2019;19(2):191-198. doi:10.1016/j.spinee.2018.06.357.
25. Lutz S, Berk L, Chang E, et al. Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. *Int J Radiat Oncol Bio Phys*. 2011;79(4):965-976. doi:10.1016/j.ijrobp.2010.11.026.

26. Management of patients with oncologic or degenerative neurologic disorders. In: Smeltzer SC, Bare BG, Hinkle JL, Cheever KH, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 12th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2010:1975-2005.
27. Miller J, Gross A, D'Sylva J, et al. Manual therapy and exercise for neck pain: a systematic review. *Man Ther*. 2010;15(4):334-354. doi:10.1016/j.math.2010.02.007.
28. Molina CA, Gokaslan ZL, Sciubba DM. Diagnosis and management of metastatic cervical spine tumors. *Orthop Clin North Am*. 2012;43(1):75-87, viii-ix.
29. Mummaneni PV, Kaiser MG, Matz PG, et al. Cervical surgical techniques for the treatment of cervical spondylotic myelopathy. *J Neurosurg Spine*. 2009;11(2):130-141. doi:10.3171/2009.3.SPINE08728.
30. Mummaneni PV, Kaiser MG, Matz PG, et al. Preoperative patient selection with magnetic resonance imaging, computed tomography, and electroencephalography: does the test predict outcome after cervical surgery? *J Neurosurg Spine*. 2009;11(2):119-129. doi:10.3171/2009.3.SPINE08717.
31. Nikolaidis I, Fouyas IP, Sandercock PA, Statham PF. Surgery for cervical radiculopathy or myelopathy. *Cochrane Database Syst Rev*. 2010;2010(1):CD001466. Published 2010 Jan 20. doi:10.1002/14651858.CD001466.pub3.
32. Panagopoulos J, Hush J, Steffens D, Hancock MJ. Do MRI Findings Change Over a Period of Up to 1 Year in Patients With Low Back Pain and/or Sciatica? *Spine*. 2017;42(7):504-512. doi:10.1097/brs.0000000000001790.
33. Rao RD, Currier BL, Albert TJ, et al. Degenerative cervical spondylosis: clinical syndromes, pathogenesis, and management. *JBJS Am*. 2007;89(6):1360-1378. doi:10.2106/00004623-200706000-00026.
34. Ries ZG, Glassman SD, Vasilyev I, Metcalfe L, Carreon LY. Updated imaging does not affect revision rates in adults undergoing spine surgery for lumbar degenerative disease. *J Neurosurg Spine*. 2018;30(2):228-223. doi:10.3171/2018.8.spine18586.
35. Rueth N, Shaw D, Groth S, et al. Management of cervical esophageal injury after spinal surgery. *Ann Thoracic Surg*. 2010;90(4):1128-1133. doi:10.1016/j.athoracsur.2010.06.045.
36. Sahai N, Changoor S, Dunn C, et al. Minimally Invasive Posterior Cervical Foraminotomy as an Alternative to Anterior Cervical Discectomy and Fusion for Unilateral Cervical Radiculopathy. *Spine (Phila Pa 1976)*. 2019;44(24):1731-1739. doi:10.1097/brs.0000000000003156.
37. Shafshak TS, Elnemr R. The Visual Analogue Scale Versus Numerical Rating Scale in Measuring Pain Severity and Predicting Disability in Low Back Pain. *J Clin Rheumatol*. 2020;27(7):1. doi:10.1097/rhu.0000000000001320.
38. Shousha M, Boehm H. Surgical treatment of cervical spondylodiscitis: a review of 30 consecutive patients. *Spine*. 2012;37(1):E30-E36. doi:10.1097/BRS.0b013e31821bfdb2.
39. Tomaras CR, Blacklock JB, Parker WD, Harper RL. Outpatient surgical treatment of cervical radiculopathy. *J Neurosurg*. 1997;87(1):41-43.
40. Tracy JA, Bartleson JD. Cervical spondylotic myelopathy. *Neurologist*. 2010;16(3):176-187.
41. Yaksi A, Özgönenel L, Özgönenel B. The Efficiency of Gabapentin Therapy in Patients With Lumbar Spinal Stenosis. *Spine*. 2007;32(9):939-942. doi:10.1097/01.brs.0000261029.29170.e6.
42. Yin L, Zhang J, Wu Y, Li J, Yang Q. Increased signal intensity of spinal cord on T2W magnetic resonance imaging for cervical spondylotic myelopathy patients. *Medicine (Baltimore)*. 2020;99(49):e23098. doi:10.1097/md.00000000000023098.