

Cigna Medical Coverage Policies – Musculoskeletal Lumbar Total Disc Arthroplasty Guidelines

Effective November 1, 2024



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by eviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

CPT[®] (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT[®] five digit codes, nomenclature and other data are copyright 2024 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT[®] book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

©Copyright 2024 eviCore healthcare

CMM-610: Lumbar Total Disc Arthroplasty

CMM-610.1: General Guidelines

CMM-610.2: Initial Primary Lumbar Total Disc Arthroplasty

CMM-610.3: Failed Cervical Total Disc Arthroplasty Implant

CMM-610.4: Non-Indications

Procedure (CPT®) Codes (CMM-610)

References (CMM-610)

CMM-610.1: General Guidelines

Application of Guideline

- The determination of medical necessity for the performance of lumbar total disc arthroplasty is always made on a case-by-case basis.
- For additional timing and documentation requirements, see **CMM-600.1: Prior Authorization Requirements**.

CMM-610.2: Initial Primary Lumbar Total Disc Arthroplasty

Initial primary lumbar total disc arthroplasty is considered **medically necessary** when **ALL** of the following are met:

- Individual is age 18 to 60 years old
- Lumbar disc prosthesis approved by the FDA or for an FDA approved indication and in accordance with FDA labeling
- No planned simultaneous fusion (hybrid surgery) at an adjacent lumbar level
- The planned implant will be used in the reconstruction of a single-level lumbar disc at only one of the following lumbar levels: L3-4, L4-L5, or L5-S1
- Absence of facet ankylosis or severe facet degeneration at the operative level
- Plain X-rays **and** advanced diagnostic imaging studies (i.e., CT, MRI) confirm **ALL** of the following:
 - ◆ Presence of moderate to severe single-level disc degeneration at the operative level (between L3-L4, L4-L5, or L5-S1)
 - ◆ Absence of degenerative disc disease at more than one level (between L3-L4, L4-L5, or L5-S1)
 - ◆ Absence of degenerative disc disease above L3-L4
 - ◆ Subjective symptoms (concordant with single-level degenerative lumbar disc disease [DDD]) include significant level of pain on a daily basis defined as clinically significant functional impairment (e.g., inability to perform household chores, prolonged standing, etc.)

- Structured physician-supervised, multi-modal, nonoperative management of medical care with licensed healthcare professionals which includes **ALL** of the following:
 - ◆ Regularly scheduled appointments
 - ◆ Follow-up evaluation
 - ◆ Less than clinically meaningful improvement with **BOTH** of the following for at least 6 consecutive months (unless contraindicated):
 - Prescription strength analgesics, steroids, gabapentinoids, and/or NSAIDs
 - Provider-directed exercise program prescribed by a physical therapist, chiropractic provider, osteopathic or allopathic physician
- Absence of unmanaged significant mental and/or behavioral health disorders (e.g., major depressive disorder, chronic pain syndrome, secondary gain, opioid and alcohol use disorders)

CMM-610.3: Failed Lumbar Total Disc Arthroplasty Implant

For a revision of a failed lumbar total disc arthroplasty to a lumbar fusion, see **CMM-609.7: Lumbar Fusion (with or without Decompression) Following Failed Lumbar Disc Arthroplasty Surgery**

CMM-610.4: Non-Indications

Not Medically Necessary

- Lumbar total disc arthroplasty performed without meeting the requirements listed in the **General Guidelines** and the criteria in the procedure-specific section (initial disc arthroplasty) is considered **not medically necessary**.
- Lumbar total disc arthroplasty is considered **not medically necessary** when performed for **ANY** of the following:
 - ◆ Lumbar partial disc prosthetics
 - ◆ As an adjunct to the treatment of primary-central or far-lateral disc herniation
- Lumbar total disc arthroplasty is considered **not medically necessary** for **ANY** of the following **contraindications**:
 - ◆ Performed for the revision of a failed lumbar artificial total disc arthroplasty
 - ◆ The individual has osteopenia or osteoporosis (T-score < -1.0)
 - ◆ There is evidence on imaging studies of **ANY** of the following:
 - Degenerative or lytic spondylolisthesis >3mm
 - Lumbar spinal stenosis
 - Pars interarticularis defect with either spondylolysis or isthmic spondylolisthesis
 - Lumbar scoliosis (>11 degrees of sagittal plane deformity)
 - Spinal fracture
 - Infection
 - Presence of tumor or active infection at the site of implantation
 - Lumbar nerve root compression or bony spinal stenosis
 - Preoperative remaining disc height <3mm

- Mid-sagittal stenosis of <8mm (by MRI)
- ◆ History of ankylosing spondylitis, rheumatoid arthritis, lupus, or other autoimmune disorder
- ◆ Allergy or sensitivity to implant materials
- ◆ Isolated radicular compression syndromes especially due to lumbar disc herniation
- ◆ Involved vertebral endplate is dimensionally smaller than the approximate dimensions of the implant in anterior/posterior width and lateral width
- ◆ Clinically compromised vertebral bodies at the affected level due to current or past trauma

Procedure (CPT®) Codes (CMM-610)

This guideline relates to the CPT® code set below. Codes are displayed for informational purposes only. Any given code's inclusion on this list does not necessarily indicate prior authorization is required.

CPT®	Code Description/Definitions
22857	Total disc arthroplasty (artificial disc), anterior approach, including discectomy to prepare interspace (other than for decompression), single interspace, lumbar
22862	Revision including replacement of total disc arthroplasty (artificial disc), anterior approach, single interspace; lumbar
22865	Removal of total disc arthroplasty (artificial disc), anterior approach, single interspace; lumbar
+0164T	Removal of total disc arthroplasty, (artificial disc), anterior approach, each additional interspace, lumbar (List separately in addition to code for primary procedure)
+0165T	Revision including replacement of total disc arthroplasty (artificial disc), anterior approach, each additional interspace, lumbar (List separately in addition to code for primary procedure)

This list may not be all-inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules

References (CMM-610)

1. Activ-L™ Artificial Disc Treatment of Degenerative Disc Disease in the Treatment of Degenerative Disc Disease - Full Text View - ClinicalTrials.gov. Clinicaltrials.gov. <https://clinicaltrials.gov/ct2/show/NCT00589797>. Published 2020.
2. Ahrens M, Tsantrizos A, Donkersloot P, et al. Nucleus replacement with the DASCOR disc arthroplasty device: interim two-year efficacy and safety results from two prospective, non-randomized multicenter European studies. *Spine (Phila Pa 1976)*. 2009;34(13):1376-1384.
3. Balsano M, Zachos A, Ruggiu A, Barca F, Tranquilli-Leali P, Doria C. Nucleus disc arthroplasty with the NUBAC™ device: 2-year clinical experience. *Eur Spine J*. 2011;20(Suppl 1):S36-S40. Epub 2011 Mar 18.
4. Bertagnoli R, Karg A, Voigt S. Lumbar partial disc replacement. *Orthop Clin N Am*. 2005;36:341-347.
5. Bertagnoli R, Yue JJ, Fenk-Mayer A, Eerulker J, Emerson JW. Treatment of symptomatic adjacent-segment degeneration after lumbar fusion with total disc arthroplasty by using the ProDisc prosthesis: a prospective study with 2-year minimum follow up. *J Neurosurg Spine*. 2006;4(2):91-97. doi:10.3171/spi.2006.4.2.91.
6. Bertagnoli R, Yue JJ, Kershaw T, et al. Lumbar total disc arthroplasty utilizing the ProDisc prosthesis in smokers versus nonsmokers. *Spine*. 2006;31(9):992-997. doi:10.1097/01.brs.0000214970.07626.68.
7. Bertagnoli R, Yue JJ, Shah RV, et al. The treatment of disabling single-level lumbar discogenic low back pain with total disc arthroplasty utilizing the ProDisc Prosthesis. *Spine*. 2005;30(19):2230-2236.
8. Blumenthal S, McAfee PC, Guyer RD, Hochschuler, Geisler RD, Holt RT. A prospective, randomized, multicenter Food and Drug Administration investigational device exemptions study of lumbar total disc replacement with the Charité™ artificial disc versus lumbar fusion. Part I: evaluation of clinical outcomes. *Spine*. 2005;30(14):1565-1575.
9. Blumenthal SL, Ohnmeiss DD, Guyer RD, Hochschuler SH. Prospective study evaluating total disc replacement: preliminary results. *J Spinal Disord Tech*. 2003;16(5):450-454.
10. Blumenthal SL, Ohnmeiss DD, Guyer R, et al. Artificial intervertebral disks and beyond: a North American Spine Society Annual Meeting Symposium. *Spine J*. 2002;2(6):460-463.
11. Boden SD, Balderston RA, Heller J, Hanley EN Jr., Zigler JE. An AOA critical issue. Disc Replacements: This Time Will We Really Cure Low-Back and Neck Pain? *J Bone Joint Surg Am*. 2004;86(2):411-422.
12. Boonstra AM, Schiphorst Preuper HR, Balk GA, Stewart RE. Cut-off points for mild, moderate, and severe pain on the visual analogue scale for pain in patients with chronic musculoskeletal pain. *Pain*. 2014;155(12):2545-2550. doi:10.1016/j.pain.2014.09.014.
13. Boswell MV, Shah RV, Everett CR, et al. Interventional techniques in the management of chronic spinal pain: evidence-based practice guidelines. *Pain Phys*. 2005;8(1):1-47. Updated 2005.
14. Bree Collaborative Spine/Low Back Pain Topic – Report and Recommendations. November 2013.
15. Chung SS, Lee CS, Kang CS. Lumbar total disc replacement using ProDisc II: A prospective study with a 2-year minimum follow-up. *J Spinal Disord Tech*. 2006;19:411-415.
16. Cinotti G, Thierry D, Postacchini F. Results of disc prosthesis after a minimum follow-up period of 2 years. *Spine*. 1996;21(8):995-1000.
17. Cohen SP, Hanling S, Bicket MC, et al. Epidural steroid injections compared with gabapentin for lumbosacral radicular pain: multicenter randomized double blind comparative efficacy study. *BMJ*. 2015;350:h1748. Published 2015 Apr 16. doi:10.1136/bmj.h1748.
18. Cunningham BW, McAfee PC, Geisler FH, et al. Distribution of in vivo and in vitro range of motion following 1-level arthroplasty with the CHARITE artificial disc compared with fusion. *J Neurosurg Spine*. 2008;8(1):7-12.
19. David T. Long-term results of one-level lumbar arthroplasty: minimum 10-year followup of the Charite artificial disc in 106 patients. *Spine*. 2007;32:661-666.
20. Davidson, M, Keating, JL. A Comparison of Five Low Back Disability Questionnaires: Reliability and Responsiveness: *Phys Ther*. 2002;82 8-24.
21. Di Silvestre M, Bakaloudis G, Lolli F, Vommaro F, Parisini P. Two-level total lumbar disc replacement. *Eur Spine J*. 2009;18(Suppl 1):64-70. Epub 2009 Apr 28.
22. Delamarter R, Zigler JE, Balderston RA, Cammisa FP, Goldstein JA, Spivak JM. Prospective, randomized, multicenter Food and Drug Administration investigational device exemption study of the ProDisc-L total disc replacement compared with circumferential arthrodesis for the treatment of two-level lumbar degenerative disc disease: results at twenty-four months. *J Bone Joint Surg Am*. 2011;93(8):705-715. doi:10.2106/JBJS.I.00680. Epub 2011 Mar 11.
23. Donnally C, Patel P, Canseco J, et al. Current incidence of adjacent segment pathology following lumbar fusion versus motion-preserving procedures: a systematic review and meta-analysis of recent projections. *Spine J*. 2020;20(10):1554-1565. doi:10.1016/j.spinee.2020.05.100.
24. ECRI Institute. *Artificial intervertebral disc replacement (AIDR) for lumbar degenerative disc disease (DDD)*. Emerging Technology Evidence Report. Plymouth Meeting (PA): ECRI Institute; 2009a.
25. FDA *Summary of Safety and Effectiveness Data Prodisc-L Total Disc Replacement*.
26. Fekete TF, Porchet F. Overview of disc arthroplasty-past, present and future. *Acta Neurochir (Wien)*. 2010;152(3):393-404.

27. Fischer, CR, Ducoffe, AR, Errico, TJ. Review Article: Posterior Lumbar Fusion: Choice of Approach and Adjunct Techniques. *J Am Acad Orthop Surg*. 2014;22:503-511.
28. Freeman BJC, Davenport J. Total disc replacement in the lumbar spine: a systematic review of the literature. *Eur Spine J*. 2006;15(Suppl. 3):S439-S447.
29. Geisler FH, Guyer RD, Blumenthal SL, McAfee PC, Cappuccino A, Bitan F, Regan JJ. Effect of previous surgery on clinical outcome following 1-level lumbar arthroplasty. *J Neurosurg Spine*. 2008;8(2):108-114. doi:10.3171/SPI/2008/8/2/108.
30. Geisler FH, Guyer RD, Blumenthal SL, McAfee PC, Cappuccino A, Bitan F, Regan JJ. Patient selection for lumbar arthroplasty and arthrodesis: the effect of revision surgery in a controlled, multicenter, randomized study. *J Neurosurg Spine*. 2008;8(1):13-16. doi:10.3171/SPI-08/01/013.
31. German JW, Foley KT. Disc arthroplasty in the management of the painful lumbar motion segment. *Spine*. 2005;30(16S):S60-S67.
32. Griffith SL, Shelokov AP, Buttner-Janzen K, LeMaire, JP, Zeegers WS. A multicenter retrospective study of the clinical results of the LINK® SB Charité intervertebral prosthesis: the initial European experience. *Spine*. 1994;19:1842-1849.
33. Guyer RD, Geisler FH, Blumenthal SL, et al. Effect of age on clinical and radiographic outcomes and adverse events following 1 level lumbar arthroplasty after a minimum 2 year follow-up. *J Neurosurg Spine*. 2008;8(2):101-107. doi:10.3171/SPI/2008/8/2/101.
34. Guyer RD, McAfee PC, Banco RJ, et al. Prospective, randomized, multicenter Food and Drug Administration investigational device exemption study of lumbar total disc replacement with the CHARITE artificial disc versus lumbar fusion: Five-year follow-up. *Spine J*. 2009;9(5):374-386. doi:10.1016/j.spinee.2008.08.007.
35. Hannibal M, Thomas DJ, Low J, Hsu KY, Zucherman J. ProDisc-L total disc replacement: a comparison of 1-level versus 2-level arthroplasty patients with a minimum 2-year follow-up. *Spine*. 2007;32(21):2322-2326.
36. Huang RC, Girardi FP, Lim MR, Cammisa FP. Advantages and disadvantages of nonfusion technology in spine surgery. *Orthop Clin N Am*. 2005;36:263-269.
37. Institutes for Clinical Systems Improvement (ICSI). Lumbar artificial intervertebral discs. Bloomington, MN: Institute for Clinical Systems Improvement (ICSI); 2005 Dec.
38. Jacobs WC, Van der Gaag NA, Kruijff MC, et al. Total disc replacement for chronic discogenic low-back pain: A Cochrane Review. *Spine (Phila Pa 1976)*. 2013;38(1):24-36. doi:10.1097/BRS.0b013e3182741b21.
39. Katsimihias M, Bailey CS, Issa K, Fleming J, Rosas-Arellano P, Bailey SI, Gurr KR. Prospective clinical and radiographic results of CHARITÉ III artificial total disc arthroplasty at 2- to 7-year follow-up: a Canadian experience. *Can J Surg*. 2010;53(6):408-4145.
40. Kostuik JP. Intervertebral disk replacement. Experimental study. *Clin Orthop Relat Res*. 1997;337:27-41.
41. Koutsogiannis P, Khan S, Phillips F, et al. A cross-sectional analysis of 284 complications for lumbar disc replacements from medical device reports maintained by the United States Food and Drug Administration. *Spine J*. 2022;22(2):278-285. doi: 10.1016/j.spinee.2021.08.001.
42. Kuhns BD, Louk S, Buchanan C, et al. Sensitivity of Magnetic Resonance Imaging in the Diagnosis of Mobile and Non-Mobile L4-5 Degenerative Spondylolisthesis. *Spine J*. 2015;15(9):1956-1962. doi:10.1016/j.spinee.2014.08.006.
43. Kurtz SM, Pelozo J, Siskey R, Villarraga ML. Analysis of a retrieved polyethylene total disc replacement component. *Spine J*. 2005;5(3):344-350.
44. Kurtz SM, van Ooij A, Ross R, Malefit JdW, Pelozo J, Ciccarelli L, Villarraga ML. Polyethylene wear and rim fracture in total disc arthroplasty. *Spine J*. 2007;7:12-21.
45. Leahy M, Zigler JE, Ohnmeiss DD, Rashbaum RF, Sachs BL. Comparison of results of total disc replacement in postdiscectomy patients versus patients with no previous lumbar surgery. *Spine*. 2008;33(15):1690-1693; discussion 1694-1695.
46. Leary SP, Regan JJ, Lanman TH, Wagner WH. Revision and Explantation Strategies Involving the Lumbar Artificial Disc Replacement. *Spine*. 2007;32(9):1001-1011.
47. Le Huec [a] J-C, Basso Y, Aunoble S, Friesem T, Bruno MB. Influence of Facet and Posterior Muscle Degeneration on Clinical Results of Lumbar Total Disc Replacement: Two-Year Follow-Up. *J Spinal Disord Tech*. 2005;18(3):219-223.
48. Le Huec [b] JC, Basso Y, Mathews H, Mehdod A, Aunoble S, Friesem T, Zdeblick T. The effect of single-level, total disc arthroplasty on sagittal balance parameters: a prospective study. *Eur Spine J*. 2005;14:480-486.
49. Lee BS, Nault R, Grabowski M, et al. Utility of repeat magnetic resonance imaging in surgical patients with lumbar stenosis without disc herniation. *Spine J*. 2019;19(2):191-198. doi:10.1016/j.spinee.2018.06.357.
50. Leivseth G, Braaten S, Frobin W, Brinckmann P. Mobility of lumbar segments instrumented with a ProDisc II Prosthesis: A two-year follow-up study. *Spine*. 2006;31(15):1726-1733.
51. LeMaire JP, Carrier H, Sari Ali E-H, Skalli W, Lavaste F. Clinical and radiological outcomes with the Charité™ artificial disc: a 10-year minimum follow-up. *J Spinal Disord Tech*. 2005;18(4):353-359.
52. Lurie JD, Tosteson TD, Tosteson AN, Zhao W, Morgan TS, Abdu WA, Herkowitz H, Weinstein, JN. Surgical versus Nonoperative Treatment for Lumbar Disc Herniation. *Spine*. 39(1):3-16.

53. Markwalder TM, Wenger M, Marbacher S. A 6.5-year follow-up of 14 patients who underwent ProDisc total disc arthroplasty for combined long-standing degenerative lumbar disc disease and recent disc herniation. *J Clin Neurosci*. 2011;18(12):1677-1681.
54. Marshman LA, Friesem T, Rampersaud YR, Le Huec JC, Krishna M. Subsidence and malplacement with the Oblique Maverick Lumbar Disc Arthroplasty: technical note. *Spine J*. 2008;8(4):650-655. Epub 2007 May 22.
55. Martin CW and the Work Comp Board (WCB) Evidence Based Practice Group. Artificial cervical and lumbar disc implants: a review of the literature. Updated Apr 2005.
56. McAfee PC, Geisler FH, Saiedy SS, et al. Revisability of the CHARITÉ artificial disc replacement. *Spine*. 2006;31(11):1217-1226.
57. McAfee PC, Cunningham B, Holsapple G, Adams K, Blumenthal S, Guyer RD. A prospective, randomized, multicenter Food and Drug device exemption study of lumbar total disc replacement with the Charité™ artificial disc versus lumbar fusion. Part II: Evaluation of radiographic outcomes and correlations of surgical technique accuracy with clinical outcomes. *Spine*. 2005;30(14):1576-1583.
58. McAfee PC, Fedder IL, Saiedy S, Shucosky EM, Cunningham BW. SB Charité disc replacement: report of 60 prospective randomized cases in a U.S. center. *J Spinal Disord Tech*. 2003;16(4):424-433.
59. Mirza SK, Deyo RA. Systematic Review of Randomized Trials Comparing Lumbar Fusion Surgery to Nonoperative Care for Treatment of Chronic Back Pain. *Spine*. 2007;32(7):816-823.
60. National Institutes for Health and Clinical Excellence (NICE). *Prosthetic intervertebral disc replacement in the cervical spine*. Interventional Procedure Guidance 341. May 2010.
61. North American Spine Society (NASS) Evidenced-Based Clinical Guidelines for Multidisciplinary Spine Care. *Diagnosis and Treatment of Degenerative Lumbar Spinal Stenosis*. Revised 2014. Burr Ridge, IL. North American Spine Society (NASS). Available at: <https://www.spine.org>.
62. North American Spine Society (NASS) Coverage Policy Recommendations. *Lumbar Artificial Disc Replacement*. 2019. Burr Ridge, IL North American Spine Society (NASS). Available at: <https://www.spine.org>.
63. Ontario Health Technology Assessment Committee (OHTAC). OHTAC Recommendation: Updated Health Technology Policy Assessment (HPTA) on Artificial Disc Replacement for Lumbar and Cervical Degenerative Disc Disease. Updated Apr 2006. *Ont Health Technol Assess Ser*. 2006;6(10):1-98.
64. Panagopoulos J, Hush J, Steffens D, Hancock MJ. Do MRI Findings Change Over a Period of Up to 1 Year in Patients With Low Back Pain and/or Sciatica? *Spine*. 2017;42(7):504-512. doi:10.1097/brs.0000000000001790.
65. Panjabi M, Malcolmson G, Teng E, Tominaga Y, Henderson G, Serhan H. Hybrid Testing of Lumbar Discs Versus Fusions. *Spine*. 2007;32(9):959-966.
66. Park CK, Ryu KS, Jee WH. Degenerative changes of discs and facet joints in lumbar total disc replacement using ProDisc II: minimum two-year follow-up. *Spine*. 2008;33(16):1755-1761.
67. Park CK, Ryu KS, Lee KY, Lee HJ. Clinical Outcome of Lumbar Total Disc Replacement Using ProDisc-L® in Degenerative Disc Disease: Minimum 5-year Follow-up Results at a Single Institute. *Spine (Phila Pa 1976)*. 2011;37(8):672-677.
68. Regan JJ. Clinical results of Charité lumbar total disc replacement. *Orthop Clin N Am*. 2005;36(3):323-340. doi:10.1016/j.ocl.2005.03.005.
69. Park SJ, Lee CS, Chung SS, Lee KH, Kim WS, Lee JY. Long-Term Outcomes Following Lumbar Total Disc Replacement Using ProDisc-II: Average 10-Year Follow-Up at a Single Institution. *Spine*. 2016;41(11):971-977.
70. Punt IM, Vissor VM, van Rhijn LW, et al. Complications and reoperations of the SB Charite lumbar disc prosthesis: experience in 75 patients. *Eur Spine J*. 2008;17(1):36-43.
71. Radcliff K, Spivak J, Darden B 2nd, Janssen M, Bernard T, Zigler J. Five-Year Reoperation Rates of 2-Level Lumbar Total Disk Replacement Versus Fusion: Results of a Prospective, Randomized Clinical Trial. *Clin Spine Surg*. 2018;31(1):37-42. doi:10.1097/BSD.0000000000000476.
72. Rasouli A, Cuellar JM, Kanim L, Delamarter R. Multiple-Level Lumbar Total Disk Replacement: A Prospective Clinical and Radiographic Analysis of Motion Preservation at 24-72 Months. *Clin Spine Surg*. 2019;32(1):38-42. doi:10.1097/BSD.0000000000000704.
73. Ries ZG, Glassman SD, Vasilyev I, Metcalfe L, Carreon LY. Updated imaging does not affect revision rates in adults undergoing spine surgery for lumbar degenerative disease. *J Neurosurg Spine*. Published online Nov 2018. 2019;30(2):228-223. doi:10.3171/2018.8.spine18586.66.
74. Ross R, Mirza AH, Norris HE, et al. Survival and clinical outcome of SB Charite III disc replacement for back pain. *J Bone Joint Surg Br*. 2007;89(6):785-789.
75. SariAli el-H, LeMaire JP, Pascal-Mousselard H, Carrier H, Skalli W. In vivo study of the kinematics in axial rotation of the lumbar spine after total intervertebral disc replacement: long-term results: a 10-14 years follow-up evaluation. *Eur Spine J*. 2006;15(10):1501-1510. doi:10.1007/s00586-005-0016-5.
76. Scott-Young MN, Lee MJ, Nielsen DE, Magno CL, Kimlin KR, Mitchell EO. Clinical and Radiological Mid-Term Outcomes of Lumbar Single-Level Total Disc Replacement. *Spine (Phila Pa 1976)*. 2018;43(2):105-113. doi:10.1097/BRS.0b013e3182345aa2.

77. Scott-Young M, Lee S, Nielsen D, Rathbone E, Rackham M, Hing W. Comparison of Mid- to Long-term Follow-up of Patient-reported Outcomes Measures After Single-level Lumbar Total Disc Arthroplasty, Multi-level Lumbar Total Disc Arthroplasty, and the Lumbar Hybrid Procedure for the Treatment of Degenerative Disc Disease. *Spine (Phila Pa 1976)*. 2021;47(5):377-386. doi:10.1097/brs.0000000000004253.
78. Shafshak TS, Elnemr R. The Visual Analogue Scale Versus Numerical Rating Scale in Measuring Pain Severity and Predicting Disability in Low Back Pain. *J Clin Rheumatol*. 2020;27(7):1. doi:10.1097/rhu.0000000000001320.
79. Sharan AD, Goldstein JA. Cervical artificial disc replacement technologies. Updated Aug 2015.
80. Shim CS, Lee S-H, Shin H-- Versus ProDisc: A Comparative Study of a Minimum 3-Year Follow-up. *Spine*. 2007;32(9):1012-1018.
81. Siepe CJ, Heider F, Haas E, et al. Influence of lumbar intervertebral disc degeneration on the outcome of total lumbar disc replacement: a prospective clinical, histological, X-ray and MRI investigation. *Eur Spine J*. 2012;21(11):2287-2299. doi:10.1007/s00586-012-2342-8.
82. Siepe CJ, Korge A, Grochulla F, Mehren C, Mayer HM. Analysis of post-operative pain patterns following total lumbar disc replacement: results from fluoroscopically guided spine infiltrations. *Eur Spine J*. 2008;17(1):44-56.
83. Siepe CJ, Mayer HM, Heinz-Leisenheimer M, Korge A. Total lumbar disc replacement: different results for different levels. *Spine*. 2007;32(7):782-790.
84. Siepe CJ, Mayer M, Wiechert K, Korge A. Clinical results of total lumbar disc replacement with ProDisc I: Three-year results for different indications. *Spine*. 2006;31:1923-1932.
85. Siepe CJ, Tepass A, Hitzl W, et al. Dynamics of improvement following total lumbar disc replacement: is the outcome predictable? *Spine (Phila Pa 1976)*. 2009;34(23):2579-2586. doi:10.1097/BRS.0b013e3181b612bd.
86. Siepe CJ, Zelenkov P, Sauri-Barraza JC, et al. The fate of facet joint and adjacent level disc degeneration following total lumbar disc replacement: a prospective clinical, x-ray, and magnetic resonance imaging investigation. *Spine (Phila Pa 1976)*. 2010;35(22):1991-2003.
87. Tropiano P, Huang RC, Giradi FP, et al. Lumbar total disc replacement: surgical technique. *J Bone Joint Surg Am*. 2006;88A(Suppl 1, Part 1):50-64. doi:10.2106/JBJS.E.01066.
88. Total disc replacement for chronic low back pain: background and a systematic review of the literature. The Cochrane Review. In: The Cochrane Library, Issue 12, 2005. Chichester, UK: John Wiley & Sons, Ltd.; 2005. Oxford: Update software.
89. Uschold TD, Fusco D, Germain R, Tumialan LM, Chang SW. Cervical and Lumbar Spinal Arthroplasty: Clinical Review. *AJNR Am J Neuroradiol*. 2012;33(9):1631-1641. doi:10.3174/ajnr.A2758.
90. U.S. Food and Drug Administration. *Orthopaedic and Rehabilitation Devices Panel of the Medical Devices Advisory Committee*. Nov 2002. Updated Nov 2002.
91. U.S. Food and Drug Administration. Center for Devices and Radiological Health (CDRH). *New Device Approval. ProDisc®-L Total Disc Replacement – P050010*. Updated Aug 2006.
92. U.S. Food and Drug Administration. Center for Devices and Radiological Health (CDRH). *New Device Approval. Charite™ Artificial Disc- P040006*. Issued October 26, 2004. Updated March 9, 2005.
93. van den Eerenbeemt KD, Ostelo RW, van Royen BJ, Peul WC, van Tulder MW. Total disc replacement surgery for symptomatic degenerative lumbar disc disease: a systematic review of the literature. *Eur Spine J*. 2010;19(8):1262-1280. doi:10.1007/s00586-010-1445-3.
94. van Ooij A, Kurtz SM, Stessels F, et al. Polyethylene wear debris and long term clinical failure of the Charite disc prosthesis: a study of 4 patients. *Spine*. 2007;32(2):223-229.
95. Wagner WH, Regan JJ, Leary SP, et al. Access strategies for revision or explanation of the Char. *J Vasc Surg*. 2006;44:1266-1272.
96. van Ooij A, Oner FC, Verbout AJ. Complications of artificial disc replacement: a report of 27 patients with the SB Charite disc. *J Spinal Disord Tech*. 2003;16(4):369-383.
97. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical Compared with Nonoperative Treatment for Lumbar Degenerative Spondylolisthesis. Four –Year Results in the Spine Patient Outcomes Research Trial (SPORT) Randomized and Observational Cohorts. *J Bone Joint Surg Am*. 2009;91:1295-1304.
98. Yaksi A, Özgönenel L, Özgönenel B. The Efficiency of Gabapentin Therapy in Patients With Lumbar Spinal Stenosis. *Spine*. 2007;32(9):939-942. doi:10.1097/01.brs.0000261029.29170.e6.
99. Yaszay B, Bendo JA, Goldstein JA, Quirno M, Spivak JM, Errico TJ. Effect of intervertebral disc height on postoperative motion and outcomes after ProDisc-L lumbar disc replacement. *Spine*. 2008;33(5):508-512; discussion 513.
100. Yue J, Garcia R, Blumenthal S, et al. Five-year Results of a Randomized Controlled Trial for Lumbar Artificial Discs in Single-level Degenerative Disc Disease. *Spine (Phila Pa 1976)*. 2019;44(24):1685-1696. doi:10.1097/brs.0000000000003171.
101. Yue J, Zhang K, Bai HX, et al. A comparison of patients who have undergone 1-Level versus 2-Level ProDisc arthroplasty: a prospective study with minimum of 5-year follow-up. *Spine (Phila Pa 1976)*. 2013;38(14):1194-1198. doi:10.1097/BRS.0b013e31828b31e8.
102. Zeller JL. Artificial spinal disk superior to fusion for treating degenerative disk disease. *JAMA*. 2006;296(22):2665-2666.

103. Zigler JE, Delamarter RB. Five-year results of the prospective, randomized, multicenter, food and drug administration investigational device exemption study of the ProDisc-L total disc replacement versus circumferential arthrodesis for the treatment of single-level degenerative disc disease. *J Neurosurg Spine*. 2012;17:493-501.
104. Zigler J, Delamater R, Spivak JM, et al. Results of the prospective randomized, multicenter food and drug administration investigational device exemption study of the ProDisc®-L total disc replacement versus circumferential fusion for the treatment of 1-level degenerative disc disease. *Spine*. 2007;32(11):1155-1162.
105. Sigler JE, Glenn J, Delamarter RB. Five year adjacent level degenerative changes in patients with single level disease treated using lumbar total disc replacement with ProDisc®-L versus circumferential fusion. *J Neurosurg Spine*. 2012;17(6):504-511. doi:10.3171/2012.9.SPINE11717.
106. Zindrick MR, Tzermiadianos MN, Voronov LI, Lorenz M, Hadjipavlou A. An evidence-based medicine approach in determining factors that may affect outcome in lumbar total disc replacement. *Spine*. 2008;33(11):1262-1269.