

Sexually Transmitted and Other Reproductive Tract Infection Testing

MOL.CS.106.Z
v2.0.2024

Introduction

Molecular testing for sexually transmitted infections (STIs) and other infections of the reproductive tract (including bacterial vaginosis) is addressed by this guideline.

Procedures addressed

The inclusion of any procedure code in this table does not imply that the code is under management or requires prior authorization. Refer to the specific Health Plan's procedure code list for management requirements.

Procedures addressed by this guideline	Procedure codes
Abbott Alinity m STI Assay	0402U
Aptima BV Assay	81513
BD MAX Vaginal Panel	81514
Bridge Women's Health Infectious Disease Detection Test	0330U
Candida species	87480 87481 87482
Candida species panel (C. albicans, C. glabrata, C. parapsilosis, C. kruseii, C. tropicalis, and C. auris), amplified probe technique with qualitative report of the presence or absence of each species	0068U
Chlamydia trachomatis	87490 87491 87492
Gardnerella	87510 87511 87512

Procedures addressed by this guideline	Procedure codes
Herpes simplex virus	87528 87529 87530
Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; direct probe technique, each organism	87797
Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; amplified probe technique, each organism	87798
Infectious agent detection by nucleic acid (DNA or RNA), not otherwise specified; quantification, each organism	87799
Mycoplasma genitalium, amplified probe technique	87563
Neisseria gonorrhoeae	87590 87591 87592
Trichomonas vaginalis	87660 87661
Xpert CT/NG	0353U
Xpert Xpress MVP	0352U

Criteria

Introduction

Requests for molecular testing for sexually transmitted infections (STIs) and other infections of the reproductive tract (including bacterial vaginosis) are reviewed using these criteria.

Please refer to the guideline, *Urinary Tract Infection Molecular Testing* for testing requested for urinary tract infections (UTIs), as this testing is not addressed here. For testing of other STIs, please refer to the guidelines *Human Immunodeficiency Virus Laboratory Testing* and *Human Papillomavirus (HPV) Molecular Testing*, as this testing is not addressed here. For any tests not addressed by an eviCore test-specific guideline, please refer to the guidelines, *Identification of Microorganisms Using Nucleic Acid Probes* and *Infectious Disease Laboratory Testing*.

Multiple Organism Detection Panels

Paragraph

Test name	Procedure code	Medical Necessity
Abbott Alinity m STI Assay	0402U	Procedure code is medically necessary when criteria are met.

Medical necessity requirements:

When billed with a single panel code, testing for the detection of multiple organisms associated with sexually transmitted infections (STIs) and other infections of the reproductive tract is considered medically necessary for individuals with clinical indications for all organisms included on the panel, as outlined in the criteria sections below.

More than one type of molecular test for the same organism for the same date of service is not medically necessary.

Bacterial Vaginosis (BV)

Test name	Procedure code	Medical Necessity
Aptima Bacterial Vaginosis Assay	81513	Procedure code is medically necessary when criteria are met
BD MAX Vaginal Panel	81514	Procedure code is medically necessary when criteria are met
Bridge Women's Health Infectious Disease Detection Test	0330U	Procedure code is experimental, investigational, or unproven
Miscellaneous Infectious Agent Detection, Direct Molecular Method	87797	Procedure code is experimental, investigational, or unproven for indications addressed by this guideline
Miscellaneous Infectious Agent Detection, Amplified Molecular Method	87798	Procedure code is experimental, investigational, or unproven for indications addressed by this guideline

Test name	Procedure code	Medical Necessity
Miscellaneous Infectious Agent Detection, Quantitative Molecular Method	87799	Procedure code is experimental, investigational, or unproven for indications addressed by this guideline
Xpert Xpress MVP	0352U	Procedure code is experimental, investigational, or unproven

Medical necessity requirements:

Testing for bacterial vaginosis (BV) using Aptima Bacterial Vaginosis Assay (CPT 81513) or BD MAX Vaginal Panel (CPT 81514) is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Routine screening for BV in asymptomatic individuals is not medically necessary

Symptomatic individuals:

- Molecular testing for BV is medically necessary when microscopy and culture are not available or unable to provide a diagnosis.

More than one type of molecular test for the same organism for the same date of service is not medically necessary.

Exclusions:

Clinical utility of testing for bacterial vaginosis using other technologies has not been demonstrated. These tests are considered Experimental, Investigational, or Unproven.

This includes but is not limited to, individual organism-specific procedure codes or agent not otherwise specified (NOS) procedure codes billed for this purpose, the proprietary laboratory analysis (PLA) codes listed as experimental, investigational, or unproven in the table above, as well as other BV panels, such as:

- NuSwab®, Vaginitis Plus (VG+) (A vaginae, BVAB-2, Megasphaera Type 1)
- OneSwab® (A vaginae, Megasphaera Type 1 and 2, BVAB-2)
- SureSwab® Vaginosis, Vaginitis Plus (G vaginalis, A vaginae, Megasphaera species)

Candida species

Test name	Procedure code	Medical Necessity
Candida Detection, Direct Method	87480	Procedure code is medically necessary when criteria are met.
Candida Detection, Amplified Method	87481	Procedure code is medically necessary when criteria are met.
Candida Detection, Quantification Method	87482	Procedure code is experimental, investigational, or unproven.
Candida Species Panel, Amplified Method	0068U	Procedure code is experimental, investigational, or unproven.

Direct or amplified Candida detection**Medical necessity requirements:**

Nucleic acid amplification testing (NAAT) for *Candida albicans* through either direct (CPT 87480) or amplified (CPT 87481) probe studies is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Evidence does not support routine screening for *Candida* species in asymptomatic pregnant women, non-pregnant women, or men unless HIV status is positive.

Symptomatic individuals:

- Molecular testing for *Candida* is medically necessary when microscopy and culture are not available or unable to provide a diagnosis.

More than one type of molecular test for the same organism for the same date of service is not medically necessary (e.g., 87480 and 87481 may not be billed together).

It should only be necessary to test one site. Therefore, only one unit per date of service is medically necessary.

Subtyping for *Candida glabrata* and other non-*albicans* *Candida* species is not routinely medically necessary. Exceptions may be considered if complicated vulvovaginal candidiasis (VVC) is diagnosed. Complicated VVC may include:

- Recurrent VVC (defined as 3 or more episodes of symptomatic VVC within 1 year), or
- Severe VVC (i.e. extensive vulvar erythema, edema, excoriation, and fissure formation)
- VVC in patients with immunosuppression or diabetes mellitus

Quantitative Candida testing

Medical necessity requirements:

Clinical utility of quantitative testing for *Candida albicans* (CPT 87482) has not been demonstrated. This test is considered Experimental, Investigational, or Unproven.

Chlamydia trachomatis

Test name	Procedure code	Medical Necessity
Chlamydia Trachomatis Detection, Direct Method	87490	Procedure code is medically necessary when criteria are met
Chlamydia Trachomatis Detection, Amplified Method	87491	Procedure code is medically necessary when criteria are met
Chlamydia Trachomatis Detection, Quantification Method	87492	Procedure code is experimental, investigational, or unproven

Direct or amplified Chlamydia trachomatis detection

Medical necessity requirements:

Nucleic acid amplification testing (NAAT) for *Chlamydia trachomatis* through either direct (CPT 87490) or amplified (CPT 87491) probe studies is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Routine annual screening of all sexually active women aged less than or equal to 25 years
- Screening of sexually active women greater than 25 years of age with risk factors (e.g., those who have a new sex partner or multiple sex partners)
- Routine screening for all pregnant women during one of the first prenatal visits
- Retesting of all pregnant women aged less than or equal to 25 years performed during the third trimester

- Retesting of all pregnant women over age 25 during the third trimester when at increased risk for Chlamydia (e.g., women who have a new or multiple sex partners, women with a history of a previous STI, high risk behavior such as inconsistent condom use, sex work)
- Screening of sexually active men with risk factors (e.g., men in correctional facilities, presenting to STI or adolescent clinics, or who have infected partner)
- Screening of all sexually active men who have sex with men (MSM), including those prescribed preexposure prophylaxis (PrEP) for prevention of HIV infection

Symptomatic individuals:

- Cervicitis
- Urethritis

Test frequency:

- Repeat testing to document eradication of infection after completing an appropriate treatment regimen is medically necessary only in the following settings: patient is pregnant, symptoms persist, re-infection is suspected, or compliance with therapy is in question. Routine test of cure is not medically necessary.
 - Non-pregnant recently infected women should be retested 3 to 12 months after treatment.
 - Based on guidelines for initial and repeat testing, no more than five screenings in a year should be necessary regardless of pregnancy or other risk factors.
- NAAT may be performed on urine, rectal, vaginal, or cervical samples. It is usually sufficient to test one site. When necessary to test more than one site, no more than 3 units of 87490 or 87491 for chlamydia trachomatis molecular testing is medically necessary for the same date of service.
- More than one type of molecular test for the same organism is not medically necessary for the same date of service.

Quantitative Chlamydia trachomatis testing

Medical necessity requirements:

Clinical utility of quantitative testing for chlamydia trachomatis (CPT 87492) has not been demonstrated. This test is considered Experimental, Investigational, or Unproven.

Gardnerella vaginalis

Test name	Procedure code	Medical Necessity
Gardnerella Vaginalis Detection, Direct Method	87510	Procedure code is medically necessary when criteria are met.
Gardnerella Vaginalis Detection, Amplified Method	87511	Procedure code is medically necessary when criteria are met.
Gardnerella Vaginalis Detection, Quantification	87512	Procedure code is experimental, investigational, or unproven

Direct or amplified Gardnerella vaginalis detection**Medical necessity requirements:**

Nucleic acid amplification testing (NAAT) for Gardnerella vaginalis through either direct (CPT 87510) or amplified (CPT 87511) probe studies is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Evidence does not support routine screening for Gardnerella vaginalis in asymptomatic pregnant women, asymptomatic non-pregnant women, or men for any indications.

Symptomatic individuals:

- Molecular testing for Gardnerella vaginalis in symptomatic females is medically necessary.

There are no medically necessary indications in males.

It should only be necessary to test one site. Therefore, only one unit per date of service is medically necessary.

More than one type of molecular test for the same organism is not medically necessary for the same date of service (e.g., 87510 and 87511 may not be billed together).

Quantitative Gardnerella vaginalis testing**Medical necessity requirements:**

Clinical utility of quantitative testing for Gardnerella vaginalis (CPT 87512) has not been demonstrated. This test is considered Experimental, Investigational, or Unproven.

Herpes simplex virus (HSV)

CPT 87528

Test name	Procedure code	Medical Necessity
Herpes Simplex Virus Detection, Direct Method	87528	Procedure code is medically necessary when criteria are met.
Herpes Simplex Virus Detection, Amplified Method	87529	Procedure code is medically necessary when criteria are met.
Herpes Simplex Virus Detection, Quantification Method	87530	Procedure code is medically necessary when criteria are met.

Direct or amplified Herpes simplex virus (HSV) detection

Medical necessity requirements:

Nucleic acid amplification testing (NAAT) for Herpes simplex virus (HSV) through either direct (CPT 87528) or amplified (CPT 87529) probe studies is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic Individuals:

- Current guidelines explicitly recommend against testing asymptomatic adults for HSV.

Symptomatic Individuals:

- New or recurrent vesicular and/or ulcerative lesions, vesicles or ulcers on or around the genitals, rectum, buttocks, thighs, back
- Recurrent genital symptoms or atypical symptoms and negative HSV cultures

It should only be necessary to test one site. Therefore, only one unit per date of service is medically necessary.

More than one type of molecular test for the same organism is not medically necessary for the same date of service (e.g., 87528 and 87529 may not be billed together).

Quantitative Herpes simplex virus testing

Quantitative testing for Herpes simplex virus (HSV) (CPT 87530) is medically necessary for monitoring disease in some circumstances.

Therefore, quantitative HSV testing is medically necessary when a diagnosis has been established and the need for monitoring is documented in the medical record. Quantitative HSV testing should not be used for the primary diagnosis of HSV.

Infectious agent, not otherwise specified

Test name	Procedure code	Medical Necessity
Miscellaneous Infectious Agent Detection, Direct Molecular Method	87797	Procedure code is experimental, investigational, or unproven for indications addressed by this guideline
Miscellaneous Infectious Agent Detection, Amplified Molecular Method	87798	Procedure code is experimental, investigational, or unproven for indications addressed by this guideline
Miscellaneous Infectious Agent Detection, Quantitative Molecular Method	87799	Procedure code is experimental, investigational, or unproven for indications addressed by this guideline

Miscellaneous infectious agent detection

Medical necessity requirements:

Molecular testing to detect a variety of organisms that do not have organism-specific procedure codes may be billed under the infectious agents not otherwise specified (NOS) codes (CPT 87797, 87798, 87799). This guideline only addresses some organisms and clinical settings. It does not apply to all testing performed under these codes.

The genitourinary organisms for which molecular testing is supported by guidelines are represented by organism-specific CPT codes. There are no clinical indications for any infectious agents billed under not otherwise (NOS) specified procedure codes that are supported by current evidence for the evaluation or management of genitourinary conditions, including bacterial vaginosis (addressed elsewhere in this guideline).

These tests are considered Experimental, Investigational, or Unproven.

Mycoplasma genitalium

Test name	Procedure code	Medical Necessity
Mycoplasma genitalium, Direct Method	87563	Procedure code is medically necessary when criteria are met

Direct Mycoplasma genitalium detection

Medical necessity requirements:

Nucleic acid amplification testing (NAAT) for Mycoplasma genitalium is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Screening for M. genitalium in sexual partners of people who are infected with M. genitalium is considered medically necessary.
- Routine screening for M. genitalium in asymptomatic individuals is otherwise not considered medically necessary, as the main benefit of testing is to determine appropriate course of treatment.

Symptomatic individuals:

- Cervicitis
- Urethritis
- Persistent PID when gonorrhea and chlamydia are negative

Test frequency:

- Repeat testing to document eradication of infection no earlier than 3 weeks after completing an appropriate treatment regimen is medically necessary in the following settings:
 - symptoms persist
 - re-infection is suspected
 - compliance with therapy is in question
- Routine test of cure in the absence of symptoms may be performed no earlier than 3 weeks after completing an appropriate treatment regimen.
- Based on guidelines for initial and repeat testing, no more than five screenings in a year should be necessary regardless of risk factors.
- NAAT for M. genitalium may be performed on urine, vaginal, cervical, male urethral, or penile meatal samples. It is usually sufficient to test one site. No

more than 3 units of 87563 for M. genitalium molecular testing are medically necessary for the same date of service.

Neisseria gonorrhoeae

Test name	Procedure code	Medical Necessity
Neisseria Gonorrhoeae, Direct Method	87590	Procedure code is medically necessary when criteria are met
Neisseria Gonorrhoeae, Amplified Method	87591	Procedure code is medically necessary when criteria are met
Neisseria Gonorrhoeae, Quantification Method	87592	Procedure code is experimental, investigational, or unproven

Direct or amplified Neisseria gonorrhoeae detection

Medical necessity requirements:

Nucleic acid amplification testing (NAAT) for Neisseria gonorrhoeae through either direct (CPT 87590) or amplified (CPT 87591) probe studies is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Annual screening of all sexually active women aged less than or equal to 25 years.
- Annual screening of women greater than 25 years who are at increased risk for infection (e.g., women with previous gonorrhea infection, other STIs, new or multiple sex partners, and inconsistent condom use, sex workers, or women living in communities with a high prevalence of disease).
- All pregnant women at increased risk for gonorrhea (as defined in the above criteria) should be screened at the first prenatal visit for N. gonorrhoeae.
- Uninfected pregnant women who remain at high risk for gonococcal infection also should be retested during the third trimester.
- Screening of sexually active individuals who have an infected partner.
- Screening of all sexually active adults prescribed preexposure prophylaxis (PrEP) for prevention of HIV infection.

Symptomatic individuals:

- Cervicitis
- Urethritis

Test frequency:

- When indicated, repeat testing to document eradication should not be performed until 3-4 weeks after the positive result. Pregnant women diagnosed with gonococcal infection during the first trimester should be retested within approximately 3–6 months, preferably in the third trimester. Recently infected individuals should be retested 3 to 12 months after treatment. When repeat testing is indicated, the following limitations apply:
 - Repeat testing is not medically necessary if performed within three weeks (less than 21 days) from a previous test.
 - Based on guidelines for initial and repeat testing, no more than five screenings in a year should be necessary regardless of pregnancy or other risk factors.
- Nucleic acid amplification test (NAAT) may be performed on urine, vaginal, or cervical samples. It is usually sufficient to test one site. When necessary to test more than one site, no more than 3 units of 87590 or 87591 for *Neisseria gonorrhoeae* molecular testing may be billed for the same date of service.
- More than one type of molecular test for the same organism is not medically necessary for the same date of service (e.g., 87590 and 87591 may not be billed together).

Quantitative *Neisseria gonorrhoeae* testing

Medical necessity requirements:

Clinical utility of quantitative testing for *Neisseria gonorrhoeae* (CPT 87592) has not been demonstrated. This test is considered Experimental, Investigational, or Unproven.

Trichomonas vaginalis

Test name	Procedure code	Medical Necessity
Trichomonas Vaginalis Detection, Direct Method	87660	Procedure code is medically necessary when criteria are met
Trichomonas Vaginalis Detection, Amplified Method	87661	Procedure code is medically necessary when criteria are met

Direct or amplified *Trichomonas vaginalis* detection

Medical necessity requirements:

Nucleic acid amplification testing (NAAT) for *Trichomonas vaginalis* through either direct (CPT 87660) or amplified (CPT 87661) probe studies is considered medically necessary for individuals with clinical indications as outlined here.

Asymptomatic individuals:

- Evidence does not support routine screening for *Trichomonas vaginalis* in asymptomatic women (pregnant or non-pregnant) or men who are not at high risk for infection.
- Screening can be considered in those at increased risk for *Trichomonas vaginalis* infection for reasons such as new or multiple sex partners, history of STIs, sex work, or drug use.
- Screening should also be performed in sexually active women who are HIV-positive at entry into care and then at least annually.

Symptomatic individuals:

- Vaginitis, abnormal vaginal discharge, cervicitis, nongonococcal urethritis, vulvar pruritis, or pelvic inflammatory disease.
- Sexually active women with trichomoniasis may be rescreened for *Trichomonas vaginalis* at 3 months following initial infection.
- Screening of sexually active individuals who have an infected partner.

Test frequency:

- Repeat testing should not be necessary more frequently than every three months.
- Nucleic acid amplification test (NAAT) may be performed on urine, vaginal, or cervical samples. It is usually sufficient to test one site. When necessary to test more than one site, no more than 3 units of 87660 or 87661 for *Trichomonas vaginalis* molecular testing are medically necessary for the same date of service.
- Based on guidelines for initial and repeat testing, no more than five screenings in a year should be necessary regardless of pregnancy or other risk factors.

Billing and Reimbursement

Introduction

This section outlines the billing requirements for tests addressed in this guideline. These requirements will be enforced during the case review process whenever appropriate. Examples of requirements may include specific coding scenarios, limits on allowable test combinations or frequency and/or information that must be provided on a claim for automated processing. Any claims submitted without the necessary information to allow for automated processing (e.g. ICD code, place of service, etc.) will not be reimbursable as billed. Any claim may require submission of medical records for post service review.

If the laboratory's testing platform consists solely of a panel of multiple targets, yet only a subset of the organisms are considered medically necessary based on the criteria included in this guideline, the laboratory can choose to bill using procedure codes that represent only the medically necessary organisms included on the panel (instead of a combined panel code).

Bacterial Vaginosis

When otherwise reimbursable, the following limitations apply:

- More than one type of molecular test for the same organism for the same date of service is not reimbursable (e.g., 81513 and 81514 may not be billed together).

Testing for bacterial vaginosis using any marker organisms or technologies not already addressed in this guideline is not eligible for reimbursement. This includes but is not limited to, the proprietary laboratory analysis (PLA) codes in the table above, as well as other BV panels, such as:

- NuSwab®, Vaginitis Plus (VG+) (A vaginae, BVAB-2, Megasphaera Type 1)
- OneSwab® (A vaginae, Megasphaera Type 1 and 2, BVAB-2)
- SureSwab® Vaginosis, Vaginitis Plus (G vaginalis, A vaginae, Megasphaera species)

Agent not otherwise specified (NOS) procedure codes will not be reimbursed for BV.

The following criteria are used to determine if testing for infectious agents NOS is being performed in the setting of genitourinary condition detection or management, including bacterial vaginosis:

- When billed with any ICD code included in Table: *ICD Codes Indicating NOS Testing Billed for Genitourinary Conditions*
- When billed on the same date of services with any other organism-specific CPT code referenced in this guideline.

Candida species

Direct or amplified Candida detection

When testing asymptomatic males or females, an ICD code that supports positive HIV status must be submitted on the claim (see Table: *ICD Codes Indicating HIV Positive Status*). Note that testing for males is only indicated when HIV positive (i.e., no symptomatic or other testing indications).

When testing symptomatic females, an ICD code that describes the common symptoms must be submitted on the claim (see Table: *ICD Codes Indicating Symptoms of Sexually Transmitted Infections and Other Infections of the Reproductive Tract*).

When testing is otherwise reimbursable, the following limitations apply:

- It should only be necessary to test one site. Therefore, only one unit per date of service is reimbursable.
- Subtyping for *Candida glabrata* and other non-*albicans* *Candida* species is not reimbursable; therefore, only one unit will be reimbursed. Exceptions may be considered on a case by case basis.
- More than one type of molecular test for the same organism will not be reimbursed for the same date of service (e.g., 87480 and 87481 may not be billed together).

Quantitative Candida testing

87482 is not reimbursable.

If the laboratory's testing platform consists of direct or amplified and quantitative testing methodologies, yet only direct or amplified testing is reimbursable based on these criteria, the lab may request reimbursement for only the reimbursable components of the test by using a procedure code that does not represent all testing methodologies performed.

Chlamydia trachomatis

Direct or amplified Chlamydia trachomatis detection

When testing is otherwise reimbursable, the following limitations apply:

- NAAT may be performed on urine, rectal, vaginal, or cervical samples. It is usually sufficient to test one site. When necessary to test more than one site, no more than 3 units of 87490 or 87491 for chlamydia trachomatis molecular testing may be billed for the same date of service.
- More than one type of molecular test for the same organism will not be reimbursed for the same date of service (e.g., 87490 and 87491 may not be billed together).

- No more than five screenings in a year are reimbursable.

Quantitative Chlamydia trachomatis testing

87492 is not reimbursable.

If the laboratory's testing platform consists of direct or amplified and quantitative testing methodologies, yet only direct or amplified testing is reimbursable based on these criteria, the lab may request reimbursement for only the reimbursable components of the test by using a procedure code that does not represent all testing methodologies performed.

Gardnerella vaginalis

Direct or amplified Gardnerella vaginalis detection

When testing symptomatic females, an ICD code that describes the common symptoms must be submitted on the claim (see Table: *ICD Codes Indicating Symptoms of Sexually Transmitted Infections and Other Infections of the Reproductive Tract*).

Testing in males is not medically necessary unless a KX modifier is included on the claim to indicate that the reported gender may not be consistent with the medical needs based on biological sex (e.g., transgender, transsexual, intersex individuals).

When testing is otherwise reimbursable, the following limitations apply:

- It should only be necessary to test one site. Therefore, only one unit per date of service is reimbursable.
- More than one type of molecular test for the same organism will not be reimbursed for the same date of service (e.g., 87510 and 87511 may not be billed together).

Quantitative Gardnerella vaginalis testing

87512 is not eligible for reimbursement.

If the laboratory's testing platform consists of direct or amplified and quantitative testing methodologies, yet only direct or amplified testing is considered reimbursable based on these criteria, the lab may request reimbursement for only the reimbursable components of the test by using a procedure code that does not represent all testing methodologies performed.

Herpes simplex virus (HSV)

Direct or amplified Herpes simplex virus (HSV) detection

When testing is otherwise reimbursable, the following limitations apply:

- It should only be necessary to test one site. Therefore, only one unit per date of service is reimbursable.
- More than one type of molecular test for the same organism will not be reimbursed for the same date of service (e.g., 87528 and 87529 may not be billed together).

Quantitative Herpes simplex virus testing

If the laboratory's testing platform consists of direct or amplified and quantitative testing methodologies, yet only direct or amplified testing is considered reimbursable based on these criteria, the lab may request reimbursement for only the reimbursable components of the test by using a procedure code that does not represent all testing methodologies performed.

Infectious agent, not otherwise specified

Miscellaneous infectious agent detection

The following criteria are used to determine if testing for infectious agents NOS is being performed in the setting of genitourinary condition detection or management, including bacterial vaginosis:

- When billed with any ICD code included in Table: *ICD Codes Indicating NOS Testing Billed for Genitourinary Conditions*.
- When billed on the same date of service with any other organism-specific CPT code referenced in this guideline.

Mycoplasma genitalium

Direct Mycoplasma genitalium detection

When testing is otherwise reimbursable, the following limitations apply:

- No more than 3 units of 87563 for M. genitalium molecular testing are reimbursable for the same date of service
- No more than five screenings in a year are reimbursable.

Neisseria gonorrhoeae

Direct or amplified Neisseria gonorrhoeae detection

When testing is otherwise reimbursable, the following limitations apply:

- When necessary to test more than one site, no more than 3 units of 87590 or 87591 for Neisseria gonorrhoeae molecular testing are reimbursable for the same date of service.

- More than one type of molecular test for the same organism will not be reimbursed for the same date of service (e.g., 87590 and 87591 may not be billed together).
- No more than five screenings in a year are reimbursable.
- Repeat testing will not be reimbursed if performed within three weeks (less than 21 days) from a previous test.

Quantitative Neisseria gonorrhoeae testing

87592 is not reimbursable.

If the laboratory's testing platform consists of direct or amplified and quantitative testing methodologies, yet only direct or amplified testing is reimbursable based on these criteria, the lab may request reimbursement for only the reimbursable components of the test by using a procedure code that does not represent all testing methodologies performed.

Trichomonas vaginalis

Direct or amplified Trichomonas vaginalis detection

When testing asymptomatic individuals, an ICD code that supports increased risk, infected partner, or positive HIV status must be submitted on the claim. See Tables: *ICD Codes Indicating High Risk Indications*, *ICD Codes Indicating Infected Partner*, *ICD Codes Indicating HIV Positive Status*.

When testing symptomatic individuals, an ICD code that describes the common symptoms, as defined in Table: *ICD Codes Indicating Symptoms of Sexually Transmitted Infections and Other Infections of the Reproductive Tract* must be submitted on the claim.

When testing is otherwise reimbursable, the following limitations apply:

- When necessary to test more than one site:
 - Additional units must be billed with modifier 59.
 - No more than 3 units of 87660 or 87661 for *Trichomonas vaginalis* molecular testing are reimbursable for the same date of service.
- More than one type of molecular test for the same organism will not be reimbursed for the same date of service (e.g., 87660 and 87661 may not be billed together).
- No more than five screenings in a year are reimbursable.
- Repeat testing is not reimbursable more frequently than every three months.

ICD Codes

ICD codes used for automated claims processing for this guideline.

Table: ICD Codes Indicating High Risk Indications

Codes and descriptions

Code or Range	Description
F10.X	Alcohol related disorders
F11.X	Opioid related disorders
F12.X	Cannabis related disorders
F13.X	Sedative, hypnotic, or anxiolytic related disorders
F14.X	Cocaine related disorders
F15.X	Other stimulant related disorders
F16.X	Hallucinogen related disorders
F18.X	Inhalant related disorders
F19.X	Other psychoactive substance related disorders
O99.32X	Drug use complicating pregnancy, childbirth, and the puerperium
Z72.5X	High risk sexual behavior
Z77.9	Other contact with and (suspected) exposures hazardous to health

Table: ICD Codes Indicating Infected Partner

Codes and descriptions

Code or Range	Description
Z20.2	Contact with and (suspected) exposure to infections with a predominantly sexual mode of transmission
Z20.6	Contact with and (suspected) exposure to human immunodeficiency virus [HIV]
Z20.8X	Contact with and (suspected) exposure to other communicable diseases
Z20.9	Contact with and (suspected) exposure to unspecified communicable disease

Table: ICD Codes Indicating HIV Positive Status

Codes and descriptions

Code or Range	Description
B20	Human immunodeficiency virus [HIV] disease
B97.35	Human immunodeficiency virus, type 2 [HIV-2]
O98.7X	Human immunodeficiency virus [HIV] disease complicating pregnancy, childbirth and the puerperium
R75	Inconclusive laboratory evidence of human immunodeficiency virus [HIV]
Z21	Asymptomatic human immunodeficiency virus [HIV] infection status

Table: ICD Codes Indicating Symptoms of Sexually Transmitted Infections and Other Infections of the Reproductive Tract

The ICD codes in the following table suggest medical necessity for the procedure codes for trichomonas vaginalis (87661), candida (87480, 87481), and gardnerella vaginalis (87510, 87511).

Code or Range	Description
A56.X	Other sexually transmitted chlamydial diseases
A59.X	Urogenital trichomoniasis
B37.3X	Candidiasis of vulva and vagina
B37.4X	Candidiasis of other urogenital sites
L29.X	Pruritus
N34.X	Urethritis and urethral syndrome
N35.1X	Postinfective urethral stricture, not elsewhere classified
N37	Urethral disorders in diseases classified elsewhere
N72	Inflammatory disease of cervix uteri
N73.X	Other female pelvic inflammatory diseases
N75.X	Diseases of Bartholin's gland

Code or Range	Description
N76.X	Other inflammation of vagina and vulva
N77.X	Vulvovaginal ulceration and inflammation in diseases classified elsewhere
N89.8	Other specified noninflammatory disorders of vagina
N89.9	Noninflammatory disorder of vagina, unspecified
N94.1X	Dyspareunia
N95.2	Postmenopausal atrophic vaginitis
O23.X	Infections of genitourinary tract in pregnancy
O86.X	Other puerperal infections
R10.2	Pelvic and perineal pain

Table: ICD Codes Indicating NOS Testing Billed for Genitourinary Conditions

The ICD codes in the following table indicate when procedure codes for infectious agent detection by nucleic acid not otherwise specified (NOS) (87797-87799) are billed for GU organisms.

Code or Range	Description
A50.X	Congenital syphilis
A51.X	Early syphilis
A52.X	Late syphilis
A53.X	Other and unspecified syphilis
A54.X	Gonococcal infection
A55	Chlamydial lymphogranuloma (venereum)
A56.X	Other sexually transmitted chlamydial diseases
A57	Chancroid
A58	Granuloma inguinale
A59.X	Trichomoniasis
A60.X	Anogenital herpesviral [herpes simplex] infections
A63.X	Other predominantly sexually transmitted diseases, not elsewhere classified

Code or Range	Description
A64	Unspecified sexually transmitted disease
A74.89	Other chlamydial diseases
A74.9	Chlamydial infection, unspecified (includes childbirth and postpartum)
B37.3X	Candidiasis of vulva and vagina
B37.4X	Candidiasis of other urogenital sites
B97.7	Papillomavirus as the cause of diseases classified elsewhere
L29.X	Pruritus
M02.30	Reiter's disease, unspecified site
N34.X	Urethritis and urethral syndrome
N35.X	Urethral stricture
N37	Urethral disorders in diseases classified elsewhere
N39.0	Urinary tract infection, site not specified
N39.9	Disorder of urinary system, unspecified
N70.X	Salpingitis and oophoritis
N71.X	Inflammatory disease of uterus, except cervix
N72	Inflammatory disease of cervix uteri
N73.X	Other female pelvic inflammatory diseases
N74	Female pelvic inflammatory disorders in diseases classified elsewhere
N75.X	Diseases of Bartholin's gland
N76.X	Other inflammation of vagina and vulva
N77.X	Vulvovaginal ulceration and inflammation in diseases classified elsewhere
N87.X	Dysplasia of cervix uteri
N94.1X	Dyspareunia
O09.X	Supervision of high risk pregnancy
O23.X	Infections of genitourinary tract in pregnancy
O86.X	Other puerperal infections

Code or Range	Description
R87.5	Abnormal microbiological findings in specimens from female genital organs
R87.6X	Abnormal cytological findings in specimens from female genital organs
R87.8X	Other abnormal findings in specimens from female genital organs
Z00.00	Encounter for general adult medical examination without abnormal findings
Z00.8	Encounter for other general examination
Z01.4X	Encounter for gynecological examination
Z11.3	Encounter for screening for infections with a predominantly sexual mode of transmission
Z11.51	Encounter for screening for human papillomavirus (HPV)
Z11.59	Encounter for screening for other viral diseases
Z11.8	Encounter for screening for other infectious and parasitic diseases
Z11.9	Encounter for screening for infectious and parasitic diseases, unspecified
Z12.4	Encounter for screening for malignant neoplasm of cervix
Z20.2	Contact with and (suspected) exposure to infections with a predominantly sexual mode of transmission
Z20.6	Contact with and (suspected) exposure to human immunodeficiency virus [HIV]
Z20.8X	Contact with and (suspected) exposure to communicable diseases
Z20.9	Contact with and (suspected) exposure to unspecified communicable disease
Z30.X	Encounter for contraceptive management
Z31.X	Encounter for procreative management
Z32.X	Encounter for pregnancy test and childbirth and childcare instruction
Z33.X	Pregnant state

Code or Range	Description
Z34.X	Encounter for supervision of normal pregnancy
Z36	Encounter for antenatal screening of mother
Z39.X	Encounter for maternal postpartum care and examination
Z64.0	Problems related to unwanted pregnancy
Z64.1	Problems related to multiparity
Z71.7	Human immunodeficiency virus [HIV] counseling
Z72.5X	High risk sexual behavior
Z77.9	Other contact with and (suspected) exposures hazardous to health
Z97.5	Presence of (intrauterine) contraceptive device

Test information

Introduction

Molecular testing for sexually transmitted infections (STIs) and other infections of the reproductive tract may include nucleic acid testing, nucleic acid amplification testing (NAAT), or other specialized molecular studies.

For a more detailed description of these techniques, please refer to the guideline, *Infectious Disease Laboratory Testing*.

Guidelines and evidence

Introduction

This section includes relevant guidelines and evidence pertaining to molecular testing for sexually transmitted infections (STIs) and other infections of the reproductive tract.

Bacterial vaginosis

American College of Obstetricians and Gynecologists

The American College of Obstetricians and Gynecologists (ACOG, 2020) Practice Bulletin Vaginitis in Nonpregnant Patients¹ stated that "[b]ecause the normal vaginal

flora is heterogeneous, routine bacterial culture of the vagina is not specific for bacterial vaginosis. For this reason, bacterial culture is not recommended for the diagnosis."

ACOG further stated that a clinical diagnosis for bacterial vaginosis is based on the Amsel criteria and requires the presence of three out of four clinical criteria (abnormal discharge, pH >4.5, positive KOH whiff test result, and presence of more than 20% clue cells on microscopy). Gram stain with Nugent scoring is considered the reference standard for a bacterial vaginosis diagnosis, however, it is generally limited to research settings.

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines stated that:²

- "BV can be diagnosed by using clinical criteria (i.e., Amsel's diagnostic criteria or by determining the Nugent score from a vaginal Gram stain. Vaginal Gram stain, considered the reference standard laboratory method for diagnosing BV, is used to determine the relative concentration of lactobacilli (i.e., long gram-positive rods), small gram-negative and gram-variable rods (i.e., *G. vaginalis* or *Bacterioides*), and curved gram-negative rods (i.e., *Mobiluncus*) characteristic of BV."
- "Multiple BV NAATs are available for BV diagnosis among symptomatic women. These tests are based on detection of specific bacterial nucleic acids and have high sensitivity and specificity for BV (i.e., *G. vaginalis*, *A. vaginae*, BVAB2, or *Megasphaera* type 1) and certain lactobacilli (i.e., *Lactobacillus crispatus*, *Lactobacillus jensenii*, and *Lactobacillus gasseri*)."
- "BV NAATs should be used among symptomatic women only (e.g., women with vaginal discharge, odor, or itch) because their accuracy is not well defined for asymptomatic women. Despite the availability of BV NAATs, traditional methods of BV diagnosis, including the Amsel criteria, Nugent score, and the Affirm VP III assay, remain useful for diagnosing symptomatic BV because of their lower cost and ability to provide a rapid diagnosis."

Society of Obstetricians and Gynecologists of Canada

The Society of Obstetricians and Gynecologists of Canada (SOGC) has published guidelines for the screening and management of bacterial vaginosis (2015)³ and additional guidelines for screening and management during pregnancy (2017) that state the following:⁴

- "Bacterial vaginosis should be diagnosed using either clinical (Amsel's) or laboratory (Gram stain with objective scoring system) criteria. (II-2A)"

Candida species

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines recommended the following in regard to candida species testing:²

- "Examination of a wet mount with KOH preparation should be performed for all women with symptoms or signs of VVC, and women with a positive result should be treated. For those with negative wet mounts but existing signs or symptoms, vaginal cultures for Candida should be considered. If Candida cultures cannot be performed for these women, empiric treatment can be considered. Identifying Candida by culture in the absence of symptoms or signs is not an indication for treatment because approximately 10%–20% of women harbor Candida species and other yeasts in the vagina. The majority of PCR tests for yeast are not FDA cleared, and providers who use these tests should be familiar with the performance characteristics of the specific test used. Yeast culture, which can identify a broad group of pathogenic yeasts, remains the reference standard for diagnosis."

Chlamydia trachomatis

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines recommended the following in regard to chlamydia trachomatis testing:²

- "Annual screening of all sexually active women aged <25 years is recommended, as is screening of older women at increased risk for infection (e.g., women aged ≥25 who have a new sex partner, more than one sex partner, a sex partner with concurrent partners, or a sex partner who has an STI... Although chlamydia incidence might be higher in some women aged ≥25 years in certain communities, overall, the largest proportion of infection is among women aged <25 years."
- "Although evidence is insufficient to recommend routine screening for C. trachomatis among sexually active young men because of certain factors (i.e., feasibility, efficacy, and cost-effectiveness), screening of sexually active young men should be considered in clinical settings with a high prevalence of chlamydia (e.g., adolescent clinics, correctional facilities, or STD specialty clinics) or for populations with a high burden of infection (e.g., MSM)."
- "NAATs are . . . the recommended test for detecting C. trachomatis infection."
- "Among symptomatic patients, POC tests for C. trachomatis can optimize treatment by limiting unnecessary presumptive treatment at the time of clinical decision-making and improve antimicrobial stewardship. Thus, using a POC test

will likely be a cost-effective diagnostic strategy for *C. trachomatis* infection. Newer NAAT-based POC tests have promising performance and are becoming commercially available.”

The Centers for Disease Control and Prevention (CDC, 2021) Preexposure Prophylaxis for the Prevention of HIV Infection in the United States Clinical Practice Guideline recommended the following:⁵

- "Tests to screen for chlamydia are recommended for all sexually active MSM prescribed PrEP, both at screening prior to initiation and at quarterly visits."
- "Chlamydia is very common, especially in young women and does not correlate strongly with risk of HIV acquisition so does not serve as an indication for initiating PrEP. However, because it is a frequent infection among sexually active women at high risk, screening for chlamydia is recommended at initiation and every 12 months for all sexually active women as a component of PrEP care."

U.S. Preventive Services Task Force

The U.S. Preventive Services Task Force (USPSTF, 2021) further recommended:⁶

- “The USPSTF recommends screening for chlamydia in sexually active females aged 24 years or younger and in older women who are at increased risk for infection. (B recommendation)”
- “The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for chlamydia and gonorrhea in men. (I statement)”

Gardnerella Vaginalis

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Diseases Treatment Guidelines stated the following in regard to *Gardnerella vaginalis* testing:²

- "BV can be diagnosed by using clinical criteria (i.e., Amsel's diagnostic criteria) or by determining the Nugent score from a vaginal Gram stain. Vaginal Gram stain, considered the reference standard laboratory method for diagnosing BV, is used to determine the relative concentration of lactobacilli (i.e., long gram-positive rods), small gram-negative and gram-variable rods (i.e., *G. vaginalis* or *Bacteroides*), and curved gram-negative rods (i.e., *Mobiluncus*) characteristic of BV. A Nugent score of 0–3 is consistent with a *Lactobacillus*-predominant vaginal microbiota, 4–6 with intermediate microbiota (emergence of *G. vaginalis*), and 7–10 with BV."

- "In addition to the Amsel criteria, multiple POC tests are available for BV diagnosis. The Osom BV Blue test (Sekisui Diagnostics) detects vaginal sialidase activity. The Affirm VP III (Becton Dickinson) is an oligonucleotide probe test that detects high concentrations of *G. vaginalis* nucleic acids (>5 x 10⁵ CFU of *G. vaginalis*/mL of vaginal fluid) for diagnosing BV, *Candida* species, and *T. vaginalis*. This test has been reported to be most useful for symptomatic women in conjunction with vaginal pH measurement and presence of amine odor (sensitivity of 97%); specificity is 81% compared with Nugent."
- "Despite the availability of BV NAATs, traditional methods of BV diagnosis, including the Amsel criteria, Nugent score, and the Affirm VP III assay, remain useful for diagnosing symptomatic BV because of their lower cost and ability to provide a rapid diagnosis. Culture of *G. vaginalis* is not recommended as a diagnostic tool because it is not specific."

Herpes simplex virus

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines recommended the following in regard to herpes simplex virus testing:²

- "[A]ll persons who have genital, anal, or perianal ulcers should be evaluated. Specific evaluation of genital, anal, or perianal ulcers includes syphilis serology tests and darkfield examination from lesion exudate or tissue, or NAAT if available; NAAT or culture for genital herpes type 1 or 2; and serologic testing for type-specific HSV antibody."
- "HSV PCR of the blood should not be performed to diagnose genital herpes infection, except in cases in which concern exists for disseminated infection (e.g., hepatitis)."

U.S. Preventive Services Task Force

The U.S. Preventive Services Task Force (USPSTF, 2023) further recommended:⁷

- "The USPSTF recommends against routine serologic screening for genital herpes simplex virus infection in asymptomatic adolescents and adults, including pregnant persons. (D statement)"

Mycoplasma genitalium

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines stated that the main use of testing for *M. genitalium* is in patients with persistent or recurrent urethritis, cervicitis or pelvic inflammatory

disease (PID).² The CDC noted that the main method for testing is NAAT.² The guideline stated:

- "M. genitalium is an extremely slow-growing organism. Culture can take up to 6 months, and technical laboratory capacity is limited to research settings. NAAT for M. genitalium is FDA cleared for use with urine and urethral, penile meatal, endocervical, and vaginal swab samples."
- "M. genitalium should be suspected in cases of persistent or recurrent urethritis and or cervicitis and considered for PID."

Other Guidelines

The British Association for Sexual Health and HIV (2018) published a national guideline for the management of infection with M. genitalium.⁸ The guideline does not recommend screening asymptomatic individuals, but it does recommend testing for sexual partners of infected individuals. It stated:⁸

- "The evidence suggests that the majority of people infected with M. genitalium in the genital tract do not develop disease....Current treatments are imperfect and associated with development of antimicrobial resistance... There is no evidence that screening asymptomatic individuals will be of benefit, and indeed is likely to do harm at a population level."
- "We recommend testing current sexual partners of persons infected with *M. genitalium*."

Neisseria gonorrhoeae

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines recommended the following in regard to neisseria gonorrhoeae testing:²

- "Annual screening for N. gonorrhoeae infection is recommended for all sexually active women aged <25 years and for older women at increased risk for infection (e.g., those aged ≥25 years who have a new sex partner, more than one sex partner, a sex partner with concurrent partners, or a sex partner who has an STI)."

The Centers for Disease Control and Prevention (CDC, 2021) Preexposure Prophylaxis for the Prevention of HIV Infection in the United States Clinical Practice Guideline recommended the following:⁵

- "Tests to screen for gonorrhea are recommended for all sexually active adults prescribed PrEP, both at screening, for MSM at quarterly visits, and for women at semi-annual visits."

U.S. Preventive Services Task Force

The U.S. Preventive Services Task Force (USPSTF, 2021) further recommended:⁶

- “The USPSTF recommends screening for gonorrhea in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection. (B recommendation)”
- “The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for chlamydia and gonorrhea in men. (I statement)”

Trichomonas vaginalis

Centers for Disease Control and Prevention

The Centers for Disease Control and Prevention (CDC, 2021) Sexually Transmitted Infections Treatment Guidelines recommended the following in regard to *Trichomonas vaginalis* testing:²

- “Symptomatic pregnant women, regardless of pregnancy stage, should be tested and treated. . . . The benefit of routine screening for *T. vaginalis* in asymptomatic pregnant women has not been established.”
- "Because of the high prevalence of *T. vaginalis* among women with HIV and the potential for adverse reproductive health, poor birth outcomes, and possibly amplified HIV transmission, routine screening and prompt treatment are recommended for all women with HIV infection; screening should occur at entry to care and then at least annually thereafter."

Multiple Organism Detection Panels

Infectious Diseases Society of America

An IDSA (2018) Clinical Practice Guideline for Laboratory Diagnosis of Infectious Diseases, Section XI. Genital Infections recommended the following:⁹

- "Testing simultaneously for CT [*Chlamydia trachomatis*], GC [*Neisseria gonorrhoeae*], and *Trichomonas* is optimal for detection of the most common treatable STIs in female patients."

References

Introduction

These references are cited in this guideline.

1. ACOG Committee on Practice Bulletins-Gynecology. ACOG Practice Bulletin, Clinical management guidelines for obstetrician-gynecologists, Number 215: Vaginitis in nonpregnant patients. *Obstet Gynecol*. 2020;135(1):e1-e17.
2. Workowski KA, Bachmann LH, Chan PA, et al.; Centers for Disease Control and Prevention (CDC). Sexually transmitted infections treatment guidelines, 2021. *MMWR Recomm Rep*. 2021 70(4):1-187. doi: 10.15585/mmwr.rr7004a1
3. van Schalkwyk J, Yudin MH, Infectious Disease Committee. Vulvovaginitis – Screening for and management of trichomoniasis, vulvovaginal candidiasis, and bacterial vaginosis. *J Obstet Gynaecol Can*. 2015;37(3):266–274.
4. Yudin MH, Money DM. No. 211 – Screening and management of bacterial vaginosis in pregnancy. *J Obstet Gynaecol Can*. 2017;39(8):e184-e191.
5. Centers for Disease Control and Prevention (CDC). Preexposure Prophylaxis for the Prevention of HIV Infection in the United States- 2021 Update: A Clinical Practice Guideline. Available at: <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf> <https://www.cdc.gov/hiv/pdf/risk/prep/cdc-hiv-prep-guidelines-2021.pdf>
6. U.S. Preventive Services Task Force. Chlamydia and Gonorrhea Screening: U.S. Preventive Services Task Force Recommendation Statement. Available at: <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/chlamydia-and-gonorrhea-screening#fullrecommendationstart>
7. U.S. Preventive Services Task Force. Genital Herpes Infection: Serologic Screening: U.S. Preventive Services Task Force Recommendation Statement. Available at: <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/genital-herpes-serologic-screening>
8. Soni S, Homer P, Rayment M. et al. British Association for Sexual Health and HIV national guideline for the management of infection with *Mycoplasma genitalium* (2018). *Int J STD AIDS*. 2019;30(10):938-950.
9. Miller JM, Binnicker MJ, Campbell S, et al. A Guide to Utilization of the Microbiology Laboratory for Diagnosis of Infectious Diseases: 2018 Update by the Infectious Disease Society of America and the American Society for Microbiology. *Clin Infect Dis*. 2018;67(6):e1-e94. doi: 10.1093/cid/ciy381
10. NuSwab New Approach to Testing for Infections of the Genital Tract. Available at: <https://www.labcorp.com/tests/related-documents/L9603>
11. LabCorp. Vaginitis Plus (VG+), NuSwab®. Updated 2023. Available at: <https://www.labcorp.com/tests/180021/vaginitis-plus-vg-nuswab>
12. Medical Diagnostic Laboratories L.L.C. Testing Menu. Updated 2023. Available at: <http://www.mdlab.com/resources/testing-menu>

13. Medical Diagnostic Laboratories L.L.C. Bacterial vaginosis (with Lactobacillus profiling) qPCR panel. Updated February 22, 2018. Available at:
http://www.mdlab.com/pdf/test_bulletins/BV_Lacto_Tech_Bulletin.pdf
14. Quest Diagnostics Test Center. Available at:
<http://www.questdiagnostics.com/testcenter/BUOrderInfo.action?tc=17333&labCode=DAL>