



HOME HEALTHCARE

Supplementary Guidelines

Effective: September 03, 2024

VERSION 1.0.2024

EviCore
By **EVERNORTH**

EviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Imaging requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

EviCore's Clinical Review Criteria ("CRC") and related content is made available for the limited uses of: reference; and individual use, only limited to facilitating the determination of medically necessary and appropriate clinical treatment by clinicians for specific delegated patients under their care. The CRC and related content is proprietary information of eviCore, and copyrighted to the full extent of the law. Except as expressly permitted, you may not modify, copy, reproduce, republish, upload, post, transmit, hyperlink to or from, or distribute in any way the CRC, nor may you sell, transfer, distribute, assign, lease, reproduce, or otherwise use the CRC in commerce, in a manner that competes with us or infringes upon our rights, or for any public or commercial endeavor without our prior and express written consent.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2024 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Administrative Guidelines

Guideline	Page
Definitions and abbreviations.....	3

Definitions and abbreviations

HHC.AD.101
v1.0.2024

Non-skilled Maintenance Care	Services not requiring assistance of a licensed nurse or therapist
Skilled Maintenance Care	Services required to maintain the individual's current condition or to prevent or slow deterioration of the individual's condition
Palliative Care	Care typically performed to manage or alleviate symptoms and does not provide corrective benefit to the condition

Clinical Use Guidelines

Guideline	Page
Home Health Aide Visits.....	5
Social Worker Visits.....	8

Supplementary Guidelines

Home Health Aide Visits

HHC.CU.100.A**v1.0.2024**

- When non-skilled (not requiring assistance of licensed nurse or therapist) assistance is needed for personal care or other health-related services to maintain the individual's health or facilitate treatment, a Home Health Aide (HHA) referral is indicated if **all** of the following apply:
 - Non-skilled services will support a skilled treatment plan
 - There is documentation of significant functional limitations and **both**:
 - The individual is unable to perform the service independently
 - There is no caregiver available to provide supportive care or the individual will not allow the available caregiver to perform the needed care
 - There is an expectation that the individual's condition will improve significantly in a reasonable period of time

Initial services

Once criteria for home health aide referral has been satisfied, visit timing and frequency for initial services is guided based on the level of functional impairment.

- For **initial services** the following visits can be approved over a 2-week period:
 - 6 visits (12 hours) for an individual with minimal functional impairment documented by one or more of the following conditions:
 - Individual requires assistance with ADLs or IADLs
 - Individual ambulates with assistance of a person or device
 - Individual requires cognitive or communication assistance from a person or device
 - Incontinence of bowel or bladder one or more times per week
 - Individual requires some assistance with transfers
 - 10 visits (20 hours) for an individual with moderate functional impairment documented by two or more of the following conditions:
 - Individual requires assistance with ADLs or IADLs
 - Individual ambulates with assistance of a person or device
 - Individual requires cognitive or communication assistance from a person or device
 - Incontinence of bowel or bladder one or more times per week
 - Individual requires some assistance with transfers
 - 14 visits (28 hours) for an individual with maximal functional impairment documented by three or more of the following conditions:

- Individual requires total assistance with ADLs or IADLs
- Bowel or bladder incontinence
- Individual is non-ambulatory
- Individual is non-communicating or unable to follow safety measures
- Individual requires 1 or 2 people to assist with transfers
- 28 visits (56 hours) for an individual without available caregiver support when there is maximal functional impairment documented by three or more of the following conditions:
 - Individual requires total assistance with ADLs or IADLs
 - Bowel or bladder incontinence
 - Individual is non-ambulatory
 - Individual is non-communicating or unable to follow safety measures
 - Individual requires 1 or 2 people to assist with transfers

Continued services

- Once criteria for home health aide referral has been satisfied, visit timing and frequency for continuing services is guided based on the level of functional impairment.
 - For **continued services** the following visits can be approved over a two-week period:
 - 5 visits (10 hours) for an individual with minimal functional impairment documented by one or more of the following conditions:
 - Individual requires assistance with ADLs or IADLs
 - Individual ambulates with assistance of a person or device
 - Individual requires cognitive or communication assistance from a person or device
 - Incontinence of bowel or bladder one or more times per week
 - Individual requires some assistance with transfers
 - 10 visits (20 hours) for an individual with moderate functional impairment documented by two or more of the following conditions:
 - Individual requires assistance with ADLs or IADLs
 - Individual ambulates with assistance of a person or device
 - Individual requires cognitive or communication assistance from a person or device
 - Incontinence of bowel or bladder one or more times per week
 - Individual requires some assistance with transfers
 - 14 visits (28 hours) for an individual with maximal functional impairment documented by three or more of the following conditions:

- Individual requires total assistance with ADLs or IADLs
- Bowel or bladder incontinence
- Individual is non-ambulatory
- Individual is non-communicating or unable to follow safety measures
- Individual requires 1 or 2 people to assist with transfers
- 28 visits (56 hours) for an individual without available caregiver support when there is maximal functional impairment documented by three or more of the following conditions:
 - Individual requires total assistance with ADLs or IADLs
 - Bowel or bladder incontinence
 - Individual is non-ambulatory
 - Individual is non-communicating or unable to follow safety measures
 - Individual requires 1 or 2 people to assist with transfers

References

1. American Academy of Pediatrics Council on Children with Disabilities. Care coordination in the medical home: integrating health and related systems of care for children with special health care needs. *Pediatrics*. 2005;116(5):1238-1244. doi:10.1542/peds.2005-2070.
2. Brayman SJ, Roley SS, Clark GF, et. al. Standards of Practice for Occupational Therapy. *American Journal of Occupational Therapy*. 2005;59(6):663-665. doi:10.5014/ajot.59.6.663.
3. Bureau of Labor Statistics, U.S. Department of Labor, Occupational Outlook Handbook, Social Workers, at <https://www.bls.gov/ooh/community-and-social-service/social-workers.htm>.
4. Centers for Medicare & Medicaid Services. Chapter 7 - Home Health Services. In: *Medicare Benefit Policy Manual*. Bethesda, MD: Centers for Medicare & Medicaid Services; 2017.
5. Centers for Medicare & Medicaid Services (CMS). Home health agency manual, section 204.1A. Centers for Medicare & Medicaid Services, U.S. Department of Health and Human Services; 2014. Available from: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c07.pdf>.
6. Grilli L, Feldman DE, Majnemer A, et al. Associations between a functional independence measure (WeeFIM) and the pediatric quality of life inventory (PedsQL4.0) in young children with physical disabilities. *Quality of Life Research*. 2006;15(6):1023-1031. doi:10.1007/s11136-006-0041-9.
7. Institute of Medicine (US) Committee on Palliative and End-of-Life Care for Children and Their Families, Field MJ, Behrman RE, eds. When Children Die: Improving Palliative and End-of-Life Care for Children and Their Families. Washington (DC): *National Academies Press* (US); 2003.
8. Msall ME, DiGaudio K, Duffy LC, et. al. WeeFIM. Normative sample of an instrument for tracking functional independence in children. *Clin Pediatr* (Phila). 1994 Jul;33(7):431-8. doi: 10.1177/000992289403300709.
9. Msall ME, Digaudio K, Rogers BT, et al. The Functional Independence Measure for Children (WeeFIM). *Clinical Pediatrics*. 1994;33(7):421-430. doi:10.1177/000992289403300708.
10. National Guideline Centre (UK). Evidence review: Barriers to accessing end of life care services: End of life care for adults: service delivery. London: *National Institute for Health and Care Excellence (UK)*; October 2019.
11. Ottenbacher et al. The reliability of the functional independence measure: a quantitative review. *Arch Phys Med Rehabil* 1996. 77(12):1226-32.
12. What is Hospice? *Hospice Foundation of America*. <https://hospicefoundation.org/Hospice-Care/Hospice-Services>.

Social Worker Visits

HHC.CU.101.A

v1.0.2024

For the purpose of this guideline, social service visits are considered to be medically necessary when there are social or emotional barriers to the effective medical treatment or rate of recovery of an individual and there is a plan of care defining intervention that requires services of a social worker in order to be performed safely and effectively.

Social Worker Visits

- An individual qualifies for two visits with a licensed social worker during a two-week period when both of the following apply
 - To qualify for home health care, the patient must be:
 - Confined to the home
 - Under the care of a physician
 - Receiving services with an established plan of care that is periodically reviewed by a physician
 - In need of skilled care on an intermittent basis by a nurse, physical therapist, or speech-language pathologist
 - In need of continued treatment by occupational therapy
 - There is a care plan establishing the required services for:
 - Evaluation of barriers to the individual's medical treatment or recovery due to:
 - home environment
 - financial resources
 - community resource needs
 - Assistance with obtaining community services
 - Assistance with long-term planning, durable power of attorney, or legal guardian
- Two visits with a licensed social worker during a two-week period can be approved for patient, family, or caregiver evaluation and intervention when
 - Previously established caregiver is no longer available
 - There is suspected or identified abuse or neglect
 - There is increased interpersonal conflict or other evidence of ineffective coping

References

1. American Academy of Pediatrics Council on Children with Disabilities. Care coordination in the medical home: integrating health and related systems of care for children with special health care needs. *Pediatrics*. 2005;116(5):1238-1244. doi:10.1542/peds.2005-2070.

2. Brayman SJ, Roley SS, Clark GF, et. al. Standards of Practice for Occupational Therapy. *American Journal of Occupational Therapy*. 2005;59(6):663-665. doi:10.5014/ajot.59.6.663.
3. Bureau of Labor Statistics, U.S. Department of Labor, Occupational Outlook Handbook, Social Workers, at <https://www.bls.gov/ooh/community-and-social-service/social-workers.htm>.
4. Centers for Medicare & Medicaid Services. Chapter 7 - Home Health Services. In: *Medicare Benefit Policy Manual*. Bethesda, MD: Centers for Medicare & Medicaid Services; 2017.
5. Centers for Medicare & Medicaid Services (CMS). Home health agency manual, section 204.1A. Centers for Medicare & Medicaid Services, U.S. Department of Health and Human Services; 2014. Available from: <https://www.cms.gov/Regulations-and-Guidance/Guidance/Manuals/Downloads/bp102c07.pdf>.
6. Grilli L, Feldman DE, Majnemer A, et al. Associations between a functional independence measure (WeeFIM) and the pediatric quality of life inventory (PedsQL4.0) in young children with physical disabilities. *Quality of Life Research*. 2006;15(6):1023-1031. doi:10.1007/s11136-006-0041-9.
7. Institute of Medicine (US) Committee on Palliative and End-of-Life Care for Children and Their Families, Field MJ, Behrman RE, eds. When Children Die: Improving Palliative and End-of-Life Care for Children and Their Families. Washington (DC): *National Academies Press* (US); 2003.
8. Msall ME, DiGaudio K, Duffy LC, et. al. WeeFIM. Normative sample of an instrument for tracking functional independence in children. *Clin Pediatr* (Phila). 1994 Jul;33(7):431-8. doi: 10.1177/000992289403300709.
9. Msall ME, Digaudio K, Rogers BT, et al. The Functional Independence Measure for Children (WeeFIM). *Clinical Pediatrics*. 1994;33(7):421-430. doi:10.1177/000992289403300708.
10. National Guideline Centre (UK). Evidence review: Barriers to accessing end of life care services: End of life care for adults: service delivery. London: *National Institute for Health and Care Excellence (UK)*; October 2019.
11. Ottenbacher et al. The reliability of the functional independence measure: a quantitative review. *Arch Phys Med Rehabil* 1996. 77(12):1226-32.
12. What is Hospice? *Hospice Foundation of America*. <https://hospicefoundation.org/Hospice-Care/Hospice-Services>.