

Liquid Biopsy Testing

MOL.TS.194.A
v1.0.2025

Introduction

Liquid biopsy testing is addressed by this guideline.

Procedures addressed

The inclusion of any procedure code in this table does not imply that the code is under management or requires prior authorization. Refer to the specific Health Plan's procedure code list for management requirements.

Procedures addressed by this guideline	Procedure codes
ABL1 Mutation Analysis	81170
APC Sequencing	81201
ASXL1 Full Gene Sequencing	81175
ASXL1 Mutation Analysis	81176
BRAF V600 Targeted Mutation Analysis	81210
BRCA1/2 Sequencing	81163
BRCA1 Sequencing	81165
BRCA2 Sequencing	81216
CALR Exon 9 Mutation Analysis	81219
Caris Assure	0485U
CCND1/IGH (t(11;14)) Translocation Analysis, Major Breakpoint	81168
CEBPA Full Gene Sequencing	81218
EGFR Targeted Mutation Analysis	81235
Epic Sciences ctDNA Metastatic Breast Cancer Panel	0428U
EZH2 Common Variant(s) (e.g. codon 646)	81237
EZH2 Full Gene Sequencing	81236
FLT3 Mutation Analysis (internal tandem duplication variants)	81245
FLT3 Mutation Analysis (tyrosine kinase domain variants)	81246

Procedures addressed by this guideline	Procedure codes
FoundationOne Liquid CDx	0239U
Guardant360 CDx	0242U
Guardant360 LDT	0326U
Guardant360 Response	0422U
Hematolymphoid Neoplasm Molecular Profiling; 5-50 Genes	81450
IDH1 Mutation Analysis	81120
IDH2 Mutation Analysis	81121
IGH@/BCL2 (t(14;18)) Translocation Analysis, Major Breakpoint Region (MBR) and Minor Cluster Region (mcr) Breakpoints	81278
InVisionFirst-Lung Liquid Biopsy, Inivata, Inc.	0388U
JAK2 Targeted Mutation Analysis (e.g exons 12 and 13)	81279
JAK2 V617F Targeted Mutation Analysis	81270
KIT D816 Targeted Mutation Analysis	81273
KIT Targeted Sequence Analysis	81272
KRAS Exon 2 Targeted Mutation Analysis	81275
KRAS Targeted Mutation Analysis, Additional Variants	81276
LiquidHALLMARK	0409U
MGMT Promoter Methylation Analysis	81287
MLH1 Sequencing	81292

Procedures addressed by this guideline	Procedure codes
Molecular Tumor Marker Test	81400 81401 81402 81403 81405 81406 81407 81408 81479
Molecular Tumor Marker Test	88271
MPL Common Variants (e.g. W515A, W515K, W515L, W515R)	81338
MPL Mutation Analysis, Exon 10	81339
MSH2 Sequencing	81295
MSH6 Sequencing	81298
NeoLAB Prostate	0011M
Northstar Select	0487U
NPM1 Exon 12 Targeted Mutation Analysis	81310
NRAS Exon 2 and Exon 3 Analysis	81311
NTRK1 Translocation Analysis	81191
NTRK2 Translocation Analysis	81192
NTRK3 Translocation Analysis	81193
NTRK Translocation Analysis	81194
PDGFRA Targeted Sequence Analysis	81314
PMS2 Sequencing	81317
PTEN Sequencing	81321
Resolution ctDx Lung	0179U
RUNX1 Mutation Analysis	81334
SF3B1 Common Variants (e.g. A672T, E622D, L833F, R625C, R625L)	81347

Procedures addressed by this guideline	Procedure codes
Solid Organ Neoplasm, Genomic Sequence Analysis Panel, Cell-free Nucleic Acid (eg, plasma), Interrogation for Sequence Variants; DNA Analysis or Combined DNA and RNA Analysis, Copy Number Variants and Rearrangements	81462
Solid Organ Neoplasm, Genomic Sequence Analysis Panel, Cell-free Nucleic Acid (eg, plasma), Interrogation for Sequence Variants; DNA Analysis, Copy Number Variants, and Microsatellite Instability	81463
Solid Organ Neoplasm, Genomic Sequence Analysis Panel, Cell-free Nucleic Acid (eg, plasma), Interrogation for Sequence Variants; DNA Analysis or Combined DNA and RNA Analysis, Copy Number Variants, Microsatellite Instability, Tumor Mutation Burden, and Rearrangements	81464
SRSF2 Common Variants (e.g. P95H, P95L)	81348
TERT Targeted Sequence Analysis	81345
therascreen PIK3CA RGQ PCR Kit	0177U
TP53 Sequencing	81351
TP53 Targeted Sequence Analysis	81352
U2AF1 Common Variants (e.g. S34F, S34Y, Q157R, Q157P)	81357
ZRSR2 Common Variants (e.g. E65fs, E122fs, R448fs)	81360

Criteria

Introduction

Requests for liquid biopsy testing are reviewed using the following criteria.

Companion Diagnostic (CDx) Liquid Biopsy Assay

Liquid biopsy-based companion diagnostic assays are considered medically necessary when the member meets ALL of the following criteria:

- Member has a diagnosis of cancer, AND
- Treatment with a medication for which there is a liquid biopsy-based FDA-approved companion diagnostic is being considered, AND
- FDA approval for the CDx being requested must include the member's specific cancer type as an approved indication, AND
- FDA label for the drug and indication being considered states companion diagnostic testing is necessary for member selection, AND
- Member has not had previous somatic and/or germline testing that would have identified the genetic change required to prescribe the medication under consideration, AND
- Family history:
 - Member does not have a close (1st or 2nd degree) biological relative with a known germline mutation in a gene that is a target of the requested companion diagnostic test (e.g. known familial mutation in BRCA1/2 and requested test is myChoice CDx), or
 - Member has a close (1st or 2nd degree) biological relative with a known germline mutation in a gene that is a target of the requested companion diagnostic test (e.g. known familial mutation in BRCA1/2 and requested test is myChoice CDx), and the member's germline test was negative, AND
- Rendering laboratory is a qualified provider of service per the Health Plan policy.

Note Not all indications for medications with an FDA-approved companion diagnostic liquid biopsy test require the results of that test prior to prescribing. Testing would not be considered medically necessary when prescribed for indications that do not require the companion diagnostic.

Guardant360 LDT

When Guardant360 LDT (laboratory developed test) is being requested for indications that are outside the scope of a companion diagnostic (i.e.: non-CDx), the panel will be considered medically necessary when the following criteria are met:

- The member has a diagnosis of metastatic or recurrent non-small cell lung cancer (NSCLC), AND
- NSCLC diagnosis has been confirmed based on a histopathologic assessment of tumor tissue, AND
- No previous multi-gene panel testing has been performed for NSCLC, AND

- Insufficient tumor tissue is available for broad molecular profiling and member is unable to undergo an additional standard tissue biopsy due to documented medical reasons (i.e., invasive tissue sampling is contraindicated due to the member's clinical condition)

Liquid Biopsy Tests for Other Non-CDx Indications

These tests are considered Experimental, Investigational, or Unproven.

- Experimental, Investigational, or Unproven (E/I/U) refers to tests, or uses of tests, that have insufficient data to demonstrate an overall health benefit. This typically means there is insufficient data to support that a test accurately assesses the outcome of interest (analytical and clinical validity) and significantly improves patient health outcomes (clinical utility). Such tests are also not generally accepted as the standard of care in the evaluation or management of a particular condition.
- In the case of laboratory testing, FDA approval or clearance is not a reliable standard given the number of laboratory developed tests that currently fall outside of FDA oversight. In addition, FDA approval or clearance often does not include an assessment of clinical utility.

Billing and Reimbursement

Introduction

This section outlines the billing requirements for tests addressed in this guideline. These requirements will be enforced during the case review process whenever appropriate. Examples of requirements may include specific coding scenarios, limits on allowable test combinations or frequency and/or information that must be provided on a claim for automated processing. Any claims submitted without the necessary information to allow for automated processing (e.g. ICD code, place of service, etc.) will not be reimbursable as billed. Any claim may require submission of medical records for post service review.

When otherwise reimbursable, the following will apply:

- When a panel is being performed, it is only reimbursable when billed with a single, appropriate panel procedure code (e.g., 81462, 81463, 81464*).

Note *The panel code(s) listed here may not be all-inclusive. For further discussion of what is considered an appropriate panel code, please refer to the guideline *Laboratory Billing and Reimbursement*.

What is liquid biopsy testing?

Definition

The use of circulating tumor DNA (ctDNA) to identify genetic mutations present in a tumor is also referred to as a liquid biopsy.

- The National Cancer Institute defines a liquid biopsy as “a laboratory test done on a sample of blood, urine, or other body fluid to look for cancer cells from a tumor or small pieces of DNA, RNA, or other molecules released by tumor cells into a person's body fluids. Liquid biopsy allows multiple samples to be taken over time, which may help doctors understand what kind of genetic or molecular changes that are taking place in a tumor.”¹
- Circulating tumor DNA is released into circulation by tumors.² It can be found in various substances, including blood, urine, saliva, etc.
- Analysis of ctDNA can be performed to help identify indicators of disease recurrence or disease progression. It can also help to determine if a specific treatment is indicated.
- Liquid biopsies can be used to more easily obtain serial sampling of a tumor. This is particularly useful since somatic mutations that are used in treatment decisions can change as the tumor progresses.² ctDNA is also thought to provide a more representative sample of the entire tumor genome as well as any metastases that may be present.²
- Traditional methods of performing biopsies on tumor tissue pose the following problems:^{2,3}
 - Biopsies are invasive, involve risks, are typically costly, and are typically difficult to obtain.
 - Treatment decisions often rely on one single biopsy, while tumors are usually heterogeneous in nature, tumor characteristics can evolve, and information regarding metastases may not be known.²
- The use of liquid biopsies can help overcome some of the above problems with traditional biopsies since they can be completed in a noninvasive manner.
- This guideline will only address the use of ctDNA as a liquid biopsy in solid tumors and hematologic malignancies. Circulating tumor cells (CTCs) can be used to help obtain information about an individual's cancer prognosis and treatment options. CTC assays are not addressed by this guideline.

Test information

Introduction

Liquid biopsy testing is an assay that utilizes ctDNA to assist with monitoring disease status and potentially determining sensitivity to certain treatments.

Liquid biopsy methodology

Testing methodology relies on the presence of ctDNA in circulation, which is typically analyzed by one of the following methods:

- Standard testing methodologies, such as polymerase chain reaction (PCR) or sequencing, are used to identify targeted mutations commonly present in tumors of a specific type.
- Methodologies such as next-generation sequencing (NGS) or array comparative genomic hybridization (aCGH) are used to identify both novel and recurrent mutations. These include whole genome sequencing or whole exome sequencing. These approaches analyze single genes, panels of genes, exomes, or genomes. Use of these approaches allows testing with no prior knowledge of genetic mutations that are present in the individual's tumor.
- Several liquid biopsy tests have been designated by the Food and Drug Administration (FDA) as companion diagnostic (CDx) assays deemed necessary for the effective use of a specific medication in the context of a specific clinical indication. Within this guideline, liquid biopsy tests that do not have the designation of companion diagnostics are referred to as non-CDx assays.

Note Tests that extract DNA from nucleated cells in the blood or bone marrow for hematologic malignancies are not considered liquid biopsies. For information on these assays, please refer to the guideline *Somatic Mutation Testing*, as this testing is not addressed here.

Guidelines and evidence

Introduction

This section includes relevant guidelines and evidence pertaining to liquid biopsy testing.

American Society of Clinical Oncology and College of American Pathologists

Based on a comprehensive systematic review of 77 scientific studies on ctDNA assays for solid tumors, an expert panel assembled by the American Society of Clinical Oncology (ASCO, 2018) and the College of American Pathologists (CAP, 2018) concluded that there is currently insufficient evidence of clinical validity and clinical utility for most ctDNA assays being used in advanced cancer.⁴ There are some ctDNA

assays that have demonstrated clinical validity and clinical utility with certain types of cancers, such as non-small cell lung cancer. There is no evidence for use in early stage cancer, treatment monitoring, or residual disease detection. They also stated that there is no evidence of clinical value for cancer screening outside of a clinical trial.

To establish clinical validity and clinical utility of ctDNA analyses, the expert panel recommended the following:

- "Future research studies to establish clinical validity and utility of ctDNA assays should include a patient cohort that matches the intended-use population as closely as possible and samples collected from a prospective study with defined entry criteria. Data will most frequently come from a phase II or phase III study in the patient population where it is anticipated the assay would be used in subsequent clinical practice, with the frequency of the variant under study approximately equal to that in an unselected clinical population. In prospective studies of targeted therapies, the entry criteria should allow inclusion of patients in which the variant under study is observed in the plasma, but not in the tissue analysis, to evaluate the treatment response of this population with discordant genotyping results."

European Society for Medical Oncology

The European Society for Medical Oncology (ESMO, 2022) stated the following regarding liquid biopsies (LBs) for testing in individuals with advanced cancer:⁵

- "LB assays with very high analytical and clinical specificity, and therefore positive predictive values, may be used in routine practice when the results will affect standard treatment options. The limitations of ctDNA assays, however, must be taken into account."
- "...the clinical utility of ctDNA is very much context-dependent, contingent on disease types and stages, available treatment that could effectively eradicate MRD [minimal residual disease] and intended use..."
- Tissue-based testing is the most appropriate test for the majority of individuals, while clinical scenarios exist where ctDNA assays are recommended. These include certain aggressive tumors or when tumor tissue is insufficient or not appropriate.

The guidelines also stated that insufficient evidence exists for implementing use of ctDNA assays for cancer screening, monitoring of treatment response, or detection of molecular relapse or MRD.

International Association for the Study of Lung Cancer

A consensus statement from the International Association for the Study of Lung Cancer (IASLC, 2021) on usage of liquid biopsy for advanced NSCLC made the following recommendations:⁶

- For treatment-naïve individuals with advanced NSCLC, tumor genotyping with a tissue sample should be performed first if available. If the tumor tissue is of

uncertain adequacy or there is concern for incomplete tumor genotyping, ctDNA genotyping can be performed concurrently or as reflex testing. When a tissue sample is unavailable for these individuals, plasma ctDNA genotyping is the recommended test.

- Usage of plasma ctDNA is recommended "[a]t the time of acquired resistance after tyrosine kinase inhibitor (TKI) therapy in an oncogene-driven NSCLC."

National Comprehensive Cancer Network

The National Comprehensive Cancer Network (NCCN, 2024) stated the following regarding liquid biopsies for testing in individuals with non-small cell lung cancer:⁷

- "ctDNA testing should not be used in lieu of a histologic tissue diagnosis."
- "The use of cell-free/circulating tumor DNA testing can be considered in specific clinical circumstances, most notably:"
 - "If a patient is medically unfit for invasive tissue sampling"
 - "In the initial diagnostic setting, if following pathologic confirmation of a NSCLC diagnosis there is insufficient material for molecular analysis, cell-free/circulating tumor DNA can be used; however, follow-up tissue-based analysis for all patients in which an oncogenic driver is not identified should be planned."
 - "In the initial diagnostic setting, if tissue-based testing does not completely assess all recommended biomarkers owing to tissue quantity or testing methodologies available, consider repeat biopsy and/or cell-free/circulating tumor DNA testing."
 - "In the initial diagnostic setting, if the feasibility of timely tissue-based testing is uncertain, concurrent cfDNA [cell-free DNA] testing may aid in biomarker evaluation for treatment selection, provided negative results are considered per above limitations."
- "...the panel feels that plasma ctDNA testing should not be used to diagnose NSCLC; tissue should be used to diagnose NSCLC. Standards and guidelines for plasma ctDNA testing for somatic variants/mutations have not been published, there is up to a 30% false-negative rate, and variants can be detected that are not related to the tumor....careful consideration is required to determine whether ctDNA findings reflect a true oncogenic driver or an unrelated finding."
- "Data suggest that plasma ctDNA testing is a useful minimally invasive test that can be used to identify ALK, BRAF, EGFR, HER2, MET exon 14 skipping, RET, ROS1, and other oncogenic biomarkers that would not otherwise be identified in patients with metastatic NSCLC."
- "The NCCN Guidelines for NSCLC provide recommendations for individual biomarkers that should be tested and recommend testing techniques but do not endorse any specific commercially available biomarker assays or commercial laboratories."

Selected Relevant Publications

Many laboratories are developing liquid biopsy assays. For many of these assays, analytical validity studies have been performed; however, data regarding the clinical validity and clinical utility of these tests is still emerging.^{3,8-41}

The TRACERx study (Tracking Non-small cell lung cancer evolution through therapy (Rx)) is a large, prospective clinical trial being conducted to evaluate “the relationship between intra-tumour heterogeneity and clinical outcome following surgery and adjuvant therapy.”⁴² Researchers plan to analyze the individual's tumors before surgery and multiple times after surgery during their treatment regimen. Tumor tissue and ctDNA in individual's blood will be examined in approximately 840 individuals with NSCLC. This trial is expected to continue until 2026.⁴²

Limited evidence suggests that liquid biopsy with Guardant360, in individuals with advanced NSCLC, may be a reasonable non-invasive alternative to tumor biopsy, particularly in individuals unable to undergo standard tissue biopsy or in cases where tumor tissues are lacking or insufficient for proper mutation analysis.⁴³⁻⁵⁸

Several systematic reviews and meta-analyses have synthesized the findings of multiple studies to evaluate the clinical validity and clinical utility of cell-free circulating tumor DNA (ctDNA) to detect a variety of advanced cancer (excluding non-small cell lung cancer and hematological malignancies).^{8-12,16-41,59-63} With the exception of FDA-approved ctDNA assays, the majority of assays have limited evidence of clinical validity and very limited-to-no evidence of clinical utility for use in individuals with advanced cancer.⁶³ Some studies have also reported relatively high rates of discordance between ctDNA assays and tissue-based testing. There is even less evidence regarding the validity of ctDNA testing in early stage disease, during treatment monitoring, or minimal residual disease (MRD) detection.⁶³ Additional well-designed prospective studies are needed to establish the clinical validity and clinical utility of ctDNA assays before ctDNA assays (liquid biopsy) can be widely adopted in clinical practice.

Note This benefit/harm statement only applies to those jurisdictions that do not have Medicare guidance. Based upon the guidelines and evidence provided in the clinical policy, following EviCore's criteria for liquid biopsy testing will ensure that testing will be available to those members most likely to benefit from the information provided by the assays. For those not meeting criteria, it ensures alternate diagnostic/management strategies are considered. However, it is possible that some members who would benefit from the testing, but do not meet clinical criteria, will not receive an immediate approval for testing.

References

Introduction

These references are cited in this guideline.

1. National Cancer Institute. NCI dictionary of cancer terms: liquid biopsy. Available at: <http://www.cancer.gov/publications/dictionaries/cancer-terms?cdrid=779095>.
2. Corcoran RB, Chabner BA. Application of Cell-free DNA Analysis to Cancer Treatment. *N Engl J Med*. 2018;379(18):1754-1765.
3. Schwaederle M, Husain H, Fantal PT, et al. Detection rate of actionable mutations in diverse cancers using a biopsy-free (blood) circulating tumor cell DNA assay. *Oncotarget*. 2016.
4. Merker JD, Oxnard GR, Compton C, et al. Circulating Tumor DNA Analysis in Patients With Cancer: American Society of Clinical Oncology and College of American Pathologists Joint Review. *J Clin Oncol*. 2018;36(16):1631-1641. doi: 10.1200/jco.2017.76.8671 .
5. Pascual J, Attard G, Bidard FC, et al. ESMO recommendations on the use of circulating tumour DNA assays for patients with cancer: a report from the ESMO Precision Medicine Working Group. *Ann Oncol*. 2022;33(8):750-768. doi:10.1016/j.annonc.2022.05.520
6. Rolfo C, Mack P, Scagliotti GV, et al. Liquid biopsy for advanced NSCLC: A Consensus Statement from the International Association for the Study of Lung Cancer. *J Thorac Oncol*. 2021;16(10):1647-1662. doi:10.1016/j.jtho.2021.06.017
7. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines™) for Non-Small Cell Lung Cancer – v1.2024. © National Comprehensive Cancer Network, Inc. 2024. All rights reserved. Available at: https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf
8. Barata PC, Mendiratta P, Heald B, et al. Targeted next-generation sequencing in men with metastatic prostate cancer: a pilot study. *Target Oncol*. 2018;13(4):495-500.
9. Chung JH, Pavlick D, Hartmaier R, et al. Hybrid capture-based genomic profiling of circulating tumor DNA from patients with estrogen receptor-positive metastatic breast cancer. *Ann Oncol*. 2017;28(11):2866-2873.
10. Schrock AB, Welsh A, Chung JH, et al. Hybrid capture-based genomic profiling of circulating tumor DNA from patients with advanced non-small cell lung cancer. *J Thorac Oncol*. 2019;14(2):255-264.
11. Sokol ES, Feng YX, Jin DX, et al. Loss of function of NF1 is a mechanism of acquired resistance to endocrine therapy in lobular breast cancer. *Ann Oncol*. 2019;30(1):115-123.
12. Zhou C, Yuan Z, Ma W, et al. Clinical utility of tumor genomic profiling in patients with high plasma circulating tumor DNA burden or metabolically active tumors. *J Hematol Oncol*. 2018;11(1):129.

13. Helman E, Nguyen M, Karlovich CA, et al. Cell-Free DNA Next-Generation Sequencing Prediction of Response and Resistance to Third-Generation EGFR Inhibitor. *Clin Lung Cancer*. 2018;19(6):518-530.e517.
14. Wang Z, Cheng Y, An T, et al. Detection of EGFR mutations in plasma circulating tumour DNA as a selection criterion for first-line gefitinib treatment in patients with advanced lung adenocarcinoma (BENEFIT): a phase 2, single-arm, multicentre clinical trial. *Lancet Respir Med*. 2018;6(9):681-690.
15. Park C-K, Cho H-J, Choi Y-D, Oh I-J, Kim Y-C. A Phase II Trial of Osimertinib in the Second-Line Treatment of Non-small Cell Lung Cancer with the EGFR T790M Mutation, Detected from Circulating Tumor DNA: Liquid Lung-O-Cohort 2. *Cancer Res Treat*. 2019;51(2):777-787.
16. Nakamura S, Yokoyama K, Yusa N, et al. Circulating tumor DNA dynamically predicts response and/or relapse in patients with hematological malignancies. *Int J Hematol*. 2018;108(4):402-410.
17. Yan L, Chen Y, Zhou J, Zhao H, Zhang H, Wang G. Diagnostic value of circulating cell-free DNA levels for hepatocellular carcinoma. *Int J Infect Dis*. 2018;67:92-97.
18. Chung JH, Pavlick D, Hartmaier R, et al. Hybrid capture-based genomic profiling of circulating tumor DNA from patients with estrogen receptor-positive metastatic breast cancer. *Ann Oncol*. 2017;28(11):2866-2873.
19. Bunduc S, Gede N, Váncsa S, et al. Prognostic role of cell-free DNA biomarkers in pancreatic adenocarcinoma: A systematic review and meta-analysis. *Crit Rev Oncol Hematol*. 2021;169:103548. doi:10.1016/j.critrevonc.2021.103548.
20. Campo F, Zocchi J, Moretto S, et al. Cell-Free Human Papillomavirus-DNA for Monitoring Treatment Response of Head and Neck Squamous Cell Carcinoma: Systematic Review and Meta-Analysis. *Laryngoscope*. 2021. doi:10.1002/lary.29739.
21. Chen C, Chen C, Sadeghi M. Evaluation of cell-free DNA accuracy as diagnostic biomarker for prostate cancer: A systematic review and meta-analysis. *Biotechnol Appl Biochem*. 2021. doi:10.1002/bab.2149.
22. Chidambaram S, Markar SR. Clinical utility and applicability of circulating tumor DNA testing in esophageal cancer: a systematic review and meta-analysis. *Dis Esophagus*. 2021. doi:10.1093/dote/doab046.
23. de Kruijff IE, Beijer N, Martens JWM, et al. Liquid Biopsies to Select Patients for Perioperative Chemotherapy in Muscle-invasive Bladder Cancer: A Systematic Review. *Eur Urol Oncol*. 2021;4(2):204-214. doi:10.1016/j.euo.2020.01.003.
24. Duque G, Manterola C, Otzen T, et al. Clinical utility of liquid biopsy in breast cancer: A systematic review. *Clin Genet*. 2021. doi:10.1111/cge.14077.

25. Feng SN, Cen XT, Tan R, et al. The prognostic value of circulating tumor DNA in patients with melanoma: A systematic review and meta-analysis. *Transl Oncol*. 2021;14(6):101072. doi:10.1016/j.tranon.2021.101072.
26. Gracie L, Pan Y, Atenafu EG, et al. Circulating tumour DNA (ctDNA) in metastatic melanoma, a systematic review and meta-analysis. *Eur J Cancer*. 2021;158:191-207. doi:10.1016/j.ejca.2021.09.019.
27. Gally TB, Aleluia MM, Borges GF, et al. Circulating MicroRNAs as Novel Potential Diagnostic Biomarkers for Osteosarcoma: A Systematic Review. *Biomolecules*. 2021;11(10). doi:10.3390/biom11101432.
28. Guo Q, Hua Y. The assessment of circulating cell-free DNA as a diagnostic tool for breast cancer: an updated systematic review and meta-analysis of quantitative and qualitative ssays. *Clin Chem Lab Med*. 2021;59(9):1479-1500. doi:10.1515/cclm-2021-0193
29. Guven DC, Sahin TK, Yildirim HC, et al. A systematic review and meta-analysis of the association between circulating tumor DNA (ctDNA) and prognosis in pancreatic cancer. *Crit Rev Oncol Hematol*. 2021;168:103528. doi:10.1016/j.critrevonc.2021.103528.
30. Herranz R, Oto J, Plana E, et al. Circulating Cell-Free DNA in Liquid Biopsies as Potential Biomarker for Bladder Cancer: A Systematic Review. *Cancers (Basel)*. 2021;13(6). doi:10.3390/cancers13061448
31. Hudečková M, Koucký V, Rottenberg J, et al. Gene Mutations in Circulating Tumour DNA as a Diagnostic and Prognostic Marker in Head and Neck Cancer- A Systematic Review. *Biomedicines*. 2021;9(11). doi:10.3390/biomedicines9111548.
32. Huerta M, Roselló S, Sabater L, et al. Circulating Tumor DNA Detection by Digital-Droplet PCR in Pancreatic Ductal Adenocarcinoma: A Systematic Review. *Cancers (Basel)*. 2021;13(5). doi:10.3390/cancers13050994.
33. Jones RP, Pugh SA, Graham J, et al. Circulating tumour DNA as a biomarker in resectable and irresectable stage IV colorectal cancer; a systematic review and meta-analysis. *Eur J Cancer*. 2021;144:368-381. doi:10.1016/j.ejca.2020.11.025.
34. Jongbloed EM, Deger T, Sleijfer S, et al. A Systematic Review of the Use of Circulating Cell-Free DNA Dynamics to Monitor Response to Treatment in Metastatic Breast Cancer Patients. *Cancers (Basel)*. 2021;13(8). doi:10.3390/cancers13081811.
35. Martuszewski A, Paluszkiewicz P, Król M, et al. Donor-Derived Cell-Free DNA in Kidney Transplantation as a Potential Rejection Biomarker: A Systematic Literature Review. *J Clin Med*. 2021;10(2). doi:10.3390/jcm10020193.

36. Morais M, Pinto DM, Machado JC, et al. ctDNA on liquid biopsy for predicting response and prognosis in locally advanced rectal cancer: A systematic review. *Eur J Surg Oncol*. 2021. doi:10.1016/j.ejso.2021.08.034.
37. Palmela Leitão T, Miranda M, Polido J, et al. Circulating tumor cell detection methods in renal cell carcinoma: A systematic review. *Crit Rev Oncol Hematol*. 2021;161:103331. doi:10.1016/j.critrevonc.2021.103331
38. Thusgaard CF, Korsholm M, Koldby KM, et al. Epithelial ovarian cancer and the use of circulating tumor DNA: A systematic review. *Gynecol Oncol*. 2021;161(3):884-895. doi:10.1016/j.ygyno.2021.04.020
39. Uhe I, Hagen ME, Ris F, et al. Cell-free DNA liquid biopsy for early detection of gastrointestinal cancers: A systematic review. *World J Gastrointest Oncol*. 2021;13(11):1799-1812. doi:10.4251/wjgo.v13.i11.1799.
40. Van der Eecken K, Vanwelkenhuyzen J, Deek MP, et al. Tissue- and Blood-derived Genomic Biomarkers for Metastatic Hormone-sensitive Prostate Cancer: A Systematic Review. *Eur Urol Oncol*. 2021;4(6):914-923. doi:10.1016/j.euo.2021.10.005.
41. Xiao H, Gao F, Pang Q, et al. Diagnostic Accuracy of Donor-derived Cell-free DNA in Renal-allograft Rejection: A Meta-analysis. *Transplantation*. 2021;105(6):1303-1310. doi:10.1097/tp.0000000000003443
42. Clinical Trials.gov. Tracking non-small cell lung cancer evolution through therapy (Rx) (TRACERx). Available at: <https://clinicaltrials.gov/ct2/show/NCT01888601>.
43. Mao C, Yuan JQ, Yang ZY, Fu XH, Wu XY, Tang JL. Blood as a substitute for tumor tissue in detecting EGFR mutations for guiding EGFR TKIs treatment of non small cell lung cancer: a systematic review and meta-analysis. *Medicine*. 2015;94(21):e775.
44. Thompson JC, Yee SS, Troxel AB, et al. Detection of therapeutically targetable driver and resistance mutations in lung cancer patients by next-generation sequencing of cell-free circulating tumor DNA. *Clin Cancer Res*. 2016;22(23):5772-5782.
45. Santos ES RL, Castillero LDC, Maran C, Hunis B. Genomic tissue analysis and liquid biopsy profiles from patients diagnosed with advanced adenocarcinoma of the lung. *Clinics in Oncology*. 2016;1:Article 1099.
46. Rozenblum AB, Ilouze M, Dudnik E, et al. Clinical impact of hybrid capture-based next-generation sequencing on changes in treatment decisions in lung cancer. *J Thorac Oncol*. 2017;12(2):258-268.
47. Yang M, Topaloglu U, Petty WJ, et al. Circulating mutational portrait of cancer: manifestation of aggressive clonal events in both early and late stages. *J Hematol Oncol*. 2017;10(1):100.

48. Kim ST BK, Lee SH, Kim K, Park JO, et al. Prospective feasibility study for using cell-free circulating tumor DNA-guided therapy in refractory metastatic solid cancers: an interim analysis. *Precis Oncol.* 2017.
49. Schwaederle M, Chattopadhyay R, Kato S, et al. Genomic alterations in circulating tumor DNA from diverse cancer patients identified by next-generation sequencing. *Cancer Res.* 2017;77(19):5419-5427.
50. Odegaard JI. Validation of a plasma-based comprehensive cancer genotyping assay utilizing orthogonal tissue- and plasma-based methodologies. *Clin Cancer Res.* 2018;24(15):3539-3549.
51. Laufer-Geva S, Rozenblum AB, Twito T, et al. The clinical impact of comprehensive genomic testing of circulating cell-free DNA in advanced lung cancer. *J Thorac Oncol.* 2018;13(11):1705-1716.
52. Aggarwal C, Thompson JC, Black TA, et al. Clinical implications of plasma-based genotyping with the delivery of personalized therapy in metastatic non-small cell lung cancer. *JAMA Oncol.* 2018.
53. Fan G, Zhang K, Ding J, Li J. Prognostic value of EGFR and KRAS in circulating tumor DNA in patients with advanced non-small cell lung cancer: a systematic review and meta-analysis. *Oncotarget.* 2017;8(20):33922-33932.
54. Leighl NB, Page RD, Raymond VM, et al. Clinical utility of comprehensive cell-free DNA analysis to identify genomic biomarkers in patients with newly diagnosed metastatic non-small cell lung cancer. *Clin Cancer Res.* 2019;25(15):4691-4700.
55. Schwaederle MC, Patel SP, Husain H, et al. Utility of genomic assessment of blood-derived circulating tumor DNA (ctDNA) in patients with advanced lung adenocarcinoma. *Clin Cancer Res.* 2017;23(17):5101-5111.
56. Villafior V, Won B, Nagy R, et al. Biopsy-free circulating tumor DNA assay identifies actionable mutations in lung cancer. *Oncotarget.* 2016;7(41):66880-66891.
57. Wang H, Zhou F, Qiao M, et al. The Role of Circulating Tumor DNA in Advanced Non-Small Cell Lung Cancer Patients Treated With Immune Checkpoint Inhibitors: A Systematic Review and Meta-Analysis. *Front Oncol.* 2021;11:671874. doi:10.3389/fonc.2021.671874
58. Wang N, Zhang X, Wang F, et al. The Diagnostic Accuracy of Liquid Biopsy in EGFR-Mutated NSCLC: A Systematic Review and Meta-Analysis of 40 Studies. *SLAS Technol.* 2021;26(1):42-54. doi:10.1177/2472630320939565
59. Bos MK, Angus L, Nasserinejad K, et al. Whole exome sequencing of cell-free DNA - A systematic review and Bayesian individual patient data meta-analysis. *Cancer Treat Rev.* 2020;83:101951. doi:10.1016/j.ctrv.2019.101951.

60. Cervena K, Vodicka P, Vymetalkova V. Diagnostic and prognostic impact of cell-free DNA in human cancers: Systematic review. *Mutat Res*. 2019;781:100-129. doi:10.1016/j.mrrev.2019.05.002.
61. Esagian SM, Grigoriadou G, Nikas IP, et al. Comparison of liquid-based to tissue-based biopsy analysis by targeted next generation sequencing in advanced non-small cell lung cancer: a comprehensive systematic review . *J Cancer Res Clin Oncol*. 2010;146(8):2015-2066. doi:10.1007/s00432-020-03267-x.
62. Cree IA, Uttley L, Buckley Woods H, et al. The evidence base for circulating tumour DNA blood-based biomarkers for the early detection of cancer: a systematic mapping review. *BMC Cancer*. 2017;17(1):697. doi:10.1186/s12885-017-3693-7.
63. Merker JD, Oxnard GR, Compton C, et al. Circulating Tumor DNA Analysis in Patients With Cancer: American Society of Clinical Oncology and College of American Pathologists Joint Review. *Arch Pathol Lab Med*. 2018;142(10):1242-1253. doi:10.5858/arpa.2018-0901-SA.