

Repeat Colonoscopy for Inadequate Preparation (COLON-13)

- Bowel preparation adequacy is defined as BBPS ≥ 6 with each segment score of 2 or 3.
- Colonoscopy as requested within 1 year if bowel preparation is inadequate.

References

1. Calderwood AH, Holub JL, Greenwald DA. Recommendations for follow-up interval after colonoscopy with inadequate bowel preparation in a national colonoscopy quality registry. *Gastrointest Endosc*. 2021. (Epub ahead of print) doi:10.1016/j.gie.2021.09.027.
2. Clark BT, Rustagi T, Laine L. What level of bowel prep quality requires early repeat colonoscopy: systematic review and meta-analysis of the impact of preparation quality on adenoma detection rate. *Am J Gastroenterol*. (2014) 109:1714–23. doi:10.1038/ajg.2014.232.
3. Lieberman DA, Rex DK, Winawer SJ, Giardiello FM, Johnson DA, Levin TR. Guidelines for colonoscopy surveillance after screening and polypectomy: a consensus update by the US Multi-Society Task Force on colorectal cancer. *Gastroenterology*. 2012;143:844-857.
4. Parmar, R, Martel M, Rostom A, Barkun AN. Validated scales of colon cleansing: a systematic review. *Am J Gastroenterol*. 2016;111:197-204.
5. Kewani RN, Crockett SD, Calderwood AH. AGA clinical practice update on strategies to improve quality of screening and surveillance colonoscopy: expert review. *Gastroenterology*. 2021;161(2):701-711.