

Evidence Discussion – Gastric Neuroendocrine Neoplasms (EGD -1.12)

Gastric neuroendocrine neoplasms (g-NENs) are a type of tumor arising from the neuroendocrine cells in the stomach. These tumors can vary in behavior from benign to highly malignant. After resection, regular surveillance is critical to monitor for recurrence or progression.

Following resection, g-NENs can be re-evaluated at frequent intervals via endoscopy initially and then annually. This interval allows for the early detection of any recurrence or new neoplastic growths. Ongoing monitoring helps ensure any changes are identified promptly, allowing for timely intervention and management.

References

1. ENETS Guidelines for the Management of Neuroendocrine Tumors. Available at: <https://link.springer.com/article/10.1007/s12020-022-03143-3>.
2. Exarchou K, Hu H, Stephens NA, Moore AR, Kelly M, Lamarca A, Mansoor W, Hubner R, McNamara MG, Smart H, Howes NR, Valle JW, Pritchard DM. Endoscopic surveillance alone is feasible and safe in type I gastric neuroendocrine neoplasms less than 10 mm in diameter. *Endocrine*. 2022 Oct;78(1):186-196. doi: 10.1007/s12020-022-03143-3. Epub 2022 Jul 27. PMID: 35895180; PMCID: PMC9474380.
3. ESGE Guidelines on Endoscopic Management of Subepithelial Lesions. Available at: <https://www.esge.com/endoscopic-management-of-subepithelial-lesions-esge-guideline>.
4. Evans J, Decker G, Foley K, et al. ASGE Guideline: The role of endoscopy in the management of premalignant and malignant conditions of the stomach. *Gastrointestinal Endosc*. 2015;82(1):1-8. doi:10.1016/j.gie.2015.03.1967
5. Li QL, Zhang YQ, Chen WF, et al. Endoscopic submucosal dissection for foregut neuroendocrine tumors: an initial study. *World J Gastroenterol* 2012; 18: 5799–806.
6. Basuroy R, Srirajaskanthan R, Prachalias A, et al. Review article: the investigation and management of gastric neuroendocrine tumours. *Aliment Pharmacol Ther* 2014;39:1071-84.