

Nucala® (mepolizumab)

When requesting Nucala® (mepolizumab), the individual requiring treatment must be diagnosed with one of the following FDA-approved indications and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-Approved Indications

Nucala is indicated for the:

- Add-on maintenance treatment of adults and pediatric patients aged 6 years and older with severe asthma and with an eosinophilic phenotype.
- Treatment of adult patients with eosinophilic granulomatosis with polyangiitis.
- Treatment of adults and pediatric patients aged 12 years and older with hypereosinophilic syndrome (HES) for at least 6 months without an identifiable non-hematologic secondary cause.
- Add-on maintenance treatment of chronic rhinosinusitis with nasal polyps in adults patients 18 years of age and older with inadequate response to nasal corticosteroids.

Coverage Guidelines

Severe Asthma

The individual must meet all of the following criteria for initial authorization:

- Has a blood eosinophil count of greater than or equal to 150 cells per microliter within the previous 6 weeks or prior to treatment with Nucala or another monoclonal antibody therapy that may lower blood eosinophil levels (e.g., Cinqair, Dupixent, Adbry, Ebglyss, Nemludio, Fasenra, Tezspire, Xolair);
- Has received at least 3 consecutive months of combination therapy with an inhaled corticosteroid and one additional asthma controller/maintenance medication (e.g., monoclonal antibody therapies, inhaled long-acting beta₂-agonists, inhaled long-acting muscarinic antagonists, a combination inhaler containing both an inhaled corticosteroid and a long acting beta₂-agonist);
- Asthma is/was uncontrolled prior to starting Nucala or another monoclonal antibody therapy (e.g., Fasenra, Dupixent, Cinqair, Tezspire, Xolair) defined by ONE of the following:
 - o Has experienced two or more asthma exacerbations requiring treatment with systemic corticosteroids in the previous year; OR
 - o Has experienced one or more asthma exacerbation(s) requiring hospitalization, emergency room visits, or urgent care visits in the previous year; OR
 - o Has a forced expiratory volume in 1 second (FEV₁) less than 80% predicted; OR

- o Has an FEV₁/forced vital capacity (FVC) less than 0.80; OR
 - o Asthma worsens upon tapering of oral corticosteroid therapy;
- Is 6 years of age or older;
- Nucala is prescribed by or in consultation with an allergist, immunologist, or pulmonologist.

The individual must meet all of the following criteria for re-authorization:

- Has already received at least 6 months of Nucala therapy (*NOTE: Patients who have received less than 6 months of therapy or those who are restarting therapy with Nucala should be considered under initial criteria*);
- Continues to receive therapy with one inhaled corticosteroid (ICS) or one ICS-containing combination inhaler (e.g., Flovent Diskus/HFA, ArmonAir RespiClick, Arnuity);
- Has responded to Nucala therapy (e.g., decreased asthma exacerbations; decreased asthma symptoms; decreased hospitalizations, emergency room visits, urgent care visits, or medical clinic visits due to asthma; decreased requirement for oral corticosteroids therapy).

Eosinophilic Granulomatosis with Polyangiitis (EGPA) [Formerly known as Churg-Strauss Syndrome]

The individual must meet all of the following criteria for initial authorization:

- Has active, non-severe disease defined as vasculitis without life- or organ-threatening manifestations (examples of symptoms of non-severe disease include rhinosinusitis, asthma, mild systemic symptoms, uncomplicated cutaneous disease, mild inflammatory arthritis);
- Is currently receiving a systemic corticosteroid (e.g., prednisone) and has been on therapy for a minimum of 4 weeks;
- Has/had a blood eosinophil level of greater than or equal to 150 cells per microliter within the previous 4 weeks or prior to treatment with Nucala or another monoclonal antibody therapy that may lower blood eosinophil levels (e.g., Cinqair, Dupixent, Adbry, Ebglyss, Nemluvio, Fasenra, Tezspire, Xolair);
- Is 18 years of age or older;
- Nucala is prescribed by or in consultation with an allergist, immunologist, rheumatologist, or pulmonologist.

The individual must meet both of the following criteria for re-authorization:

- Has already received at least 9 months of Nucala therapy (*NOTE: Patients who have received less than 9 months of therapy or those who are restarting therapy with Nucala should be considered under initial criteria*);
- Has responded to Nucala therapy (e.g., reduced rate of relapse, reduced corticosteroid dose, reduced eosinophil levels).

Hypereosinophilic Syndrome (HES)

The individual must meet all of the following criteria for initial authorization:

- Has had hypereosinophilic syndrome for at least 6 months;
- Has FIP1L1-PDGFR α -negative disease;
- Does NOT have an identifiable non-hematologic secondary cause of HES (e.g., drug hypersensitivity, parasitic helminth infection, human immunodeficiency virus infection, non-hematologic malignancy);
- Has/had a blood eosinophil level of greater than or equal to 1,000 cells per microliter prior to treatment with any monoclonal antibody therapy that may lower blood eosinophil levels (e.g., Cinqair, Dupixent, Adbry, Ebglyss, Nemludio, Fasenna, Tezspire, Xolair);
- Has tried at least one other treatment for HES (e.g., systemic corticosteroids, hydroxyurea, cyclosporine, imatinib, pegylated-interferon) for a minimum of 4 weeks;
- Is 12 years of age or older;
- Nucala is prescribed by or in consultation with an allergist, immunologist, rheumatologist, or pulmonologist.

The individual must meet both of the following criteria for re-authorization:

- Has already received at least 8 months of Nucala therapy (*NOTE: Patients who have received less than 8 months of therapy or those who are restarting therapy with Nucala should be considered under initial criteria*);
- Has responded to Nucala therapy (e.g., decreased number of flares, improved fatigue, reduced corticosteroid requirements, decreased eosinophil levels).

Chronic Rhinosinusitis with Nasal Polyps

The individual must meet all of the following criteria for initial authorization:

- Has chronic rhinosinusitis with nasal polyps as evidenced by direct examination, endoscopy, or sinus computed tomography (CT) scan;
- Has experienced two or more of the following symptoms for at least 6 months: nasal congestion, nasal obstruction, nasal discharge, and/or reduction/loss of smell;
- Has received at least 4 weeks of therapy with an intranasal corticosteroid and will continue to receive an intranasal corticosteroid concomitantly with Nucala;
- Meets ONE of the following:
 - Has received at least one course of treatment with a systemic corticosteroid for 5 days or more within the previous 2 years; OR
 - Has a contraindication to systemic corticosteroid therapy; OR
 - Has had prior surgery for nasal polyps;
- Is 18 years of age or older;
- Nucala is prescribed by or in consultation with an allergist, immunologist, or an otolaryngologist (ear, nose and throat [ENT] physician specialist).

The individual must meet all of the following criteria for re-authorization:

- Has already received at least 6 months of Nucala therapy (*NOTE: Patients who have received less than 6 months of therapy or those who are restarting therapy with Nucala should be considered under initial criteria*);
- Continues to receive an intranasal corticosteroid therapy;
- Has responded to Nucala therapy (e.g., reduced nasal polyp size, improved nasal congestion, reduced sinus opacification, decreased sino-nasal symptoms, improved sense of smell).

Approval duration:

Initial authorization: HES 8 months; EGPA 9 months; other indications 6 months

Reauthorization: 12 months

Dosing Recommendations

Severe Asthma

Adults and adolescents aged 12 years and older:

The recommended dose of Nucala is 100 mg administered once every 4 weeks by subcutaneous injection.

Pediatric patients aged 6 to 11 years:

The recommended dose of Nucala is 40 mg administered once every 4 weeks by subcutaneous injection.

Eosinophilic Granulomatosis with Polyangiitis; Hypereosinophilic Syndrome

The recommended dose of Nucala is 300 mg administered once every 4 weeks by subcutaneous injection as 3 separate 100-mg injections.

Chronic Rhinosinusitis with Nasal Polyps

The recommended dose of Nucala is 100 mg administered once every 4 weeks by subcutaneous injection.

References

1. Nucala® subcutaneous injection [prescribing information]. Research Triangle Park, NC: GlaxoSmithKline; March 2023.
2. Pavord ID, Korn S, Howarth P, et al. Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet*. 2012;380:651-659.
3. Ortega HG, Liu MC, Pavord ID, et al. Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med*. 2014;371:1198-1207.
4. Bel EH, Wenzel SE, Thompson PJ, et al. Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N Engl J Med*. 2014;371(13):1189-1197.
5. Han JK, Bachert C, Fokkens W, et al. Mepolizumab for chronic rhinosinusitis with nasal polyps (SYNAPSE): a randomized, double-blind, placebo-controlled, phase 3 trial. *Lancet Respir Med*. 2021;9(10):1141-1153.
6. Wechsler ME, Akuthota P, Jayne D. Mepolizumab or placebo for eosinophilic granulomatosis with polyangiitis. *N Engl J Med*. 2017;376(20):1921-1932.
7. Roufosse F, Kahn JE, Rothenberg ME, et al. Efficacy and safety of mepolizumab in hypereosinophilic syndrome: a phase III, randomized, placebo-controlled trial. *J Allergy Clin Immunol*. 2020;146(6):1397-1405.
8. Global Initiative for Asthma. Global strategy for asthma management and prevention. Updated 2024. Available at: <http://www.ginasthma.org>. Accessed on: October 1, 2024.

9. Chung KF, Wenzel SE, Brozek JL, et al. International ERS/ATS guidelines on definition, evaluation and treatment of severe asthma. *Eur Respir J*. 2014;43:343-373.
10. Holguin F, Cardet JC, Chung KF, et al. Management of severe asthma: a European Respiratory Society/American Thoracic Society Guideline. *Eur Respir J*. 2020;55:1900588.
11. Rank MA, Chu DK, Bognanni A, et al. Joint Task Force on Practice Parameters GRADE guidelines for the medical management of chronic rhinosinusitis with nasal polyposis. *J Allergy Clin Immunol*. 2023;151(2):386-398.
12. Peters AT, Spector S, Hsu J, et al. Diagnosis and management of rhinosinusitis: a practice parameter update. *Ann Allergy Asthma Immunol*. 2014;347-385.
13. Dykewicz MS, Wallace DV, Baroody F, et al. Treatment of seasonal allergic rhinitis: an evidenced-based focused 2017 guideline update. *Ann Allergy Asthma Immunol*. 2017;119(6):489-511.
14. Joint Task Force on Practice Parameters: American Academy of Allergy, Asthma and Immunology and the American College of Allergy, Asthma and Immunology. Rhinitis 2020: a practice parameter update. *J Allergy Clin Immunol*. 2020;146:721-767.
15. Rosenfeld RM, Piccirillo JF, Chandrasekhar SS, et al. Clinical practice guideline (update): adult sinusitis. *Otolaryngol Head Neck Surg*. 2015;152(2S):S1-S39.
16. Chung SA, Langford CA, Maz M, et al. 2021 American College of Rheumatology/Vasculitis Foundation guideline for the management of antineutrophil cytoplasmic antibody-associated vasculitis. *Arthritis Rheumatol*. 2021;73(8):1088-1105.
17. Hellmich B, Sanchez-Alamo B, Schirmer JH, et al. EULAR recommendations for the management of ANCA-associated vasculitis: 2022 update. *Ann Rheum Dis*. 2024;83(1):30-47.
18. Shomali W, Gotlib J. World Health Organization and international consensus classification of eosinophilic disorders: 2024 update on diagnosis, risk stratification, and management. *Am J Hematol*. 2024;99(5):946-968.
19. Oldhoff JM, Darsow U, Werfel T, et al. Anti-IL-5 recombinant humanized monoclonal antibody (mepolizumab) for the treatment of atopic dermatitis. *Allergy*. 2005;60(5):693-696.
20. Oldhoff JM, Darsow U, Werfel T, et al. No effect of anti-interleukin-5 therapy (mepolizumab) on the atopy patch test in atopic dermatitis patients. *Int Arch Allergy Immunol*. 2006;141(3):290-294.
21. Kang EG, Narayana PK, Pouliguen IJ, et al. Efficacy and safety of mepolizumab administered subcutaneously for moderate to severe atopic dermatitis. *Allergy*. 2020;75(4):950-953.
22. Pavord ID, Chanez P, Criner GJ, et al. Mepolizumab for eosinophilic chronic obstructive pulmonary disease. *N Engl J Med*. 2017;377(17):1613-1629.
23. GSK reports on outcome of the FDA Advisory Committee on mepolizumab for the treatment of COPD patients on maximum inhaled therapy [press release]. London, UK: GlaxoSmithKline; July 25, 2018. Available at: <https://www.gsk.com/en-gb/media/press-releases/gsk-reports-on-outcome-of-the-fda-advisory-committee-on-mepolizumab-for-the-treatment-of-copd-patients-on-maximum-inhaled-therapy/>. Accessed on April 9, 2024.
24. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: 2024 report. Global Initiative for Chronic Obstructive Lung Disease. Available from: <http://goldcopd.org/>. Accessed on April 9, 2024.
25. Stein ML, Collins MH, Villanueva JM, et al. Anti-IL-5 (mepolizumab) therapy for eosinophilic esophagitis. *J Allergy Clin Immunol*. 2006;118(6):1312-1319.
26. Straumann A, Conus S, Grzonka P, et al. Anti-interleukin-5 antibody treatment (mepolizumab) in active eosinophilic esophagitis: a randomized, placebo-controlled, double-blind trial. *Gut*. 2010;59:21-30.
27. Assa'ad AH, Gupta SK, Collins MH, et al. An antibody against IL-5 reduces numbers of esophageal intraepithelial eosinophils in children with eosinophilic esophagitis. *Gastroenterology*. 2011;141(5):1593-1604.
28. Dellon ES, Peterson KA, Mitlyng BL, et al. Mepolizumab for treatment of adolescents and adults with eosinophilic oesophagitis: a multicenter, randomized, double-blind, placebo-controlled clinical trial. *Gut*. 2023;72(10):1828-1837.
29. Hirano I, Chan ES, Rank MA, et al. AGA Institute and Joint Task Force on Allergy-Immunology Practice Parameters Clinical Guidelines for the Management of Eosinophilic Esophagitis. *Gastroenterology*. 2020;158(6):1776-1786.
30. Wechsler ME, Nair P, Terrier B, et al. Benralizumab versus mepolizumab for eosinophilic granulomatosis with polyangiitis. *New Engl J Med*. 2024;390(10):911-921.