

# CLINICAL GUIDELINES

## Cerebrovascular Intervention

Effective Date: June 2, 2025



---

### Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by EviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

These guidelines include procedures EviCore does not review for Cigna. Please refer to the **Cigna CPT code list** for the current list of high-tech imaging procedures that EviCore reviews for Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five-digit codes, nomenclature and other data are copyright 2025 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

# Table of Contents

Guideline	Page
<b>General Information.....</b>	<b>3</b>
Glossary.....	4
General Information for Vascular Intervention Requests.....	7
<b>Cerebrovascular Endovascular Embolization and Stents.....</b>	<b>9</b>
Cerebrovascular Endovascular Embolization and Stents.....	10
References.....	14

# General Information

Guideline	Page
Glossary.....	4
General Information for Vascular Intervention Requests.....	7

# Glossary

## Terms and abbreviations

<b>Aneurysm</b>	Defined as a diameter 1.5x the normal arterial diameter.
<b>Angioplasty</b>	A procedure that utilizes a catheter with a balloon that is inflated to enlarge a stenotic area.
<b>Ankle-Brachial Index (ABI)</b>	Ratio of the systolic blood pressure (SBP) measured at the ankle to the brachial (arm) SBP.
<b>Atherectomy</b>	A procedure that utilizes a catheter with a sharp blade or laser on the end of the catheter to remove plaque from a blood vessel.
<b>Crescendo TIA</b>	Multiple recurrent episodes of TIA over hours to days.
<b>Critical limb ischemia</b>	Severe stenosis or occlusion in the vessels supplying the lower extremity such that limb loss will result without treatment. Symptoms of critical limb ischemia in the lower extremities include but are not limited to non-healing wounds, gangrene and ischemic rest pain.
<b>Dissection</b>	Disruption of the media layer of the aorta with bleeding within and along the wall of the aorta.
<b>Graft</b>	Synthetic material used to replace or repair a segment of an artery or bypass an occluded segment of artery.
<b>High-grade stenosis</b>	A high grade stenosis is defined as at least a 50% narrowing of an artery.

<b>Ischemic rest pain</b>	Pain arises from severe arterial occlusive disease in the lower extremities such that the patient experiences pain in the distal aspect of the foot and toes while the limb is in the supine position as would occur with sleep. The pain is relieved with the limb in the dependent position or "dangling from the bed" as the limb is depending on gravity to assist with perfusion.
<b>NASCET</b>	North American Symptomatic Carotid Endarterectomy Trial
<b>Pseudo-aneurysm</b>	Outpouching of blood resulting from disruption of the arterial wall with extravasation of blood contained by periarterial connective tissue and not by the arterial wall layers.
<b>PTA</b>	Percutaneous transluminal angioplasty.
<b>Spider veins</b>	Enlarged, tortuous veins that are usually distributed in a web like cluster. These veins are typically <3mm in diameter.
<b>Stent</b>	A metal scaffold placed inside the artery to maintain patency.
<b>Stent-graft</b>	A metal scaffold covered by fabric material placed inside an artery.
<b>Symptomatic carotid stenosis</b>	Characterized by either a transient ischemic attack or cerebrovascular accident that is in the distribution of known severe carotid stenosis, e.g. transient right sided upper and lower extremity paralysis in the setting of 70% left internal carotid artery stenosis.
<b>Symptomatic aneurysm</b>	Unrelenting non-positional back pain in the setting of a known abdominal or thoracic aortic aneurysm. Patients with a symptomatic aneurysm may or may not have evidence of a free or contained rupture. The presence of symptoms indicate impending rupture.
<b>Varicose veins</b>	Enlarged, tortuous veins often caused by incompetent valves. Veins are typically ≥3mm in diameter.

<b>Velocity ratio (V1/V2)</b>	Ratio of peak systolic velocity in the diseased segment of blood vessel demonstrating elevated flow velocities to the peak systolic velocity of blood flow in normal vessel just proximal to area of concern in arteries, or just distal in veins.
<b>Venous reflux</b>	Characterized by incompetent or "leaky" valves that no longer function as one way valves facilitating the flow of blood from the lower extremities to the heart. This results in pooling of blood in the lower extremities leading to distended engorged veins when the lower extremities are in the dependent position as in sitting or standing.

# General Information for Vascular Intervention Requests

PVI.100.A  
v1.0.2025

## Documentation Requirements for Vascular Intervention Requests

Documentation needed to complete a prior authorization request for vascular intervention include **all** of the following:

- Procedure proposed
- Condition being treated
- Detailed documentation of provider-directed conservative treatment, duration and frequency of treatment, and the response to such treatments, if applicable
- Detailed documentation of any previous intervention and the response
- Detailed documentation of location and size of aneurysmal disease, if present
- Detailed documentation regarding nature of the critical limb ischemia: non-healing wound or ischemic rest pain, if applicable
- Recent (within 6 months) written reports of any of the following diagnostic imaging modalities and studies acceptable for purposes of the peripheral vascular intervention guidelines:
  - Ankle-brachial indices, segmental pressures and pulse volume recordings as applicable
  - Duplex ultrasound including carotid, lower extremity and abdominal
  - CTA or CTV abdomen/pelvis with or without lower extremity run-off
  - MRA or MRV abdomen/pelvis with or without lower extremity run-off
  - Arteriogram or venogram
  - IVUS – intravascular ultrasound
- Recent (within 6 months) clinical evaluation documenting:
  - Symptoms (if lifestyle-limiting, detailed documentation regarding quality of life parameters that are affected)
  - Physical exam findings

## Emergent and Urgent Requests

Individuals being evaluated for vascular/endovascular surgery should be screened for the presence of a medical condition that warrants urgent/emergent definitive surgical treatment. Provider directed non-surgical management is **not** required when there is

documentation, supported by imaging studies or clinical assessment, of any of the following urgent/emergent conditions:

- Critical limb ischemia
- Symptomatic carotid stenosis
- Crescendo TIA's (multiple recurrent episodes of TIA over hours to days)
- Symptomatic or ruptured aneurysms

An urgent/emergent request based on 2018 NCQA standards for utilization management occurs when the time frame for making routine or non-life threatening determinations on care **either**:

- Could seriously jeopardize the life, health, or safety of the member or others, due to the member's psychological state
- In the opinion of a practitioner with knowledge of the member's medical or behavioral condition, would subject the member to adverse health consequences without the care or treatment that is the subject of the request.

Procedures to treat arterial disease may be indicated on an intra-operative basis

## Background and Supporting Information

Prior-authorization requests should be submitted at least two weeks prior to the anticipated date of an elective surgery.



# Cerebrovascular Endovascular Embolization and Stents

Guideline	Page
Cerebrovascular Endovascular Embolization and Stents.....	10
References.....	14

# Cerebrovascular Endovascular Embolization and Stents

**PVI.105.A**  
**v1.0.2025**

## Procedures Included

- Endovascular procedures may include:
  - Embolization (including coiling)
  - Balloon angioplasty
  - Stent placement
  - Flow diverters

## Cerebrovascular Embolization and Stent

Procedure Description	CPT®
Transcatheter permanent occlusion or embolization (e.g., for tumor destruction, to achieve hemostasis, to occlude a vascular malformation), percutaneous, any method, central nervous system (intracranial, spinal cord)	61624
Balloon angioplasty, intracranial (e.g. atherosclerotic stenosis), percutaneous [not covered for prophylactic percutaneous transluminal angioplasty of intracranial arteries after aneurysmal subarachnoid hemorrhage] [dual diagnosis needed – subarachnoid hemorrhage and ischemia]	61630
Transcatheter placement of intravascular stent(s), intracranial (e.g. atherosclerotic stenosis), including balloon angioplasty if performed [not covered for prophylactic percutaneous transluminal angioplasty of intracranial arteries after aneurysmal subarachnoid hemorrhage] [dual diagnosis needed – subarachnoid hemorrhage and ischemia]	61635
Transcatheter placement of an intravascular stent(s), open or percutaneous, including radiological supervision and interpretation and including angioplasty within the same vessel, when performed; initial vein	37238
Transcatheter placement of an intravascular stent(s), open or percutaneous, including radiological supervision and interpretation and including angioplasty within the same vessel, when performed; each additional vein (List separately in addition to code for primary procedure)	37239

## Endovascular treatment of intracerebral pathology

### Indications

- Endovascular treatment of intracerebral pathology is medically necessary when recent clinical history as described in **General Information for Vascular Intervention Requests** documents **any** of the following:
  - Unruptured Aneurysms: Treatment is medically necessary at >5mm
  - Ruptured Aneurysms and/or Subarachnoid Hemorrhage at any size
  - Arteriovenous Malformations for any size

### Non-indications

- Endovascular treatment is not considered medically necessary for intracranial atherosclerosis

### Evidence Discussion

Cerebral aneurysm is a bulging, weakened area in the wall of a blood vessel resulting in an abnormal widening or ballooning greater than 50% of the vessel's normal diameter (width).

The causes of aneurysms are varied. They may be congenital or hereditary, or may be caused by other medical conditions or injury.

The risk of rupture for an asymptomatic aneurysm is 1% per year or less, this risk increases with size, size increase over time, family history, and history of prior subarachnoid hemorrhage (SAH). Endovascular treatment options for aneurysm include coil embolization, balloon remodeling, stent-assisted coil embolization, and/or flow diverters. Treatments for ruptured cerebral aneurysms include surgical clipping, endovascular coiling and/or use of flow diverters. Stenting of a ruptured aneurysm is associated with increased morbidity and mortality and is only considered when less risky options are not available. Treatment should be done early to prevent re-rupture. With conservative management, the risk of aneurysm re-bleeding is 20% to 30% in the first month and then approximately 3% per year.

Most brain AVMs are sporadic and do not have an underlying genetic cause. Conservative management, endovascular embolization, radiation and operative resection are four modalities that can be considered in the treatment of brain AVM. These modalities may be performed either in isolation or in combination.

Intracranial atherosclerosis: Stroke or TIA (transient ischemic attack) can be caused by symptomatic intracranial atherosclerosis. For individuals with cerebral ischemia attributable to stenosis of an intracranial artery, the mainstay of treatment consists of risk factor modification and medications. Angioplasty and/or stenting is generally not

recommended given the low rate of stroke on medical management and the inherent peri-procedural risk of endovascular treatment.

## **Middle Meningeal Artery Embolization for Chronic Subdural Hematoma**

### **Indications**

Middle Meningeal Artery embolization (MMAE) is considered medically necessary in an individual with any of the following clinical scenarios:

- Asymptomatic chronic subdural hematoma measuring  $\geq 8$ mm on imaging
- Symptomatic chronic subdural hematoma as an adjunct to surgical decompression
- Chronic subdural hematoma in the setting of a coagulopathy on antiplatelet therapy and/or unable to undergo surgical drainage
- Prophylactic embolization after surgical evacuation (even without evidence of post-operative chronic subdural hematoma recurrence)

### **Evidence Discussion**

Chronic subdural hematoma (cSDH) is a common neurological disorder primarily affecting the elderly and has been associated with substantial morbidity and mortality. Surgical drainage has historically been the definitive management strategy for symptomatic cSDH. This has been associated with significant morbidity and recurrence rates up to 5-30%. Middle meningeal artery embolization (MMAE) was initially reported as an option for treatment of cSDH over two decades ago. Since that time, multiple studies have been performed comparing the safety and efficacy of MMAE compared to surgical drainage. A meta-analysis pooling 20 studies with 1416 patients, 718 who underwent MMAE and 698 who were conventionally managed, reported lower rates of recurrence (4.8% vs 21.5%), reoperation (4.4% vs 16.4%), and complications (1.7%-4.9%), in the groups of patients who underwent MMAE. This has been supported even in patients with thrombocytopenia or on anticoagulation.

A proposed treatment algorithm for cSDH with the use of MMAE has been published. The algorithm supports MMAE alone for asymptomatic (including headaches) cSDH with subdural  $\geq 8$ mm in size and  $\pm$  midline shift. For patients with symptomatic or large cSDH with  $>5$  mm midline shift, MMAE is supported as adjunct to surgery to decrease risk of recurrence.

Additional studies have evaluated the safety of MMAE in the elderly (65-79 years old) and advanced elderly ( $\geq 80$  years) with cSDH, and found it to be safe and effective in management of cSDH.

A recent systematic review and meta-analysis comparing middle meningeal artery (MMA) embolization to conventional management, found that middle meningeal artery embolization decreased treatment failure and the need for surgical rescue without furthering the risk of morbidity and mortality, compared to conventional management. The study authors recommended considering MMA embolization in the management of chronic subdural hematoma.

### **Idiopathic intracranial hypertension**

Treatment of idiopathic intracranial hypertension with venous angioplasty and/or stenting is considered not medically necessary.

# References

PVI.105.A

v1.0.2025

1. Chimowitz MI, Lynn MJ, Derdeyn CP, et al. Stenting versus aggressive medical therapy for intracranial arterial stenosis [published correction appears in *N Engl J Med*. 2012 Jul 5;367(1):93]. *N Engl J Med*. 2011;365(11):993-1003. doi:10.1056/NEJMoa1105335.
2. Chung DY, Abdalkader M, Nguyen TN. Aneurysmal Subarachnoid Hemorrhage. *Neurol Clin*. 2021;39(2):419-442. doi:10.1016/j.ncl.2021.02.006.
3. Connolly ES Jr, Rabinstein AA, Carhuapoma JR, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2012;43(6):1711-1737. doi:10.1161/STR.0b013e3182587839.
4. Etminan N, de Sousa DA, Tiseo C, et al. European Stroke Organization (ESO) guidelines on management of unruptured intracranial aneurysms. *Eur Stroke J*. 2022;7(3):V. doi:10.1177/23969873221099736.
5. Hoh BL, Ko NU, Amin-Hanjani S, et al. 2023 Guideline for the Management of Patients With Aneurysmal Subarachnoid Hemorrhage: A Guideline From the American Heart Association/American Stroke Association. *Stroke*. 2023;54(7):e314-e370. doi:10.1161/STR.0000000000000436.
6. Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association [published correction appears in *Stroke*. 2015 Feb;46(2):e54]. *Stroke*. 2014;45(7):2160-2236. doi:10.1161/STR.0000000000000024.
7. Nguyen TN. Management of Unruptured Intracranial Aneurysms and Brain Arteriovenous Malformations. *Continuum (Minneapolis)*. 2023;29(2):584-604. doi:10.1212/CON.0000000000001247.
8. Tonetti DA, Thomas AJ, Bulsara KR. Middle Meningeal Artery Embolization for Chronic Subdural Hematoma: A Review. *Oper Neurosurg (Hagerstown)*. 2023 May 1;24(5):469-475. doi: 10.1227/ons.0000000000000656.
9. Ironside N, Nguyen C, Do Q, et al. Middle meningeal artery embolization for chronic subdural hematoma: a systematic review and meta-analysis. *J Neurointerv Surg*. 2021;13(10):951-957. doi:10.1136/neurintsurg-2021-017352.
10. Catapano JS, Nguyen CL, Wakim AA, et al. Middle Meningeal Artery Embolization for Chronic Subdural Hematoma. *Front Neurol*. 2020 Oct 20;11:557233. doi: 10.3389/fneur.2020.557233. Erratum in: *Front Neurol*. 2021 Mar 22;12:666701. doi: 10.3389/fneur.2021.666701.
11. Joyce E, Bounajem MT, Scoville J, et al. Middle meningeal artery embolization treatment of nonacute subdural hematomas in the elderly: a multiinstitutional experience of 151 cases. *Neurosurg Focus*. 2020 Oct;49(4):E5. doi: 10.3171/2020.7.FOCUS20518.
12. Sattari SA, Yang W, Shahbandi A, et al. Middle Meningeal Artery Embolization Versus Conventional Management for Patients With Chronic Subdural Hematoma: A Systematic Review and Meta-Analysis. *Neurosurgery*. 2023 Jun 1;92(6):1142-1154. doi: 10.1227/neu.0000000000002365.