



CLINICAL GUIDELINES

CMM-209: Regional Sympathetic Blocks Guidelines

Effective July 1, 2025

VERSION 1.0.2025

EviCore
By **EVERNORTH**

EviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist and/or individual's Primary Care Physician (PCP) may provide additional insight.

EviCore's Clinical Review Criteria (CRC) and related content is made available for the limited uses of: reference; and individual use, only limited to facilitating the determination of medically necessary and appropriate clinical treatment by clinicians for specific delegated patients under their care. The CRC and related content is proprietary information of EviCore, and copyrighted to the full extent of the law. Except as expressly permitted, you may not modify, copy, reproduce, republish, upload, post, transmit, hyperlink to or from, or distribute in any way the CRC, nor may you sell, transfer, distribute, assign, lease, reproduce, or otherwise use the CRC in commerce, in a manner that competes with us or infringes upon our rights, or for any public or commercial endeavor without our prior and express written consent.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2025 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

©2025 EVICORE HEALTHCARE. ALL RIGHTS RESERVED

CMM-209: Regional Sympathetic Blocks

Definitions**General Guidelines****Indications****Non-Indications****Codes (CMM-209)****References (CMM-209)**

Definitions

- **Complex Regional Pain Syndrome (CRPS):** (as defined by the International Association for the Study of Pain [IASP]): a variety of painful conditions following injury which appear regionally having a distal predominance of abnormal findings, exceeding in both magnitude and duration the expected clinical course of the inciting event and often resulting in significant impairment of motor function, and showing variable progression over time. In addition to injury, CRPS can also occur as a result of various medical disorders or illnesses.
- **Regional sympathetic blocks:** (i.e., Stellate Ganglion Blocks and Lumbar Sympathetic Blocks): the injection of local anesthetic along the sympathetic ganglia using image guidance to reduce sympathetic nervous system activity.

General Guidelines

Application of Guideline

- This guideline does not apply to injections/blocks of other autonomic nerves (e.g., sphenopalatine ganglion, carotid sinus, superior hypogastric plexus, celiac plexus, Gasserian ganglion [trigeminal nerve], splanchnic nerve, Ganglion of Impar, rami communicans).
- All regional sympathetic blocks in recalcitrant cases of CRPS should be performed with the intent of facilitating involvement and advancement in an active rehabilitation/functional restoration program.
 - ◆ This is due to insufficient evidence that regional sympathetic blocks (Stellate Ganglion Blocks and Lumbar Sympathetic Chain Blocks) performed as an isolated treatment alter the long-term outcome of CRPS.
- The determination of medical necessity for the performance of regional sympathetic blocks is always made on a case-by-case basis.

Injectates

- Regional sympathetic blocks may only be performed with anesthetic, corticosteroid, and/or contrast agent

Image Guidance

- Stellate Ganglion blocks must be performed using fluoroscopy or ultrasound for image guidance
- Lumbar sympathetic blocks may only be performed using fluoroscopy for image guidance

Frequency & Number of Injections/Procedures

- Only one invasive modality or procedure will be performed on the same date of service.
- When criteria have been met in the **Indications** section, up to 10 regional sympathetic blocks (4 diagnostic, 6 therapeutic) in the prior 12 months are permitted

Indications

Initial Diagnostic Regional Sympathetic Block

- The performance of an initial diagnostic regional sympathetic block is considered **medically necessary** to establish the presence or absence of sympathetically mediated complex regional pain syndrome (CRPS) when **ALL** the following diagnostic criteria for CRPS have been met:
 - ◆ Continuing pain that is disproportionate to any inciting event
 - ◆ Must report at least one (1) of the symptoms in **THREE of the four** following categories:
 - Sensory: reports of hyperesthesia
 - Vasomotor: reports of temperature asymmetry, skin color changes, and/or skin color asymmetry
 - Sudomotor/edema: reports of edema, sweating changes, and/or sweating asymmetry
 - Motor/trophic: reports of decreased range of motion, motor dysfunction (weakness, tremor, dystonia), and/or trophic changes (hair, nails, skin).
 - ◆ Must display at least one (1) sign on physical examination at the time of evaluation in **TWO or MORE** the following categories:
 - Sensory: evidence of hyperalgesia (to pinprick) and/or allodynia (to light touch)
 - Vasomotor: evidence of temperature asymmetry, skin color changes, and/or asymmetry
 - Sudomotor/edema: evidence of edema, sweating changes, and/or sweating asymmetry
 - Motor/trophic: evidence of decreased range of motion, motor dysfunction (weakness, tremor, dystonia), and/or trophic changes (hair, nails, skin)
 - ◆ There is(are) no other medical or psychological diagnoses that are concordant with the presenting symptoms, signs, and results of relevant studies (e.g., imaging, electrodiagnostic testing, laboratory testing, etc.).
 - ◆ Diagnosis is limited to only the extremities and not to the head/face/neck, trunk, perineum/pelvis, or abdominal viscera.

Additional Diagnostic Regional Sympathetic Blocks

- Following a positive initial diagnostic regional sympathetic block, three (3) additional regional sympathetic blocks, performed within the first two (2) weeks of the initial block, may be considered **medically necessary** to diagnose the individual's pain and obtain a therapeutic response.
 - ◆ A positive response to a diagnostic regional sympathetic block is evidenced by at least 50% reduction in pain and improvement in function for the duration of the local anesthetic used.

Therapeutic Regional Sympathetic Blocks

- Therapeutic regional sympathetic blocks are considered **medically necessary** when **ALL** of the following criteria have been met:
 - ◆ There is a documented positive response to the prior therapeutic regional sympathetic block as evidenced by **ALL** of the following:
 - Decreased use of pain medication
 - Increased functional ability (e.g., increased range of motion, strength, and use of the extremity in activities of daily living)
 - Increased tolerance to touch (e.g., decreased allodynia)
 - ◆ Conservative treatment includes **BOTH** of the following:
 - The therapeutic regional sympathetic block is provided as part of a comprehensive pain management program
 - Ongoing participation in an active rehabilitation/functional restoration program
 - ◆ Therapeutic regional sympathetic blocks are performed at a frequency of no more than one time per week
 - ◆ No more than six (6) total therapeutic regional sympathetic blocks are performed in a 12-month period

Non-Indications

Not Medically Necessary

- Regional sympathetic blocks performed without meeting the criteria listed in the **Definitions**, the **General Guidelines**, and the **Indications** sections are considered **not medically necessary**.
- Regional sympathetic blocks performed for a diagnosis of CRPS in the head/face/neck, trunk, perineum/pelvis, or abdominal viscera are considered **not medically necessary**.

Experimental, Investigational, or Unproven (EIU)

- Regional Sympathetic blocks performed by **EITHER** of the following methods are considered **experimental, investigational, or unproven (EIU)**:
 - ◆ Radiofrequency ablation (RFA)
 - ◆ Chemical neurolysis (with phenol [carbolic acid] or ethyl alcohol injections)

Codes (CMM-209)

The inclusion of any code in this table does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

CPT®	Code Description/Definition
64510	Injection, anesthetic agent; stellate ganglion (cervical sympathetic)
64520	Injection, anesthetic agent; lumbar or thoracic (paravertebral sympathetic)

References (CMM-209)

1. Ackerman W, Zhang J. Efficacy of stellate ganglion blockade for the management of type 1 complex regional pain syndrome. *S Med J*. 2006;99(10):1084-1088.
2. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*. 2nd ed. 2008.
3. American Medical Association. *Current Procedural Terminology—Professional Edition*.
4. Benzon HT, Elmofty D, Shankar H, et al. Use of corticosteroids for adult chronic pain interventions: sympathetic and peripheral nerve blocks, trigger point injections - guidelines from the American Society of Regional Anesthesia and Pain Medicine, the American Academy of Pain Medicine, the American Society of Interventional Pain Physicians, the International Pain and Spine Intervention Society, and the North American Spine Society. *Reg Anesth Pain Med*. doi:10.1136/rapm-2024-105593.
5. Cepeda M, Carr D, Lau J. Local anesthetic sympathetic blockade for complex regional pain syndrome. *Cochrane Database Syst Rev*. 2005;4:CD004598. doi:10.1002/14651858.cd004598.pub2.
6. Cepeda M, Lau J, Carr D. Defining the therapeutic role of local anesthetic sympathetic blockade in complex regional pain syndrome: a narrative and systematic review. *Clin J Pain*. 2002;18:216-233.
7. Chou R, Huffman L. American Pain Society. American College of Physicians. Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med*. 2007;147(7):505-514.
8. Chou R, Huffman LH American Pain Society. American College of Physicians. Nonpharmacologic therapies for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med*. 2007;147(7):492-504.
9. Chou R, Qaseem A, Snow V, et al. Clinical Efficacy Assessment Subcommittee of the American College of Physicians. American College of Physicians. American Pain Society Low Back Pain Guidelines Panel. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. 2007;147(7):478-491.
10. Chou R. Using evidence in pain practice: Part I: Assessing quality of systematic reviews and clinical practice guidelines. *Pain Med*. 2008;9(5):518-530.
11. Dworkin R, O'Connor A, Kent J et al. Interventional management of neuropathic pain: NeuPSIG recommendations. *Pain*. 2013;154(11):2249-2261. doi:10.1016/j.pain.2013.06.004.
12. Forouzanfar T, Köke A, van Kleef M, Weber W. Treatment of complex regional pain syndrome type I. *Eur J Pain*. 2002;6:105-122.
13. Furlan A, Mailis A, Papagapiou M. Are we paying a high price for surgical sympathectomy? A systemic literature review of late complications. *J Pain*. 2000;1:245-257.
14. Galer B, Bruehl S, Harden R. IASP diagnostic criteria for complex regional pain syndrome: a preliminary empirical validation study. *Clin J Pain*. 1998;14:48-54.
15. Gofeld M, Shankar H, Benzon H. Chapter 84: Fluoroscopy and Ultrasound-Guided Sympathetic Blocks. Section IX: Other Nerve Blocks in Pain Medicine. In: *Essentials of Pain Medicine*. 4th ed. Elsevier. 2018:789-804.e2.
16. Harden RN, McCabe CS, Goebel A, et al. Complex Regional Pain Syndrome: Practical Diagnostic and Treatment Guidelines, 5th ed. *Pain Med*. 2022;23(Suppl 1):S1-S53. doi:10.1093/pm/pnac046.
17. Hartrick C, Kovan J, Naismith P. Outcome prediction following sympathetic block for complex regional pain syndrome. *Pain Pract*. 2004;4:222-228.
18. Harvey AM. Classification of chronic pain—descriptions of chronic pain syndromes and definitions of pain terms. *Clin J Pain*. 1995;11(2):179.
19. Janig W, Baron R. Complex regional pain syndrome: mystery explained? *Lancet Neurol*. 2003;2:687-697.
20. Leis S, Weber M, Schmelz M, Birklein F. Facilitated neurogenic inflammation in unaffected limbs of patients with complex regional pain syndrome. *Neurosci Lett*. 2004;359:163-166.
21. O'Connell N, Wand B, Gibson W, Carr D, Birklein F, Stanton T. Local anaesthetic sympathetic blockade for complex regional pain syndrome. *Cochrane Database Syst Rev*. 2016. doi:10.1002/14651858.cd004598.pub4.
22. Ozturk E, Mohur H, Arslan N, et al. Quantitative three-phase bone scintigraphy in the evaluation of intravenous regional blockade treatment in patients with stage-I reflex sympathetic dystrophy of upper extremity. *Ann of Nucl Med*. 2004;18(8):653-658.
23. Paraskevas K, Michaloglou A, Briana D, Samara M. Treatment of complex regional pain syndrome type I of the hand with a series of intravenous regional sympathetic blocks with guanethidine and lidocaine. *Clin Rheumatol*. 2005;7:1-7.
24. Perez R, Kwakkel G, Zuurmond W, de Lange J. Treatment of reflex sympathetic dystrophy (CRPS type 1): a research synthesis of 21 randomized clinical trials. *J Pain Symptom Manage*. 2001;21:511-526.
25. Schurmann M, Gradl G, Wizgal I, et al. Clinical and physiologic evaluation of stellate ganglion blockade for complex regional pain syndrome type I. *Clin J Pain*. 2001;17:94-100.
26. Severens J, Oerlemans H, Weegels A, et al. Cost-effectiveness analysis of adjuvant physical or occupational therapy for patients with reflex sympathetic dystrophy. *Arch Phys Med Rehabil*. 1999;80:1038-1043.

27. Sharma A, Williams K, Raja S. Advances in treatment of complex regional pain syndrome: recent insights on a perplexing disease. *Curr Opin Anaesthesiol*. 2006;19:566-572.
28. Stanton-Hicks M. A report on the 2nd IASP Research Symposium, Cardiff, Wales. Complex regional pain syndrome: current research on mechanisms and diagnosis. In: *International Association for the Study of Pain*. Special Interest Group on Pain and the Sympathetic Nervous System. 2000;1–2.
29. Stanton-Hicks M. In: Wakefield CA, Bajwa JH, eds. *Principles and Practice of Pain Medicine*. 2nd ed. 2004.
30. Stanton-Hicks M, Baron R, et al. Consensus report: complex regional pain syndromes: guidelines for therapy. *Clin J Pain*. 1998;14:155–166.
31. Stanton-Hicks M, Burton A, Bruehl S, et al. An updated interdisciplinary clinical pathway for CRPS: report of an expert panel. *Pain Pract*. 2005;84(3):S4-S16.
32. Stanton-Hicks M. Complex regional pain syndrome: manifestations and the role of neurostimulation in its management. *J Pain Symptom Manage*. 2006;(4 Suppl):S20-S24.
33. Straube S, Derry S, Moore R, McQuay H. Cervico-thoracic or lumbar sympathectomy for neuropathic pain and complex regional pain syndrome. *Cochrane Database Syst Rev*. 2010. doi:10.1002/14651858.cd002918.pub2.
34. Suresh S, Wheeler M, Patel A. Case series: IV regional anesthesia with ketorolac and lidocaine: is it effective for the management of complex regional pain syndrome 1 in children and adolescents? *Anesth Analg*. 2003;96:694-695.
35. Turner J, Loeser J, Deyo R, Sanders S. Spinal cord stimulation for patients with failed back surgery syndrome or complex regional pain syndrome: a systematic review of effectiveness and complications. *Pain*. 2004;108:137-147.
36. Varrassi G, Paladini A, Marinangeli F, Racz G. Neural modulation by blocks and infusions. *Pain Pract*. 2006;6:34-38.
37. Wang L, Chen H, Chang P, et al. Axillary brachial plexus block with patient controlled analgesia for complex regional pain syndrome type I: a case report. *Reg Anesth Pain Med*. 2001;26(1):68-71.
38. Workloss Data Institute. Official Disability Guidelines 2009.
39. Zernikow B, Wager J, Brehmer H, Hirschfeld G, Maier C. Invasive Treatments for Complex Regional Pain Syndrome in Children and Adolescents. *Anesthesiology*. 2015;122(3):699-707. doi:10.1097/aln.0000000000000573.