

Cigna Medical Coverage Policies – Musculoskeletal Spinal Cord and Dorsal Root Ganglion Stimulation

Effective July 1, 2025



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by eviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five digit codes, nomenclature and other data are copyright 2025 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

©Copyright 2025 eviCore healthcare

CMM-211: Spinal Cord and Dorsal Root Ganglion Stimulation

Definitions

General Guidelines

Indications

Replacement

Non-Indications

Appendices

Codes (CMM-211)

References (CMM-211)

Definitions

- **Complex Regional Pain Syndrome (CRPS)** (as defined by the International Association for the Study of Pain [IASP]): a variety of painful conditions following injury which appear regionally having a distal predominance of abnormal findings, exceeding in both magnitude and duration the expected clinical course of the inciting event and often resulting in significant impairment of motor function, and showing variable progression over time. In addition to injury, CRPS can also occur as a result of various medical disorders or illnesses.
- **Critical Limb Ischemia (CLI)**: clinical syndrome of ischemic pain at rest and ischemic tissue loss such as non-healing ulcers or gangrene, related to peripheral artery disease (PAD) of the lower limbs. Spinal stimulators may be appropriate for the treatment of intractable rest pain secondary to chronic limb ischemia.
 - ◆ **Ischemic Rest Pain**: pain that occurs in the toes or in the area of the metatarsal heads. Occasionally, it occurs in the foot proximal to the metatarsal heads. Elevation of the limb above or at the horizontal position aggravates the pain and pendency, to some degree at least, brings relief. The pain is secondary to severe arterial insufficiency resulting in inadequate perfusion to the distal lower extremity.
- **Dorsal Root Ganglion (DRG) Stimulation**: an emerging method of treatment for neuropathic pain. With DRG stimulation, leads are placed percutaneously into the epidural space under fluoroscopic guidance directly over the targeted dorsal root ganglion within the lumbar or sacral region of the spine. The procedure initially involves a short-term trial (i.e., greater than 48 hours) using an external pulse generator; upon success of the trial a permanent pulse generator may then be implanted.
 - ◆ At this time, the evidence in the peer-reviewed scientific literature is insufficient to support long-term safety and efficacy. The use of this technology for treatment of pain conditions remains under investigation.
- **Failed Back Surgery Syndrome (FBSS)**: lumbar spinal pain of unknown origin despite surgical intervention or appearing after surgical intervention for spinal pain originally in the same spinal region. Procedures/surgery that do not encroach into the spinal canal (e.g., interspinous/interlaminar/facet distraction, kyphoplasty/vertebroplasty surgery, etc.) are not considered surgical interventions associated with FBSS.
- **High-Frequency Spinal Cord Stimulation (HF-SCS)** (also referred to as kilohertz frequency spinal cord stimulation or HF10): a type of spinal cord stimulation (SCS) providing a higher frequency than traditional spinal cord stimulator systems. The HF10 SCS uses low-amplitude, high-frequency, and short-duration pulses. HF10 SCS does not generate paresthesia and operates at a frequency of 10,000 Hz to provide pain relief in comparison to traditional spinal cord stimulation systems which operate at a frequency in the range of 40-60 Hz and do generate paresthesia. As an alternative to traditional dorsal spinal column stimulation, HF10 SCS is proven safe and effective for treatment of chronic, intractable low back and leg pain in individuals with failed back surgery syndrome (FBSS).

- **Peripheral Nerve Field Stimulation:** a technology that involves placement of electrodes subcutaneously within an area of maximal pain, with the objective of stimulating a region of affected nerves to reduce pain. Depending on the targeted nerve, leads may be placed percutaneously just under the skin or via an open approach for larger deeper peripheral nerves.
 - ◆ The use of this technology (used alone or in combination with spinal cord stimulation) for treatment of pain conditions is under investigation.
- **Peripheral Nerve Stimulation:** involves implantation of electrodes near or on a peripheral nerve to reduce pain.
 - ◆ The use of this technology (used alone or in combination with spinal cord stimulation) for treatment of pain conditions is under investigation.
- **Spinal Cord Stimulation (SCS)** (also known as dorsal column stimulation or neuromodulation): a reversible therapy applied for neuropathic pain with techniques that include multi-output implanted pulse generators and a choice of electrodes, some of which can be placed percutaneously. The technical goal of this therapy is to achieve stimulation of paresthesia of the dorsal horn of the spinal cord at a subjectively comfortable level, overlapping an individual's topography of pain. The procedure initially involves a short-term trial (e.g., greater than 48 hours) of percutaneous (temporary) spinal cord stimulation, prior to the subcutaneous (permanent) implantation of the spinal cord stimulation device, to determine whether the spinal cord stimulator device will induce sufficient pain relief to render it medically necessary.

General Guidelines

Application of Guideline

- A dorsal column stimulator capable of **using either high-frequency or non-high-frequency stimulation** (dual-mode) is considered an equally effective alternative (when the device uses non-high-frequency stimulation) for the treatment of any of the conditions listed in the **Indications** section below.
- A dorsal column stimulator **using high frequency** is considered an equally effective alternative to non-high-frequency stimulation **only** for the treatment of chronic intractable pain secondary to failed back surgery syndrome (FBSS).
- This guideline **does not apply** to simple or complex brain or peripheral (i.e., cranial nerve, peripheral nerve, autonomic nerve, neuromuscular) neurostimulator pulse generator/transmitter.
- The determination of medical necessity for implantation of a dorsal column spinal cord stimulator (SCS) is always made on a case-by-case basis.

Indications

Chronic Intractable Pain Secondary to Failed Back Surgery Syndrome (FBSS)

Trial

- A short-term trial (i.e., greater than 48 hours) spinal cord stimulation (i.e., non-high-frequency **or** high-frequency [HF10 SCS]) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to failed back surgery syndrome (FBSS) with intractable neuropathic leg pain (after prior surgery in the same spinal region) when **ALL** of the following criteria are met:
 - ◆ There has been a failure of at least six (6) consecutive months of physician-supervised conservative medical management (e.g., pharmacotherapy, physical therapy, cognitive behavioral therapy, or activity lifestyle modification)
 - ◆ Surgical intervention is not indicated **or** the individual does not wish to proceed with spinal surgery
 - ◆ An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental and/or behavioral health conditions/issues (e.g., substance use disorders, depression, or psychosis) that would impact perception of pain, and/or negatively impact the success of a SCS or contraindicate placement of the device
- If the initial short-term trial fails, a repeat trial is considered not **medically necessary**.

Permanent Implant

- Permanent implantation of a spinal cord stimulator (i.e., non-high-frequency or high-frequency [HF10 SCS]) is considered **medically necessary** when **BOTH** of the following criteria are met:
 - ◆ Must meet **ALL** criteria for the short-term trial spinal cord stimulation (SCS) as noted above
 - ◆ There has been at least a 50% reduction in pain during a short-term trial of SCS

Complex Regional Pain Syndrome (CRPS)/Reflex Sympathetic Dystrophy (RSD)

Trial

- A short-term trial (i.e., greater than 48 hours) of a non-high-frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD) only of the upper and lower extremities when **ALL** of the following criteria are met:
 - ◆ Limited to only the extremities and not to the head/face/neck, trunk, perineum/pelvis, or abdominal viscera.

- ◆ Diagnosis of CRPS/RSD as evidenced by **ALL** of the following:
 - Continuing pain which is disproportionate to any inciting event
 - Must report at least **ONE** (1) of the symptoms in **THREE of the four** following categories:
 - Sensory: reports of hyperesthesia
 - Vasomotor: reports of temperature asymmetry, skin color changes, and/or skin color asymmetry
 - Sudomotor/edema: reports of edema, sweating changes, and/or sweating asymmetry
 - Motor/trophic: reports of decreased range of motion, motor dysfunction (weakness, tremor, dystonia), and/or trophic changes (hair, nail, skin).
 - Must display at least **ONE** (1) sign on physical examination at the time of evaluation in **TWO or MORE** of the following categories:
 - Sensory: evidence of hyperalgesia (to pinprick) and/or allodynia (to light touch)
 - Vasomotor: evidence of temperature asymmetry, skin color changes, and/or asymmetry
 - Sudomotor/edema: evidence of edema, sweating changes, and/or sweating asymmetry
 - Motor/trophic: evidence of decreased range of motion, motor dysfunction (weakness, tremor, dystonia) and/or trophic changes (hair, nail, skin).
 - There is(are) no other medical or psychological diagnoses that are concordant with the presenting symptoms, signs, and results of relevant studies (e.g., imaging, electrodiagnostic testing, laboratory testing, etc.).
 - ◆ Failure of at least six (6) consecutive months of physician-supervised conservative medical management (e.g., pharmacotherapy, physical therapy, cognitive behavioral therapy, or activity lifestyle modification)
 - ◆ Surgical intervention is not indicated
 - ◆ An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment [with or without psychological questionnaires and/or psychological testing]) reveals no evidence of an inadequately controlled mental and/or behavioral health conditions/issues (e.g., substance use disorders, depression, or psychosis) that would impact perception of pain, and/or negatively impact the success of a SCS or contraindicate placement of the device
- If the initial short-term trial fails, a repeat trial is considered **not medically necessary**

Permanent Implant

- Permanent implantation of a non-high-frequency dorsal column SCS is considered **medically necessary** when **BOTH** of the following criteria are met:
 - ◆ Must meet **ALL** criteria for the short-term trial spinal cord stimulation (SCS) as noted above
 - ◆ There has been at least a 50% reduction in pain during a short-term trial of SCS

Chronic Critical Limb Ischemia (CLI)

Trial

- A short-term trial (i.e., greater than 48 hours) of a non-high-frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to chronic critical limb ischemia (CLI) when **ALL** of the following criteria are met:
 - ◆ Attestation from a vascular surgeon that the individual is not a suitable candidate for vascular reconstruction.
 - ◆ Diagnosis of critical limb ischemia when **ALL** of the following criteria are met:
 - Ischemic limb rest pain
 - Rutherford Classification Grade II, Category 4 (see table in **Appendix B**) ischemic rest pain characterized by **BOTH** of the following:
 - Resting ankle pressure <40mmHg, flat or barely pulsatile ankle or metatarsal pulse volume recording
 - Toe pressure <30mmHg
 - ◆ Advanced Imaging
 - Angiographic or CT/MR imaging demonstrating multilevel disease with absence of named vessel with flow into the foot
 - ◆ An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental and/or behavioral health conditions/issues (e.g., substance use disorders, depression, or psychosis) that would impact perception of pain, and/or negatively impact the success of a SCS or contraindicate placement of the device
- If the initial short-term trial fails, a repeat trial is considered **not medically necessary**

Permanent Implant

- Permanent implantation of a non-high-frequency dorsal column SCS is considered **medically necessary** when **BOTH** of the following criteria are met:
 - ◆ Must meet **ALL** criteria for the short-term trial spinal cord stimulation (SCS) as noted above
 - ◆ There has been at least a 50% reduction in pain during a short-term trial of SCS

Chronic Stable Angina Pectoris

Trial

- A short-term trial (i.e., greater than 48 hours) of a non-high-frequency dorsal column spinal cord stimulator (SCS) is considered **medically necessary** for the treatment of chronic, intractable pain secondary to chronic stable angina pectoris/myocardial ischemia when **ALL** of the following criteria are met:
 - ◆ Angina pectoris is Canadian Cardiovascular Society (CCS) functional class III or class IV (see **Appendix A**)
 - ◆ Attestation by the individual's treating cardiologist confirms coronary artery disease (CAD) AND the individual is not a suitable candidate for a revascularization procedure
 - ◆ Optimal medical treatment (OMT) that has failed to adequately improve anginal symptoms including **ALL** of the following:
 - Anti-platelet therapy
 - Statin and/or other lipid-lowering therapy
 - Anti-anginal therapy implemented to pursue a goal heart rate of 60 beats per minute
 - Anti-hypertensive therapy as may be indicated to pursue a goal systolic blood pressure (SBP) of less than 140 mmHG and a goal diastolic blood pressure (DBP) of less than 90 mmHG
 - ◆ An attestation by a behavioral health provider (i.e., a face-to-face or virtual assessment with or without psychological questionnaires and/or psychological testing) reveals no evidence of an inadequately controlled mental and/or behavioral health conditions/issues (e.g., substance use disorders, depression, or psychosis) that would impact perception of pain and/or negatively impact the success of a SCS or contraindicate placement of the device
- If the initial short-term trial fails, a repeat trial is considered **not medically necessary**.

Permanent Implant

- Permanent implantation of a non-high-frequency dorsal column SCS is considered **medically necessary** when **BOTH** of the following criteria are met:
 - ◆ Must meet **ALL** criteria for the short-term trial spinal cord stimulation (SCS) as noted above
 - ◆ There has been a beneficial clinical response during a short-term trial of SCS

Replacement

Replacement of Dorsal Column Spinal Cord Stimulator

- Replacement of an existing dorsal column spinal cord stimulator (high-frequency or non-high-frequency) is considered **medically necessary** when **ANY** of the following criteria are met:
 - ◆ The existing stimulator and/or battery/generator is malfunctioning, cannot be repaired, and is no longer under warranty

- ◆ Revision is required of the electrode percutaneous array(s)

Replacement of Dorsal Root Ganglion (DRG) Stimulator

Placement of a dorsal root ganglion (DRG) stimulation is considered **not medically necessary** for **ALL** indications except as noted below.

- Replacement of an existing dorsal root ganglion (DRG) stimulator with another dorsal root ganglion (DRG) stimulator is considered **medically necessary** when **ANY** of the following criteria are met:
 - ◆ The existing stimulator and/or battery/generator is malfunctioning, cannot be repaired, and is no longer under warranty
 - ◆ Revision is required of the electrode percutaneous array(s) or electrode plate/paddle(s)

Non-Indications

Not Medically Necessary

Spinal Cord Stimulation (SCS)

- Spinal cord stimulation (high-frequency or non-high-frequency) placed without meeting the requirements listed in the **Definitions**, the **General Guidelines**, and the **Indications** sections is considered **not medically necessary**.
- Replacement of a **functioning non-high-frequency** dorsal column SCS with a **high-frequency** SCS is considered **not medically necessary**.
- A high-frequency spinal cord stimulator (SCS) is considered **not medically necessary** for **ANY** other indication, including complex regional pain syndrome (CRPS)/reflex sympathetic dystrophy (RSD).
- A high-frequency **or** non-high-frequency dorsal column SCS is considered **not medically necessary** for **ANY** other indication including, but not limited to, the following conditions:
 - ◆ Post-amputation pain (phantom limb pain)
 - ◆ Post-herpetic neuralgia
 - ◆ Peripheral neuropathy (e.g., chronic intractable pain from diabetic sensory neuropathy)
 - ◆ Dysesthesias involving the lower extremities secondary to spinal cord injury
 - ◆ Abdominal/pelvic visceral pain
 - ◆ Chronic cervical or lumbar radiculopathy without prior spinal surgery
 - ◆ Chronic cervical, thoracic, or lumbar axial pain without prior spinal surgery
 - ◆ Failed cervical and/or thoracic spinal surgery with intractable neuropathic pain in arm(s) or trunk.
 - ◆ Abdominal pain related to celiac artery compression syndrome
 - ◆ Neuropathic pain associated with Multiple Sclerosis

- Generator modes **other than tonic-low and high-frequency** (e.g., burst-stimulation) are considered **not medically necessary**.

Dorsal Root Ganglion (DRG) Stimulator

- Replacement of a dorsal column spinal cord stimulator (SCS) with a dorsal root ganglion (DRG) stimulator is considered **not medically necessary**.
- Dorsal root ganglion (DRG) stimulation is considered **not medically necessary** for **ALL** indications except as noted above in **Replacement**.

Appendices

Appendix A

Classifications of Cardiovascular Disability

Class	New York Heart Association Functional Classification	Canadian Cardiovascular Society Functional Classification
I	Patients with cardiac disease but without resulting limitations of physical activity. Ordinary physical activity does not cause undue fatigue, palpitation, dyspnea, or anginal pain.	Ordinary physical activity does not cause angina, such as walking and climbing stairs. Angina occurs with strenuous or rapid or prolonged exertion at work or recreation.
II	Patients with cardiac disease resulting in slight limitation of physical activity. They are comfortable at rest. Ordinary physical activity results in fatigue, palpitation, dyspnea, or anginal pain.	Slight limitation of ordinary activity. Walking or climbing stairs rapidly, walking uphill, walking or stair climbing after meals, in cold, in wind, or under emotional stress, or only during the few hours after awakening. Walking more than two blocks on the level and climbing more than one flight of ordinary stairs at a normal pace and in normal conditions.
III	Patients with cardiac disease resulting in marked limitation of physical activity. They are comfortable at rest. Less than ordinary physical activity causes fatigue, palpitation, dyspnea, or anginal pain.	Marked limitation of ordinary physical activity. Walking one to two blocks on the level and climbing one flight in normal conditions and at a normal pace.
IV	Patient with cardiac disease resulting in inability to carry on any physical activity without discomfort. Symptoms of cardiac insufficiency or of the anginal syndrome may be present even at rest. If any physical activity is undertaken, discomfort is increased.	Inability to carry on any physical activity without discomfort—anginal syndrome may be present at rest.
(Heart Failure Society of America [HFSA], 2006; Gibbons, et al., 2002; American Heart Association [AHA], 1994; Canadian Cardiovascular Society [CCS], 1976).		

Appendix B

Rutherford Classification System for Staging Ischemia

Grade	Category	Clinical Description	Objective Criteria
0	0	Asymptomatic- no hemodynamically significant occlusive disease	Normal treadmill or reactive hyperemia test
	1	Mild claudication	Completes treadmill exercise; AP after exercise > 50 mmHg, but at least 20 mmHg lower than resting value
I	2	Moderate claudication	Between categories 1 and 3
	3	Severe claudication	Cannot complete standard treadmill exercise and AP after exercise < 50 mmHg
II	4	Ischemic rest pain	Resting AP < 40 mmHg, flat or barely pulsatile ankle or metatarsal PVR; TP < 30mmHg
III	5	Minor tissue loss non-healing ulcer, focal gangrene with diffuse pedal ischemia	Resting AP < 60 mmHG, ankle or metatarsal PVR flat or barely pulsatile; TP < 40 mmHg
	6	Major tissue loss- extending above TM level, functional foot no longer salvageable	Same as category 5

AP: ankle pressure; PVR: pulse volume recording; TM: transmetatarsal; TP: toe pressure

Codes (CMM-211)

The inclusion of any code in this table does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

Code	Code Description/Definition
63650	Percutaneous implantation of neurostimulator electrode array, epidural
63655	Laminectomy for implantation of neurostimulator electrodes, plate/paddle, epidural
63685	Insertion or replacement of spinal neurostimulator pulse generator or receiver, requiring pocket creation and connection between electrode array and pulse generator or receiver

This list may not be all-inclusive and is not intended to be used for coding/billing purposes. The final determination of reimbursement for services is the decision of the health plan and is based on the individual's policy or benefit entitlement structure as well as claims processing rules.

References (CMM-211)

1. Abraham M, Potdar A, Ward M, Herschman Y, Mammis A. Risk Factors Associated with Lead Migration Requiring Revision in Dorsal Root Ganglion Stimulation. *World Neurosurg*. 2019;128:e649-e652. doi:10.1016/j.wneu.2019.04.225.
2. Aiyer R, Barkin RL, Bhatia A, Gungor S. A systematic review on the treatment of phantom limb pain with spinal cord stimulation. *Pain Management*. 2017;7(1):59-69. doi:10.2217/pmt-2016-0041.
3. Al-Kaisy A, Van Buyten JP, Smet I, et al. Sustained effectiveness of 10 kHz high-frequency spinal cord stimulation for patients with chronic, low back pain: 24-month results of a prospective multicenter study. *Pain Med*. 2014;15(3):347-354.
4. Amann W, Berg P, Gersbach P, et al; European Peripheral Vascular Disease Outcome Study SCS- EPOS. Spinal cord stimulation in the treatment of non-reconstructable stable critical leg ischaemia: results of the European Peripheral Vascular Disease Outcome Study (SCS-EPOS). *Eur J Vasc Endovasc Surg*. 2003;26(3):280-286.
5. American College of Occupational and Environmental Medicine. *Occupational Medicine Practice Guideline*. 2nd ed. 2008.
6. American Medical Association. 2024 AMA CPT® Professional Edition.
7. Baber Z, Erdeck MA. Failed back surgery syndrome: current perspectives. *J Pain Res*. 2016;9:979–987. doi:10.2147/JPR.S92776.
8. Barolat G, Knobler RL, Lublin FD. Trigeminal neuralgia in a patient with multiple sclerosis treated with high cervical spinal cord stimulation. Case report. *Appl Neurophysiol*. 1988;51(6):333-337.
9. Bicket MC, Dunn RY, Ahmed SU. High-frequency spinal cord stimulation for chronic pain: Pre-clinical overview and systematic review of controlled trials. *Pain Med*. 2016;17(12):2326-2336.
10. Blonde L, Umpierrez GE, Reddy SS, et al. American Association of Clinical Endocrinology Clinical Practice Guideline: Developing a Diabetes Mellitus Comprehensive Care Plan—2022 Update. *Endocr Pract*. 2022;28(10). doi:10.1016/j.eprac.2022.08.002.
11. Boden WE, O'rourke RA, Teo KK, et al. Impact of Optimal Medical Therapy With or Without Percutaneous Coronary Intervention on Long-Term Cardiovascular End Points in Patients With Stable Coronary Artery Disease (from the COURAGE Trial). *Am J Cardiol*. 2009;104(1):1-4. doi:10.1016/j.amjcard.2009.02.059.
12. Brooker C, Russo M, Cousins MJ, et al. ECAP-Controlled Closed-Loop Spinal Cord Stimulation Efficacy and Opioid Reduction Over 24-Months: Final Results of the Prospective, Multicenter, Open-Label Avalon Study. *Pain Practice*. 2021;21(6):680-691. doi:10.1111/papr.13008.
13. Burns B, Watkins L, Goadsby PJ. Treatment of intractable chronic cluster headache by occipital nerve stimulation in 14 patients. *Neurology*. 2009;72:341-345.
14. Burns B, Watkins L, Goadsby PJ. Treatment of hemicrania continua by occipital nerve stimulation with a bion device: long-term follow-up of a crossover study. *Lancet Neurol*. 2008;7:1001-1012.
15. Carter SA. The definition of critical limb ischemia of the lower limb and distal systolic pressures. *Br J Surg*. 1983;70(3):188-189.
16. Caylor J, Reddy R, Yin S, et al. Spinal cord stimulation in chronic pain: evidence and theory for mechanisms of action. *Bioelectron Med*. 2019;5(1). doi:10.1186/s42234-019-0023-1.
17. Chen J, Hesseltine A, Nashi S, et al. A Real-World Analysis of High-Frequency 10 kHz Spinal Cord Stimulation for the Treatment of Painful Diabetic Peripheral Neuropathy. *J Diabetes Sci Technol*. 2021:193229682110603. doi: 10.1177/19322968211060316.
18. Claey's L, Berg W, Jonas S. Spinal cord stimulation in the treatment of chronic critical limb ischemia. *Acta Neurochir Suppl*. 2007;97(Pt 1):259-265.
19. Conte M, Bradbury A, Kolh P, et al. Global vascular guidelines on the management of chronic limb-threatening ischemia. *J Vasc Surg*. 2019;69(6):3S-125S.e40. doi:10.1016/j.jvs.2019.02.016.
20. Cranley J. Ischemic Rest Pain. *Arch Surg*. 1969;98(2):187. doi:10.1001/archsurg.1969.01340080079015.
21. Cruccu G, Aziz T, Garcia-Larrea L, et al. EFNS guidelines on neurostimulation therapy for neuropathic pain. *Eur J Neurol*. 2007;14(9):952-970.
22. Daousi C, Benbow SJ, MacFarlane IA. Electrical spinal cord stimulation in the long-term treatment of chronic painful diabetic neuropathy. *Diabet Med*. 2005;22:393-398.

23. De Andres J, Monsalve-Dolz V, Fabregat-Cid G et al. Prospective, Randomized Blind Effect-on-Outcome Study of Conventional vs High-Frequency Spinal Cord Stimulation in Patients with Pain and Disability Due to Failed Back Surgery Syndrome. *Pain Med.* 2017;18:2401-2421.
24. Deer T, Pope J, Hayek S, et al. Neurostimulation for the treatment of axial back pain: A review of mechanisms, techniques, outcomes, and future advances. *Neuromodulation.* 2014;17(suppl 2):52-68.
25. Deer T, Pope J, Hunter C et al. Safety Analysis of Dorsal Root Ganglion Stimulation in the Treatment of Chronic Pain. *Neuromodulation.* 2019;23(2):239-244. doi:10.1111/ner.12941.
26. Deer TR, Levy RM, Kramer J, et al. Dorsal root ganglion stimulation yielded higher treatment success rate for complex regional pain syndrome and causalgia at 3 and 12 months. *Pain.* 2017;158(4):669-681. doi:10.1097/j.pain.0000000000000814.
27. Deer TR, Pope JE, Lamer TJ, et al. The Neuromodulation Appropriateness Consensus Committee on Best Practices for Dorsal Root Ganglion Stimulation. *Neuromodulation.* 2018;22(1):1-35. doi:10.1111/ner.12845.
28. de Vos CC, Meier K, Zaalberg PB, et al. Spinal cord stimulation in patients with painful diabetic neuropathy: a multicenter randomized clinical trial. *Pain.* 2014;155:2426-2431. doi:10.1016/j.pain.2014.08.031.
29. Duarte R, Nevitt S, Maden M, et al. Spinal cord stimulation for the management of painful diabetic neuropathy: a systematic review and meta-analysis of individual patient and aggregate data. *Pain.* 2021;162(11):2635-2643. doi:10.1097/j.pain.0000000000002262.
30. Esposito M, Malayil R, Hanes M, Deer T. Unique Characteristics of the Dorsal Root Ganglion as a Target for Neuromodulation. *Pain Med.* 2019;20(S1):S23-S30. doi:10.1093/pm/pnz012.
31. Fontaine D, Christophe Sol J, Raoul S, Fabre N, Geraud G, Magne C et al. Treatment of refractory chronic cluster headache by chronic occipital nerve stimulation. *Cephalalgia.* 2011;31:1101-1105.
32. Forouzanfar T, Kemler M, Weber W, et al. Spinal cord stimulation in complex regional pain syndrome: cervical and lumbar devices are comparably effective. *Br J Anaesth.* 2004;92(3):348-353.
33. Gersbach P, Argitis V, Gardaz JP, et al. Late outcome of spinal cord stimulation for unreconstructable and limb-threatening lower limb ischemia. *Eur J Vasc Endovasc Surg.* 2007;33(6):717-724.
34. Grider JS, Manchikanti L, Carayannopoulos A, et al. Effectiveness of spinal cord stimulation in chronic spinal pain: A systemic review. *Pain Physician.* 2016;19(1):E33-E54.
35. Hara S, Andresen H, Solheim O, et al. Effect of Spinal Cord Burst Stimulation vs Placebo Stimulation on Disability in Patients With Chronic Radicular Pain After Lumbar Spine Surgery: A Randomized Clinical Trial. *JAMA.* 2022;328(15):1506–1514. doi:10.1001/jama.2022.18231.
36. Harden RN, McCabe CS, Goebel A, et al. Complex Regional Pain Syndrome: Practical Diagnostic and Treatment Guidelines. 5th ed. *Pain Med.* 2022;23(Suppl 1):S1-S53. doi:10.1093/pm/pnac046.
37. Harke H, Gretenkort P, Ladleif H, Rahman S. Spinal cord stimulation in sympathetically maintained complex regional pain syndrome type I with severe disability. A prospective clinical study. *Eur J Pain.* 2005;9(4):363-373.
38. Harvey AM. Classification of chronic pain—descriptions of chronic pain syndromes and definitions of pain terms. *Clin J Pain.* 1995;11(2):179.
39. Henson J, Varhabhatla N, Bebic Z, et al. Spinal Cord Stimulation for Painful Diabetic Peripheral Neuropathy: A Systematic Review. *Pain Ther.* 2021;10(2):895-908. doi:10.1007/s40122-021-00282-9.
40. Hoelzer BC, Edgar D, Lu SP, Taylor RS. Indirect Comparison of 10 kHz Spinal Cord Stimulation (SCS) versus Traditional Low-Frequency SCS for the Treatment of Painful Diabetic Neuropathy: A Systematic Review of Randomized Controlled Trials. *Biomedicine.* 2022;10(10):2630. doi:10.3390/biomedicine10102630.
41. Horan M, Jacobsen A, Scherer C, et al. Complications and Effects of Dorsal Root Ganglion Stimulation in the Treatment of Chronic Neuropathic Pain: A Nationwide Cohort Study in Denmark. *Neuromodulation.* 2020. doi:10.1111/ner.13171.
42. Huang KT, Martin J, Marky A, et al. A national survey of spinal cord stimulation trial-to-permanent conversion rates. *Neuromodulation.* 2015;18:133-139.
43. Huygen F, Kallewaard J, Nijhuis H, et al. Effectiveness and Safety of Dorsal Root Ganglion Stimulation for the Treatment of Chronic Pain: A Pooled Analysis. *Neuromodulation.* 2020;23(2):213-221. doi:10.1111/ner.
44. Jivegård L, Augustinsson L, Holm J, et al. Effects of spinal cord stimulation (SCS) in patients with inoperable severe lower limb ischaemia: a prospective randomised controlled study. *Eur J Vasc Endovasc Surg.* 1995;9(4):421-425.
45. Kapural L, Narouze S, Janicki TI, Mekhail N. Spinal cord stimulation is an effective treatment for the chronic intractable visceral pelvic pain. *Pain Med.* 2006;7(5):440-443.

46. Kapural L, Peterson E, Provenzano D, Staats P. Clinical Evidence for Spinal Cord Stimulation for Failed Back Surgery Syndrome (FBSS). *Spine*. 2017;42(14S): S61–S66.
47. Kapural L, Yu C, Doust MW, et al. Novel 10-kHz high-frequency therapy (HF10 Therapy) is superior to traditional low-frequency spinal cord stimulation for the treatment of chronic back and leg pain: The SENZA-RCT Randomized Controlled Trial. *Anesthesiology*. 2015;123(4):851-860.
48. Kapural L, Yu C, Doust MW, et al. Comparison of 10-kHz High-Frequency and Traditional Low-Frequency Spinal Cord Stimulation for the Treatment of Chronic Back and Leg Pain: 24-Month Results From a Multicenter, Randomized, Controlled Pivotal Trial. *Neurosurgery*. 2016;79(5):667-677. doi:10.1227/NEU.0000000000001418.
49. Kemler MA, Barendse GA, van Kleef M, van den Wildenberg FA, van Kleef M. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. *N Engl J Med*. 2000;343(9):618-624. doi:10.1056/NEJM200008313430904.
50. Kemler MA, de Vet HC, Barendse GA, van den Wildenberg FA, van Kleef M. Effect of spinal cord stimulation for chronic complex regional pain syndrome Type I: five-year final follow-up of patients in a randomized controlled trial. *J Neurosurg*. 2008;108(2):292-298. doi:10.3171/JNS/2008/108/2/0292.
51. Kemler MA, de Vet HC, Barendse GA, van den Wildenberg FA, van Kleef M. Spinal cord stimulation for chronic reflex sympathetic dystrophy – five year follow-up. *N Engl J Med*. 2006;354(22):2394-2396. doi:10.1056/NEJMc055504.
52. Kemler MA, de Vet HC, Barendse GA, van den Wildenberg FA, van Kleef M. The effect of spinal cord stimulation in patients with chronic reflex sympathetic dystrophy: two years' follow-up of the randomized controlled trial. *Ann Neurol*. 2004;55(1):13-18. doi:10.1002/ana.10996.
53. Kemler MA, Furnée CA. Economic evaluation of spinal cord stimulation for chronic reflex sympathetic dystrophy. *Neurology*. 2002;59(8):1203-1209. doi:10.1212/01.wnl.0000028686.74056.e3.
54. Klomp H, Spincemaille G, Steyerberg E, et al. Spinal-cord stimulation in critical limb ischaemia: a randomised trial. ESES Study Group. *Lancet*. 1999;353(9158):1040-1044.
55. Kumar K, Hunter G, Demeria D. Spinal cord stimulation in treatment of chronic benign pain: challenges in treatment planning and present status, a 22-year experience. *Neurosurgery*. 2006;58(3):481-496.
56. Kumar K, Taylor RS, Jacques L, et al. Spinal cord stimulation versus conventional medical management for neuropathic pain: A multicenter randomized controlled trial in patients with failed back surgery syndrome. *Pain*. 2007;132(1-2):179-188.
57. Lam CM, Monroe BR. Successful treatment of central pain and spasticity in patient with multiple sclerosis with dorsal column, paresthesia-free spinal cord stimulator: A case report. *AA Pract*. 2019;12(9):308-312.
58. Lambrou G, Matharu M. Occipital nerve stimulation in primary headache syndromes. *Ther Adv Neurol Disord*. 2012;5(1):57-67.
59. Latif O, Nedeljkovic S, Stevenson L. Spinal cord stimulation for chronic intractable angina pectoris: a unified theory on its mechanism. *Clin Cardiol*. 2001;24:533-541.
60. Levy R, Mekhail N, Kramer J, et al. Therapy Habituation at 12 Months: Spinal Cord Stimulation versus Dorsal Root Ganglion Stimulation for Complex Regional Pain Syndrome Type I and II. *J Pain*. 2020;21(3-4):399-408. doi:10.1016/j.jpain.2019.08.005.
61. Lipton RB, Goadsby PJ, Cady RK, et al. PRISM study: occipital nerve stimulation for treatment-refractory migraine. *Cephalalgia*. 2009;29:30.
62. Magis D, Allena M, Bolla M, De Pasqua V, Remacle JM, Schoenen J. Occipital nerve stimulation for drug-resistant chronic cluster headache: a prospective pilot study. *Lancet Neurol*. 2007;6:314-321.
63. Magis D, Gerardy PY, Remacle JM, Schoenen J. Sustained effectiveness of occipital nerve stimulation in drug-resistant chronic cluster headache. *Headache*. 2011;51:1191-1201.
64. Magis D, Schoenen J. Advances and challenges in neurostimulation for headaches. *Lancet Neurol*. 2012;11:708-719.
65. Mannheimer C, Eliasson T, Augustinsson LE, et al. Electrical stimulation versus coronary artery bypass surgery in severe angina pectoris: the ESBY study. *Circulation*. 1998;97:1157-1163.
66. Marin JC, Goadsby P. Response of SUNCT (short-lasting unilateral neuralgiform headaches with conjunctival injection and tearing), SUNA (short-lasting unilateral neuralgiform headaches with autonomic symptoms) and primary stabbing headaches to occipital nerve stimulation (ONS). *Neurology*. 2010;74:P04.006.
67. McGreevy K, Williams K. Contemporary insights into painful diabetic neuropathy and treatment with spinal cord stimulation. *Curr Pain Headache Rep*. 2012;16:43-49.

68. Mekhail N, Levy RM, Deer TR, et al; On behalf of the Evoke Study Group*. Long-term safety and efficacy of closed-loop spinal cord stimulation to treat chronic back and leg pain (Evoke): a double-blind, randomised, controlled trial. *Lancet Neurol.* 2020;19:123-134. doi:10.1016/S1474-4422(19)30414-4.
69. Mekhail NA, Mathews M, Nageeb F, et al. Retrospective review of 707 cases of spinal cord stimulation: indications and complications. *Pain Pract.* 2011;11:148-153.
70. Mekhail N, Visnjevac O, Azer G, Mehanny D, Agrawal P, Foorsov V. Spinal Cord Stimulation 50 Years Later. *Reg Anesth Pain Med.* 2018;43(4):391-406. doi:10.1097/aap.0000000000000744.
71. Mesaroli G, Hundert A, Birnie K, Campbell F, Stinson J. Screening and diagnostic tools for complex regional pain syndrome: a systematic review. *Pain.* 2020;162(5):1295-1304. doi:10.1097/j.pain.0000000000002146.
72. Misra S, Shishehbor M, Takahashi E, et al. Perfusion Assessment in Critical Limb Ischemia: Principles for Understanding and the Development of Evidence and Evaluation of Devices: A Scientific Statement from the American Heart Association. *Circulation.* 2019;140(12). doi:10.1161/cir.0000000000000708.
73. National Institute for Health and Care Excellence (NICE). Spinal Cord Stimulation for *chronic pain of neuropathic or ischaemic origin*. Technology appraisal guidance (TAP) 159. 2008 Oct. <https://www.nice.org.uk/guidance/ta159>.
74. North R, Calkins S, Campbell D, et al. Automated, patient-interactive, spinal cord stimulator adjustment: a randomized controlled trial. *Neurosurgery.* 2003;52(3):572-580.
75. North RB, Campbell JN, James CS, et al. Failed back surgery syndrome: Five year followup in 102 patients undergoing repeated operation. *Neurosurgery.* 1991;28;28: 685-91.
76. North R, Kidd D, Farrokhi F, Piantadosi S. Spinal cord stimulation versus repeated lumbosacral spine surgery for chronic pain: a randomized, controlled trial. *Neurosurgery.* 2005;56(1):98-106.
77. North R, Kidd D, Lee M, et al. A prospective, randomized study of spinal cord stimulation versus reoperation for failed back surgery syndrome: initial results. *Stereotact Funct Neurosurg.* 1994;62:267-272.
78. North R, Kidd H, Piantadosi S. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a prospective, randomized study design. *Acta Neurochir.* 1995;64(Suppl):106-108.
79. North R, Kidd D, Shipley J, Taylor R. Spinal cord stimulation versus reoperation for failed back surgery syndrome: a cost effectiveness and cost utility analysis based on a randomized, controlled trial. *Neurosurgery.* 2007;61(2):361-368.
80. Novo S, Coppola G, Milio G. Critical Limb Ischemia: Definition and Natural History. *Curr Drug Targets Cardiovasc Haematol Disord.* 2004;4(3):219-225. doi:10.2174/1568006043335989.
81. Pedrini L, Magnoni F. Spinal cord stimulation for lower limb ischemic pain treatment. *Interact Cardiovasc Thorac Surg.* 2007;6(4):495-500. doi:10.1510/icvts.2006.150185.
82. Perruchoud C, Eldabe S, Batterham AM, et al. Analgesic efficacy of high-frequency spinal cord stimulation: A randomized double-blind placebo controlled study. *Neuromodulation.* 2013;16(4):363-369.
83. Petersen E, Stauss T, Scowcroft J, et al. Durability of High-Frequency 10-kHz Spinal Cord Stimulation for Patients With Painful Diabetic Neuropathy Refractory to Conventional Treatments: 12-Month Results From a Randomized Controlled Trial. *Diabetes Care.* 2021;45(1):e3-e6. doi:10.2337/dc21-1813.
84. Pop-Busui R, Ang L, Boulton A, et al. Diagnosis and Treatment of Painful Diabetic Peripheral Neuropathy. *ADA Clinical Compendia.* 2022;2022(1):1-32. doi:10.2337/db2022-01.
85. Rapcan R, Mlaka J, Venglarcik M, et al. High-frequency-spinal cord stimulation. *Bratisl Lek Listy.* 2015;116(6):354-356.
86. Reed K. Peripheral neuromodulation and headaches: history, clinical approach, and considerations on underlying mechanisms. *Curr pain Headache Rep.* 2013;17:305.
87. Reed KL, Black SB, Banta CJ 2nd, Will KR. Combined occipital and supraorbital neurostimulation for the treatment of chronic migraine headaches: initial experience. *Cephalalgia.* 2010;30:260-271.
88. Russo M, Van Buyten JP. 10-kHz high-frequency SCS therapy: A clinical summary. *Pain Med.* 2015;16(5):934-942.
89. Saper JR, Dodick DW, Silberstein SD, McCarville S, Sun M, Goadsby PJ. Occipital nerve stimulation for the treatment of intractable chronic migraine headache: ONSTIM feasibility study. *Cephalalgia.* 2011;31:271-285.
90. Schwedt TJ, Dodick DW, Trentman TL, Zimmerman RS. Response to occipital nerve block is not useful in predicting efficacy of occipital nerve stimulation. *Cephalalgia.* 2007;27:271-274.
91. Shanahan P, Watkins L, Matharu M. Treatment of medically intractable short- lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT) and short-lasting unilateral neuralgiform

- headache attacks with autonomic symptoms (SUNA) with occipital nerve stimulation (ONS) in 6 patients. *Cephalalgia*. 2009;29:150.
92. Silberstein S, Dodick D, Saper J, et al. Safety and efficacy of peripheral nerve stimulation of the occipital nerves for the management of chronic migraine: Results from a randomized, multicenter, double-blinded, controlled study. *Cephalalgia*. 2012;32(16):1165-1179. doi:10.1177/0333102412462642.
 93. Sitzman B, Provenzano D. Best Practices in Spinal Cord Stimulation. *Spine*. 2017;42;14S,S67–S71.
 94. Sivanesan E, Bicket M, Cohen S. Retrospective analysis of complications associated with dorsal root ganglion stimulation for pain relief in the FDA MAUDE database. *Reg Anesth Pain Med*. 2019;44(1):100-106. doi:10.1136/rapm-2018-000007.
 95. Sivanesan E, Cohen S. Response to “Safety Analysis of Dorsal Root Ganglion Stimulation in the Treatment of Chronic Pain”. *Neuromodulation*. 2019;22(4):503-504. doi:10.1111/ner.12971.
 96. Skaribas I, Calvillo O, Delikanaki, Skaribas E. Occipital peripheral nerve stimulation in the management of chronic intractable occipital neuralgia in a patient with neurofibromatosis type 1: a case report. *J Med Case Rep*. 2011;5:174.
 97. Slangen R, Pluijms WA, Faber CG, Dirksen CD, Kessels AG, van Kleef M. Sustained effect of spinal cord stimulation on pain and quality of life in painful diabetic peripheral neuropathy. *Br J Anaesth*. 2013;111:1030-1031.
 98. Slangen R, Schaper NC, Faber CG, et al. Spinal Cord Stimulation and Pain Relief in Painful Diabetic Peripheral Neuropathy: A Prospective Two-Center Randomized Controlled Trial. *Diabetes Care*. 2014;37(11):3016-3024. doi:10.2337/dc14-0684.
 99. Stancák A, Kozák J, Vrba I, et al. Functional magnetic resonance imaging of cerebral activation during spinal cord stimulation in failed back surgery syndrome patients. *Eur J Pain*. 2008;12(2):137-148.
 100. Stanton-Hicks M, Burton A, Bruehl S, et al. An updated interdisciplinary clinical pathway for CRPS: report of an expert panel. *Pain Pract*. 2005;84(3):S4-S16.
 101. Sundaraj S, Johnstone C, Noore F, et al. Spinal cord stimulation: a seven-year audit. *J Clin Neurosci*. 2005;12(3):264-270.
 102. Taylor R, Buyten J, Buchser E. Spinal cord stimulation for complex regional pain syndrome: A systematic review of the clinical and cost-effectiveness literature and assessment of prognostic factors. *Eur J Pain*. 2006;10(2):91-101.
 103. Taylor RS, Desai MJ, Rigoard P, Taylor RJ. Predictors of pain relief following spinal cord stimulation in chronic back and leg pain and failed back surgery syndrome: A systemic review and meta-regression analysis. *Pain Pract*. 2014;14(6):489-505.
 104. Taylor R, Taylor R. Spinal cord stimulation for failed back surgery syndrome: a decision-analytic model and cost-effectiveness analysis. *Int J Technol Assess Health Care*. 2005;21(3):351-358.
 105. TenVaarwerk I, Jessurun G, DeJongste M, et al. Clinical outcome of patients treated with spinal cord stimulation for therapeutically refractory angina pectoris. The Working Group on Neurocardiology. *Heart*. 1999;82(1):82-88.
 106. Tesfaye S, Watt J, Benbow S, et al. Electrical spinal-cord stimulation for painful diabetic peripheral neuropathy. *Lancet*. 1996;348:1698-1701.
 107. Tiede J, Brown L, Gekht G, et al. Novel spinal cord stimulation parameters in patients with predominant back pain. *Neuromodulation*. 2013;16(4):370-375.
 108. Traeger AC, Gilbert SE, Harris IA, Maher CG. Spinal cord stimulation for low back pain. *Cochrane Database Syst Rev*. 2023;2023(3). doi:10.1002/14651858.cd014789.pub2.
 109. Ubbink D, Vermeulen H. Spinal cord stimulation for critical leg ischemia: a review of effectiveness and optimal patient selection. *J Pain Symptom Manage*. 2006;31:S30-S35.
 110. Ubbink D, Vermeulen H. Spinal cord stimulation for non-reconstructable chronic critical leg ischaemia. *Cochrane Database Syst Rev*. 2005;CD004001.
 111. van Beek M, Geurts J, Slangen R, et al. Severity of Neuropathy Is Associated with Long-term Spinal Cord Stimulation Outcome in Painful Diabetic Peripheral Neuropathy: Five-Year Follow-up of a Prospective Two-Center Clinical Trial. *Diabetes Care*. 2017;41(1):32-38. doi:10.2337/dc17-0983.
 112. van Beek M, Slangen R, Schaper NC, et al. Sustained treatment effect of spinal cord stimulation in painful diabetic peripheral neuropathy: 24-month follow-up of a prospective two-center randomized controlled trial. *Diabetes Care*. 2015;38(9):e132-e134. doi:10.2337/dc15-0740.

113. van Buyten JP, Al-Kaisy A, Smet I, Palmisani S, Smith T. High-frequency spinal cord stimulation for the treatment of chronic back pain patients: Results of a prospective multicenter European clinical study. *Neuromodulation*. 2013;16(1):59-65.
114. Weiner RL, Reed KL. Peripheral neurostimulation for control of intractable occipital neuralgia. *Neuromodulation*. 1999;2(3):217-222. doi:10.1046/j.1525-1403.1999.00217.x.
115. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/ AHA/ AAPA/ ABC/ ACPM/ AGS/ APhA/ ASH/ ASPC/ NMA/ PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2018;71:e127–248. doi:10.1016/j.jacc.2017.11.00.
116. Yeung AM, Huang J, Nguyen KT, et al. Spinal Cord Stimulation for Painful Diabetic Neuropathy. *J Diabetes Sci Technol*. 2024;18(1):168-192. doi:10.1177/19322968221133795.