

CIGNA MEDICAL COVERAGE POLICIES

Sleep Disordered Breathing Diagnosis and Treatment Guidelines

Effective Date: June 1, 2025



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by EviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

These guidelines include procedures EviCore does not review for Cigna. Please refer to the **Cigna CPT code list** for the current list of high-tech imaging procedures that EviCore reviews for Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five-digit codes, nomenclature and other data are copyright 2025 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Table of Contents

Guideline	Page
General Information.....	3
Abbreviations for Sleep Guidelines.....	4
Preface to the Sleep Guidelines.....	6
General Guidelines (SL-1.0).....	7
Practice Notes (SL-1.1).....	17
Sleep Diagnostics.....	19
Sleep Diagnostics-Coding (SL-1).....	20
Initial Sleep Diagnostic and Treatment Testing (SL-2.1).....	24
PSG and Multiple Sleep Latency Testing (excessive sleepiness)(SL-2.3).....	31
Maintenance of wakefulness testing (MWT)-indications and criteria (SL-2.4).....	35
Repeat sleep testing - home or attended sleep studies (SL-2.6).....	38
Diagnostic Testing pre- and post- hypoglossal nerve stimulator (HGNS) implantation (SL-2.8).....	43
Diagnostic Testing pre- and post transvenous phrenic nerve stimulator (TPNS) implantation (SL-2.9).....	48
Treatment of Sleep-related Breathing Disorders.....	54
Treatment - General information.....	55
Positive airway pressure devices (SL-4.2).....	60
Oral Appliances for the Treatment of Obstructive Sleep Apnea (SL-9).....	72
Pediatric Guidelines.....	75
Pediatric Sleep Guidelines (SL-3).....	76
Program Exclusions.....	81
Sleep Apnea Treatment Program Exclusions (SL-5).....	82
References.....	84
General References.....	85
Pediatric References.....	95
Questionnaires (SL-8).....	97

General Information

Guideline	Page
Abbreviations for Sleep Guidelines.....	4
Preface to the Sleep Guidelines.....	6
General Guidelines (SL-1.0).....	7
Practice Notes (SL-1.1).....	17

Abbreviations for Sleep Guidelines

SL.GG.104.A

v1.0.2025

Abbreviation	Full name
AASM	American Academy of Sleep Medicine
ABG	Arterial Blood Gas
AHI	Apnea-Hypopnea Index
AOSATF of AASM	Adult Obstructive Sleep Apnea Task Force of the American Academy of Sleep Medicine
APAP	Auto-adjusting positive airway pressure
ASV	Adaptive servo-ventilation
BMI	Body mass index (body weight divided by the square of the height)
BPAP	Bi-level positive airway pressure
CPAP	Continuous positive airway pressure
CSA	Central Sleep Apnea
EDS	Excessive daytime sleepiness
EtPCO ₂	End-tidal PCO ₂
HCPCS	Healthcare Common Procedural Coding System <ul style="list-style-type: none"> (Level II alphanumeric codes used to report services not included in CPT®)
HSAT	Home sleep apnea testing
IDTF	Independent Diagnostic Testing Facilities
JCAHO	Joint Commission on Accreditation of Healthcare Organizations
MSLT	Multiple Sleep Latency Test

Abbreviation	Full name
MWT	Maintenance of Wakefulness Test
OA	Oral appliance
OHS	Obesity Hypoventilation Syndrome
OSA	Obstructive sleep apnea
PAP	Positive airway pressure
PM	Portable monitoring (in home sleep studies)
PSG	Polysomnography
RDI	Respiratory disturbance index: (respiratory effort related arousals + apneas + hypopneas/total sleep time)
RERAs	Respiratory effort related arousals
Screening Tools for Sleep Disorders	Epworth Sleepiness Scale, Berlin Questionnaire (for sleep apnea), STOP-BANG questionnaire, Insomnia Severity Index
TcPCO ₂	Transcutaneous PCO ₂
VBG	Venous Blood Gas

Preface to the Sleep Guidelines

SL.GG.100.A

v1.0.2025

Overutilization of testing

A review of the testing history, including **ALL** of the following, avoids unnecessary repeat testing:

- The results of initial studies to narrow the differential diagnosis should be obtained prior to performing further tests.
- The clinical history should include a potential indication such as a known or suspected sleep disorder. These potential indications are addressed in greater detail within the applicable guidelines.
- The results of the requested study should be expected to have an impact on individual management or treatment decisions.
- Criteria for repeat studies are addressed in the applicable guideline **Repeat Sleep Testing - (Home or Attended Sleep Studies)**
- Testing when the same or similar studies have already been conducted is not medically necessary, without clear rationale that fulfills guideline criteria.

General Guidelines (SL-1.0)

SL.GG.0010.A

v1.0.2025

Clinical Evaluation

A current and comprehensive clinical evaluation by the treating medical provider (either face-to-face or telehealth), including documentation of sleep symptoms and physical examination elements relevant to the requested service, is required before a sleep study can be considered medically necessary.

Note: The rendering site must be a qualified provider of service per health plan policy

The clinical evaluation should include **ALL** of the following:

- Comprehensive sleep history including treatment history (if applicable) and **ONE** of the following:
 - Sleep symptoms relevant to the suspected diagnosis **OR**
 - Sufficient history as to allow completion of a sleep questionnaire **OR**
 - Results of a sleep questionnaire or sleep questionnaire proxy
- Physical examination
- Relevant diagnostic studies done at any time in the past (such as previous sleep studies, overnight pulse oximetry, and cardiac and pulmonary testing)
 - The provider (or medical office staff on behalf of the provider) must document a reasonable attempt to obtain results of prior sleep testing prior to requesting repeat testing
 - If a copy of the prior sleep testing report is not available from the member, a reasonable attempt includes, but is not limited to, contacting the location and/or provider where the prior study was completed, or obtaining a prior result from the individual's previous or current DME company or equipment provider
- Relevant lab results (ABG or serum bicarbonate performed within one year)

Sleep-related signs and symptoms

- Sleep testing is not medically necessary for the evaluation of insomnia in the absence of symptoms for another sleep fragmenting disorder such as obstructive sleep apnea or periodic limb movement disorder.
- Sleep testing of asymptomatic individuals prior to bariatric surgery is not medically necessary.
- A description of sleep-related signs and symptoms that are relevant to the suspected diagnosis are required before sleep testing can be considered medically necessary.

For example, if sleep-related breathing disorders are suspected, **ONE OR MORE** of the following should be described:

- Unexplained daytime sleepiness or fatigue
- Nocturnal snoring, snorting or gasping
- Nocturnal witnessed apneas or the subjective sensation of awakening short of breath, coughing or choking
- Frequent awakenings during the night
- Prior diagnosis of OSA with a description of response to therapy
- Morning headaches
- Resistant or refractory hypertension
- Observed oxygen desaturations during sleep or during drug-induced sedation in a medical setting
- Presence of atrial fibrillation
- Presence of systolic congestive heart failure
- New/recent stroke or transient ischemic attack in the last 6 months
- Other sleep related symptoms that are suggestive of a sleep disorder and may be included are:
 - Increased sleep-related movements, sleep talking, displaying confused or erratic behavior during sleep
 - Dream enactment behavior
 - Difficulty initiating sleep
 - Non-restorative sleep
 - Limited attention, or memory loss

Documented physical examination

Documented physical examination should include:

- Cardiopulmonary evaluation if visit is conducted in person
- Level of obesity
- Neck circumference
- Oropharyngeal examination including Friedman score or Mallampati classification
- Findings may include macroglossia, tonsillar hypertrophy, nasal polyps, septal deviation, turbinate hypertrophy, elongated/enlarged uvula, narrow/high arched hard palate, retrognathia (recessed mandible) or micrognathia (small mandible)

Relevant diagnostic studies done in the past

- When an individual is new to a practice, and a record of the previous diagnosis of sleep apnea is needed in order to supply new equipment, the new practice must

make an effort to obtain the prior study results from the prior sleep medicine practice or the DME supplier, and this should be documented.

- The provider (or medical office staff on behalf of the provider) must document a reasonable attempt to obtain results of prior sleep testing prior to requesting repeat testing.
- A reasonable attempt includes but is not limited to, contacting the location and/or provider where the prior study was completed, or obtaining a prior result from the individual's previous or current DME company or equipment provider; and if the prior options do not produce results, asking the individual for a copy of prior test results.

Note:

HSAT and/or PSG must be ordered by a treating medical provider and interpreted by a board-certified sleep medicine physician or a provider that is overseen by a board-certified sleep medicine physician.

Sleep Questionnaires

Three sleep questionnaires are commonly used to quantify the level of sleepiness or probability of having OSA. These validated questionnaires include (but are not limited to):

- Epworth Sleepiness Scale
- Berlin Questionnaire
- STOP BANG Questionnaire

The results of these questionnaires help formulate an individual's likelihood of having sleep-related disease, so the questionnaires must be appropriate for the specific sleep issue in question (e.g. STOP-BANG and Berlin are most appropriate for OSA evaluation).

- To view these questionnaires and their interpretation in their entirety, see **Questionnaires (SL-8)**.

Results of a questionnaire appropriate for the specific sleep issue in question are required, or any **ONE** of the following conditions can serve as a proxy for the sleep questionnaire requirement:

- Witnessed apnea, gasping, or choking
- Documented repeated desaturations during sleep by pulse oximetry (performed medically)
- Previous diagnosis of OSA confirmed in record by prior testing
- History and physical elements are provided that would permit calculation of the STOP- BANG survey

Definitions: Sleep-related breathing disorders

For the purpose of this guideline, sleep-related breathing disorders are defined by a positive diagnosis by valid sleep testing with **ONE OR MORE** of the following sleep-related breathing disorders:

Obstructive sleep apnea (OSA) as measured by valid testing is defined as:

- The predominantly obstructive apnea-hypopnea index (AHI) or respiratory disturbance index (RDI) on a PSG, or the apnea-hypopnea index (AHI) or respiratory event index (REI) on a HSAT is ≥ 15 events per hour; **OR**
- The predominantly obstructive apnea-hypopnea index (AHI) or respiratory disturbance index (RDI) on a PSG, or the apnea-hypopnea index (AHI) or respiratory event index (REI) on a HSAT is ≥ 5 and < 15 events per hour with documentation of **ANY** of the following:
 - Symptoms of sleepiness, fatigue, insomnia, or other symptoms leading to impaired sleep-related quality of life
 - Report of awakening with breath holding, gasping, or choking
 - Bed partner or other observer reports habitual snoring, breathing interruptions, or both during sleep
 - Hypertension, a mood disorder, cognitive dysfunction, coronary artery disease, congestive heart failure, atrial fibrillation, type 2 diabetes mellitus, or stroke

Central sleep apnea (CSA) is defined as ALL of the following:

- Presence of **ONE OR MORE** of the following:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or non-restorative sleep
 - Awakening short of breath
 - Witnessed apneas
- Central apnea and/or central hypopnea index ≥ 5 per hour
- Central hypopnea/apneas are $>50\%$ of the total number of apneas and hypopneas

Central sleep apnea (CSA) with Cheyne-Stokes Breathing is defined as ALL of the following:

- Presence of **ONE OR MORE** of the following:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or non-restorative sleep
 - Awakening short of breath
 - Witnessed apneas

- Known atrial fibrillation/flutter, congestive heart failure, or a neurological disorder
- PSG must show **ALL** of the following:
 - Central apnea and/or central hypopnea index ≥ 5 per hour
 - Central hypopnea/apneas are $\geq 50\%$ of the total number of apneas and hypopneas
 - Pattern of breathing meets criteria for Cheyne-Stokes Breathing (episodes of ≥ 3 consecutive central apneas and/or central hypopneas separated by a crescendo and decrescendo change in breathing amplitude with a cycle length of ≥ 40 seconds)

Treatment Emergent Central Sleep Apnea is defined as ALL of the following:

- Presence of **ONE OR MORE** of the following thought to be attributable to the central events:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or nonrestorative sleep
 - Awakening short of breath
 - Witnessed apneas
- Diagnostic PSG demonstrates ≥ 5 predominantly obstructive respiratory events per hour of sleep (obstructive or mixed apneas, hypopneas, respiratory effort related arousals [RERAs]).
- PSG during use of positive airway pressure without a backup rate shows improvement of obstructive events and emergence or persistence of central apneas/hypopneas with **BOTH** of the following:
 - Central apnea and/or hypopnea index ≥ 5 per hour
 - Central hypopnea/apneas are $\geq 50\%$ of the total number of apneas and hypopneas

Sleep-related hypoventilation is defined as when EITHER of the following occur during sleep:

- Increase in arterial PCO_2 , transcutaneous PCO_2 , or end-tidal PCO_2 to a value >55 mmHg for ≥ 10 minutes
- There is a ≥ 10 mmHg increase in arterial PCO_2 , transcutaneous PCO_2 , or end-tidal PCO_2 during sleep (compared to awake supine value) to a value >50 mmHg for ≥ 10 minutes

Site of care for sleep testing requests

Home versus facility

The guidelines herein provide evidence-based guidance for appropriate site of care for diagnostic sleep testing requests (facility sleep testing versus home sleep apnea testing) and positive airway pressure titration requests (facility titration study versus home automatic positive airway pressure use) based on co-morbid conditions and clinical characteristics. These are also provided in **In-lab PSG indications**.

Facility testing can be considered medically necessary due to **ONE** of the following reasons:

- Suspected obstructive sleep apnea with low pretest probability of moderate-to-severe OSA defined by **EITHER** of the following:
 - Lack of reported symptoms of excessive daytime sleepiness **AND AT LEAST TWO** of the following three criteria:
 - habitual loud snoring, **OR**
 - witnessed apnea or gasping or choking, **OR**
 - diagnosed hypertension
 - Validated questionnaire (STOP BANG or Berlin) indicating low risk
 - See **Questionnaires (SL-8)**
- Individual does not have the mobility, dexterity or cognitive ability to use home sleep apnea testing equipment safely at home and the ability to follow instructions.
- HSAT has been attempted and is negative, or technically inadequate (report submitted for review).
 - Technically inadequate HSAT recordings lack a minimum of 4 hours of oximetry and flow data.
- At least **ONE** of the following suspected or known co-morbid diagnoses or clinical scenarios is documented:
 - **Obesity with ONE of the following:**
 - BMI ≥ 45
 - BMI ≥ 30 kg/m² plus **EITHER** awake arterial blood gas (ABG), end-tidal PCO₂ (EtPCO₂), or transcutaneous PCO₂ (TcPCO₂) with PCO₂ ≥ 45 **OR** venous blood gas (VBG) showing a PCO₂ ≥ 50 mmHg
 - BMI ≥ 30 kg/m² plus serum bicarbonate ≥ 27 with stated concern for obesity hypoventilation syndrome (OHS)

- **Moderate to severe pulmonary disease (for example: COPD, asthma) as demonstrated by ONE OR MORE of the following:**
 - Nocturnal oxygen use
 - Documented arterial blood gases showing $PO_2 < 60$ or $PCO_2 \geq 45$
 - Documented pulmonary function tests demonstrating moderate to severe obstruction with forced expiratory volume in one second (FEV1) $\leq 69\%$ of predicted
- **Documented neurological disease**
 - **ANY** of the following:
 - The neurological disease precludes the individual's ability to perform home sleep apnea testing due to physical limitation (e.g. documented stroke with residual weakness/lack of coordination that would prevent performance of home sleep apnea testing) **OR**
 - There is stated concern for central sleep apnea or hypoventilation (e.g., neuromuscular disease such as myotonic dystrophy or ALS) due to known risk factor(s) described in clinical notes

Note:

See Background and Supporting Information for more details

- **Moderate to severe congestive heart failure**
 - With documented pulmonary congestion or known left ventricular ejection fraction $< 45\%$
- **Pulmonary HTN**
 - With documentation of a mean pulmonary artery pressure of > 20 mmHg on right heart catheterization

Note:

If right heart catheterization results are not available, echocardiography results can be provided documenting significant probability of pulmonary HTN based on a peak tricuspid regurgitation velocity of ≥ 2.9 m/s OR documented echocardiographic signs of pulmonary hypertension (Examples may include right atrial or right ventricle enlargement, interventricular septal flattening, enlarged pulmonary artery, dilated IVC, right ventricle outflow Doppler acceleration time > 105 msec and/or midsystolic notching, pulmonary regurgitation velocity > 2.2 m/sec)

- **Other critical illness that would prevent the individual from using the HSAT equipment**

- As documented in the individual's record
- **Chronic severe insomnia**
 - Documented by validated questionnaire such as Insomnia Severity Index (ISI) ≥ 22 .
 - See **Questionnaires (SL 8)** for details regarding ISI.
- **Chronic daily opioid use**
 - With stated concern for presence of central sleep apnea (typically daily high-potency opioids e.g. Methadone®, Suboxone®, Dilaudid®)
- **Titration study (in lab full night PAP titration, only)** where bi-level positive airway pressure (HCPCS E0470 or E0471) is specifically requested for **EITHER** of the following:
 - CPAP or APAP has already been tried and proven ineffective or not tolerated due to pressure discomfort for an individual with OSA **OR**
 - The individual has been diagnosed with **ONE** of the following:
 - Central sleep apnea with or without Cheyne-Stokes Breathing
 - Treatment-emergent central sleep apnea
 - Neuromuscular or restrictive thoracic disease
 - Severe COPD (arterial blood gas PaCO_2 is ≥ 52 mmHg done while awake and breathing the individual's prescribed FiO_2)
 - Obesity hypoventilation syndrome (OHS) defined as BMI ≥ 30 kg/m² plus awake arterial blood gas (ABG), end-tidal PCO_2 (EtPCO_2), or transcutaneous PCO_2 (TcPCO_2) with $\text{PCO}_2 \geq 45$ OR venous blood gas (VBG) showing a $\text{PCO}_2 \geq 50$ mmHg
 - Other Hypoventilation (defined awake arterial blood gas, end-tidal PCO_2 (EtPCO_2), or transcutaneous PCO_2 with $\text{PCO}_2 \geq 45$ OR venous blood gas showing a $\text{PCO}_2 \geq 50$ mmHg)
 - Sleep-related hypoventilation as defined in **General Guidelines (SL-1.0)**.
- **Sleep related hypoxemia (in lab full night PAP titration, only)**
 - Sustained oxygen desaturation independent of respiratory events on prior facility-based study or during prior home sleep apnea testing with documentation on the sleep study report of one or more periods of sustained oxygen saturation less than or equal to 88% lasting a minimum of 5 total minutes without significant apneas or hypopneas.
- **Documented OSA with unsuccessful APAP attempt (in lab full night PAP titration, only)**
 - Therapy trial of at least 30 days duration and **EITHER**:
 - APAP machine download with AHI ≥ 5 /hr with ongoing symptoms of OSA

- APAP use (≥ 4 hours per night on 70% of nights) with continued symptoms
- **Titration study (in lab full night PAP titration, only) is medically necessary with previously documented Central Sleep Apnea (CSA) defined as ALL of the following:**
 - Presence of **ONE OR MORE** of the following:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or non-restorative sleep
 - Awakening short of breath
 - Witnessed apneas
 - Central apnea and/or central hypopnea index ≥ 5 per hour
 - Central hypopnea/apneas are $>50\%$ of the total number of apneas and hypopneas
- **Titration study (in lab full night PAP titration, only) is medically necessary with previously documented Treatment Emergent Central Sleep Apnea defined as BOTH of the following:**
 - Diagnostic PSG demonstrates ≥ 5 respiratory events per hour of sleep
 - PSG during use of positive airway pressure shows improvement of obstructive events and emergence or persistence of central apneas/hypopneas with **BOTH** of the following:
 - Central apnea and/or hypopnea index ≥ 5 per hour
 - Central hypopnea/apneas are $\geq 50\%$ of the total number of apneas and hypopneas
- **Complicated parasomnias** (potentially injurious or when there is stated concern for nocturnal seizure activity).
 - A complete description, assessment, and discussion of signs and symptomatology must be included.
 - Polysomnography is not medically necessary for typical disorders of arousal (e.g. sleep terrors, confusional arousals, sleep walking, sleep-related eating disorder), nightmares, sleep talking, or bruxism in the absence of concern for safety, sleep disordered breathing, or nocturnal seizures.
- **Rapid Eye Movement (REM) Behavior Disorder (RBD):** Characterized by the acting out of dreams that are vivid, intense, and violent.
 - Consideration of this diagnosis requires a clear description of dream-enacting behaviors, which consist of sleep-related vocalization in addition to complex motor behaviors (such as punching or jumping from bed) correlating with sleep-related mentation.

- **Periodic limb movement disorder (PLMD)**, but not Restless Leg Syndrome (RLS)
 - Suspected PLMD is defined by periodic episodes of repetitive limb movements during sleep associated with insomnia or impaired daytime functioning (e.g. hypersomnia) not caused by another sleep disorder (such as untreated OSA, RLS, narcolepsy, REM sleep disorder), medical disorder, or mental disorder.
 - RLS is a subjective uncomfortable sensation experienced while awake in the evening that results in an urge to move the legs; it worsens during rest and improves during movement. RLS does not require a sleep study for diagnosis.
 - In the absence of comorbidities indicated in **General Guidelines (SL 1.0) (Site of Care)**, individuals who are undergoing initial diagnostic testing when there is a clinical suspicion for OSA should be evaluated for obstructive sleep apnea with an HSAT before a diagnosis of PLMD is considered.
- **Suspected narcolepsy or idiopathic hypersomnia** with MSLT

Note: See **PSG and Multiple Sleep Latency Testing (excessive sleepiness) Indications and Criteria (SL-2.3)**

Background and Supporting Information

Though CSA is an independent predictor of incident atrial fibrillation and is found in higher prevalence in established atrial fibrillation, CSA is much less common than OSA in individuals with atrial fibrillation. Therefore, atrial fibrillation on its own is not an indication for an in-lab PSG.

Sleep apnea is prevalent after strokes and transient ischemic attacks (TIA's) with lower sleepiness and BMI as compared to the non-stroke/TIA population. The predominant sleep apnea type is obstructive. Dominant central apneas occur in much lower frequency (6% per a cross-sectional survey). Type, severity, and location of stroke do not correlate clearly with central apnea severity. Severity of both OSA and CSA can reduce during the first six months post stroke/TIA. HSAT's have been shown to be feasible and accurate in stroke patients during in-hospital rehabilitation.

Per the International Classification of Sleep Disorders, PLMD cannot be diagnosed in the context of RLS, narcolepsy, untreated obstructive sleep apnea or REM sleep behavior disorder. The presence of periodic limb movements of sleep (PLMS) on sleep testing does not equate to PLMD. PLMS is common, but PLMD is rare in adults. Restless Legs Syndrome (Willis-Ekbom Disease) is a clinical diagnosis characterized by uncomfortable sensations in the legs that begin or worsen during rest, occur predominantly in the evening, and are partially or totally relieved by movement.

Preoperative sleep testing prior to bariatric surgery testing is based on these guidelines.

Practice Notes (SL-1.1)

SL.GG.101.A

v1.0.2025

Occupational Health Examinations

The guidelines herein are also applicable to cases involving medical qualifying examinations performed for occupational health purposes.

Specific government or regulatory body occupational health requirements can be considered when the regulatory rules are provided.

Technical and policy requirements of PSG

The parameters, settings, filters, technical specifications, sleep stage scoring and event scoring should be done in accordance with the AASM Manual for the Scoring of Sleep and Associated Events.

Home Sleep Apnea Testing equipment should at a minimum include the following features:

- FDA approval or clearance for the diagnosis of sleep disordered breathing
- Unique identifier for each unit
- Must meet minimum definition for CPT® 95800, 95801, or 95806
- Ability to record oximetry
- Ability to record a measure of heart rate
- Ability to display raw data for review, manual scoring or editing of automated scoring
- Ability to calculate a respiratory event index (REI) based on monitoring time (as a surrogate for AHI) **OR** based on estimated sleep time derived from actigraphy in the case of a peripheral arterial tonometry (PAT) device

HSAT can be appropriately performed by Joint Commission (JCAHO) and Medicare IDTF-approved facilities.

PSG is called Type I monitoring and involves at a minimum, measurement of the following parameters:

- Electroencephalogram (EEG)
- Electrooculogram (EOG)
- Chin electromyogram (EMG)
- Leg EMG
- Airflow signals

- Respiratory effort signals
- Oxygen saturation
- Body position
- Electrocardiogram (EKG)
- Synchronized PSG video
 - Facilities also typically record body position (with video) and snoring (via microphone).
- Calculations of the Apnea Hypopnea Index (AHI) and Respiratory Disturbance Index (RDI) are performed.

Scoring PSG:

- AHI is the number of apneas and hypopneas per hour of sleep
- RDI is the number of apneas, hypopneas, and respiratory effort related arousals (RERAs) per hour of sleep

Scoring HSAT:

- AHI (also called the REI) is the number of apneas and hypopneas per hour of estimated sleep
- The RDI cannot be used since RERAs cannot be scored on HSAT

Sleep Diagnostics

Guideline	Page
Sleep Diagnostics-Coding (SL-1).....	20
Initial Sleep Diagnostic and Treatment Testing (SL-2.1)	24
PSG and Multiple Sleep Latency Testing (excessive sleepiness)(SL-2.3).....	31
Maintenance of wakefulness testing (MWT)-indications and criteria (SL-2.4).....	35
Repeat sleep testing - home or attended sleep studies (SL-2.6).....	38
Diagnostic Testing pre- and post- hypoglossal nerve stimulator (HGNS) implantation (SL-2.8).....	43
Diagnostic Testing pre- and post transvenous phrenic nerve stimulator (TPNS) implantation (SL-2.9).....	48

Sleep Diagnostics-Coding (SL-1)

SL.ST.100.A

v1.0.2025

The inclusion of any code in the below tables does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

Home portable monitoring (PM) (home sleep testing)-coding (SL-1.4.1)

There are currently 3 levels (HCPCS G0398, G0399 and G0400) of home PMs, with varying number of monitored parameters. Each can be used with or without an attendant but are generally performed unattended in the individual's home.

Procedure codes for home sleep apnea testing (HSAT)

Home sleep studies	HCPCS	Channels
Home sleep study test (HSAT) with type II portable monitor, unattended	G0398	Minimum of 7 channels: EEG, EOG, EMG, ECG/heart rate, airflow, respiratory effort and oxygen saturation
Home sleep test (HSAT) with type III portable monitor, unattended	G0399	Minimum of 4 channels: 2 respiratory movement/ airflow, 1 ECG/heart rate, and 1 oxygen saturation
Home sleep test (HSAT) with type IV portable monitor, unattended	G0400	Minimum of 3 channels <ul style="list-style-type: none"> Device type (airflow or peripheral arterial tone) not specified in HCPCS description

PSG procedure codes for unattended sleep studies

Unattended sleep studies	CPT®	Channels
Sleep study, unattended, simultaneous recording	95800	Minimum of 4 channels: heart rate, oxygen saturation, respiratory analysis (e.g., by airflow or peripheral arterial tone), and sleep time.
Sleep study, unattended, simultaneous recording	95801	Minimum of 3 channels: minimum of heart rate, oxygen saturation, and respiratory analysis (e.g., by airflow or peripheral arterial tone)
Sleep study, unattended, simultaneous recording <ul style="list-style-type: none"> Do not report CPT® 95806 in conjunction with any of the following codes: CPT® 93041-93229, 93268-93272, or 95800-95801 	95806	Minimum of 4 channels: heart rate, oxygen saturation, respiratory airflow, and respiratory effort (e.g., thoracoabdominal movement)

Polysomnography (facility-based-PSG) - coding (SL-1.4.2)

PSG procedure codes for attended polysomnography and sleep studies

Attended polysomnography and sleep studies	CPT®
<p>Multiple sleep latency (MSLT) or Maintenance of Wakefulness Testing (MWT), recording, analysis and interpretation of physiological measurements of sleep during multiple trials to assess sleepiness.</p> <ul style="list-style-type: none"> Multiple sleep latency testing (MSLT) is performed prior to treatment when the requesting physician suspects narcolepsy. MSLT must be requested with a facility sleep study performed the night before the CPT® 95805 (CPT® 95810 or CPT® 95811). CPT® 95805 is used to report MSLT or MWT See Multiple sleep latency testing (MSLT) (SL-2.3) - See Maintenance of Wakefulness Testing (MWT)-Indications and Criteria (SL-2.4) 	95805
<p>Sleep study, simultaneous recording of ventilation, respiratory effort, ECG or heart rate, and oxygen saturation, attended by a technologist with PAP titration</p> <ul style="list-style-type: none"> May be considered experimental, investigational, OR unproven when 95807 OR 95807-52 is utilized to request a PAP-NAP. 	95807
<p>Polysomnography; (any age), sleep staging with 1-3 additional parameters of sleep, attended by a technologist</p>	95808
<p>Polysomnography; (age 6 years or older), sleep staging with 4 or more additional parameters of sleep, attended by a technologist.</p> <ul style="list-style-type: none"> CPT® 95810 is used to report in lab full night diagnostic sleep study <ul style="list-style-type: none"> CPT® 93152 is used to report interrogation and programming of implanted phrenic nerve stimulator system during polysomnography (CPT® 95810) 	95810

Attended polysomnography and sleep studies	CPT®
<p>Polysomnography; (age 6 years or older), sleep staging with 4 or more additional parameters of sleep, with initiation of continuous positive airway pressure therapy or Bi-level ventilation, attended by a technologist</p> <ul style="list-style-type: none"> • CPT® 95811 is used as EITHER an in lab split night study when both the diagnostic study and the subsequent positive airway pressure or bi-level ventilation are initiated during the same visit, OR as in lab full night PAP titration alone after CPT® 95810 or inability to complete split night sequence or as a re-titration of PAP therapy. 	95811

The following are **pediatric codes**

Attended polysomnography and sleep studies (pediatric codes)

Attended polysomnography and sleep studies	CPT
<p>Polysomnography, (younger than 6 years), sleep staging with 4 or more additional parameters of sleep, attended by a technologist.</p> <ul style="list-style-type: none"> • CPT® 95782 is used to report an in lab full night diagnostic sleep study 	95782
<p>Polysomnography, (younger than 6 years), sleep staging with 4 or more additional parameters of sleep, with initiation of continuous positive airway pressure therapy or bi-level ventilation, attended by a technologist.</p> <ul style="list-style-type: none"> • CPT® 95783 is used to report an in lab full night PAP titration 	95783

Initial Sleep Diagnostic and Treatment Testing (SL-2.1)

SL.ST.200.A

v1.0.2025

Note: For more information on technical policy requirements of HSAT and PSG as well as on scoring, see Practice Notes (SL-1.1)

Home sleep apnea testing (HSAT)

- HSAT is medically necessary when **BOTH** of the following criteria are met:
 - Increased risk of moderate-to-severe OSA as determined by **ONE** of the following:
 - Reported symptoms of excessive daytime sleepiness **AND AT LEAST TWO** of the following three criteria:
 - habitual loud snoring, **OR**
 - witnessed apnea or gasping or choking, **OR**
 - diagnosed hypertension
 - Validated questionnaire (STOP BANG or Berlin).
 - See **Questionnaires (SL-8)**.
 - HSAT can physically be performed and individual has the mobility, dexterity and cognitive ability to use the available equipment safely at home and the ability to follow instructions.
- HSAT can also be used in follow-up treatment results after **ANY** of the following:
 - Surgical treatment for moderate to severe OSA
 - OSA oral appliance trial
 - Other non-PAP supportive interventions (e.g. positional therapy)
- Individuals' who exhibit one of the co-morbid indications for attended sleep studies found in **General Guidelines (SL-1.0)**, In-Lab Polysomnography-Sleep Disordered Breathing, and In-Laboratory Polysomnography-Other Suspected Sleep Disorders can undergo facility testing or home sleep apnea testing (HSAT) if preferred by the treating provider.

In-laboratory polysomnography- sleep disordered breathing

PSG (in lab full night diagnostic sleep study **OR** in lab split night sleep study **OR** in lab full night PAP titration) is medically necessary when sleep survey or proxy symptom(s) lead to concern for sleep disordered breathing **AND ONE** of the following:

- Facility testing can be considered medically necessary rather than HSAT due to **ONE** of the following reasons:
 - Suspected obstructive sleep apnea with low pretest probability of moderate-to-severe OSA defined by **EITHER** of the following:
 - Lack of reported symptoms of excessive daytime sleepiness **AND AT LEAST TWO** of the following three criteria:
 - habitual loud snoring, **OR**
 - witnessed apnea or gasping or choking, **OR**
 - diagnosed hypertension
 - Validated questionnaire (STOP BANG or Berlin) indicating low risk.
 - See **Questionnaires (SL-8)**.
 - Individual does not have the mobility, dexterity or cognitive ability to use home sleep apnea testing equipment safely at home and the ability to follow instructions.
 - HSAT has been attempted and is negative, or technically inadequate (report submitted for review).
 - Technically inadequate HSAT recordings lack a minimum of 4 hours of oximetry and flow data.
- At least **ONE** of the following suspected or known co-morbid diagnoses or clinical scenarios is documented:

Obesity with ONE of the following:

- BMI ≥ 45
- BMI $\geq 30 \text{ kg/m}^2$ plus **EITHER** awake arterial blood gas (ABG), end-tidal PCO₂ (EtPCO₂), or transcutaneous PCO₂ (TcPCO₂) with PCO₂ ≥ 45 **OR** venous blood gas (VBG) showing a PCO₂ ≥ 50 mmHg
- BMI $\geq 30 \text{ kg/m}^2$ plus serum bicarbonate ≥ 27 with stated concern for obesity hypoventilation syndrome (OHS)

Moderate to severe pulmonary disease (for example: COPD, asthma) as demonstrated by ONE OR MORE of the following:

- Nocturnal oxygen use
- Documented arterial blood gases showing PO₂ < 60 or PCO₂ > 45

- Documented pulmonary function tests demonstrating moderate to severe obstruction with forced expiratory volume in one second (FEV1) $\leq 69\%$ of predicted

Documented neurological disease:**ANY** of the following:

- The neurological disease precludes the individual's ability to perform home sleep apnea testing due physical limitation (e.g. documented stroke with residual weakness/lack of coordination that would prevent performance of home sleep apnea testing) **OR**
- There is stated concern for central sleep apnea or hypoventilation (e.g. neuromuscular disease such as myotonic dystrophy or ALS) due to known risk factor(s) described in clinical notes

Note:

See **Background and Supporting Information (SL-1.0)** for more details

Moderate to severe congestive heart failure:

- With documented pulmonary congestion or known left ventricular ejection fraction $<45\%$

Pulmonary HTN:

- With documentation of a mean pulmonary artery pressure of >20 mmHg on right heart catheterization

Note:

If right heart catheterization results are not available, echocardiography results can be provided documenting significant probability of pulmonary HTN based on a peak tricuspid regurgitation velocity of ≥ 2.9 m/s OR documented echocardiographic signs of pulmonary hypertension (Examples may include right atrial or right ventricle enlargement, interventricular septal flattening, enlarged pulmonary artery, dilated IVC, right ventricle outflow Doppler acceleration time >105 msec and/or midsystolic notching, pulmonary regurgitation velocity >2.2 m/sec)

Other critical illness that would prevent the individual from using the HSAT equipment:

- As documented in the individual's record

Chronic severe insomnia:

- Documented by validated questionnaire (Insomnia Severity Index ≥ 22).
 - See **Questionnaires (SL 8)** for details regarding ISI.

Chronic daily opioid use:

- With stated concern for presence of central sleep apnea (typically daily high-potency opioids e.g. Methadone®, Suboxone®, Dilaudid®)

Titration Study

Titration study (in lab full night PAP titration, only) where bilevel positive airway pressure (HCPCS E0470 or E0471) are specifically requested for **EITHER** of the following:

- CPAP or APAP has already been tried and proven ineffective or not tolerated due to pressure discomfort for an individual with OSA **OR**
- The individual has been diagnosed with **ONE** of the following:
 - Central sleep apnea with or without Cheyne-Stokes Breathing
 - Treatment-emergent central sleep apnea
 - Neuromuscular or restrictive thoracic disease
 - Severe COPD (arterial blood gas PaCO₂ is ≥52 mmHg done while awake and breathing the individual's prescribed FiO₂)
 - Obesity hypoventilation syndrome (OHS) defined as BMI ≥30 kg/m₂ plus awake arterial blood gas (ABG), end-tidal PCO₂ (EtPCO₂), or transcutaneous PCO₂ (TcPCO₂) with PCO₂ ≥45 **OR** venous blood gas (VBG) showing a PCO₂ ≥50 mmHg
 - Other hypoventilation (defined awake arterial blood gas, end-tidal PCO₂ (ET PCO₂), or transcutaneous PCO₂ with PCO₂ ≥45 **OR** venous blood gas showing a PCO₂ ≥50 mmHg)
 - Sleep-related hypoventilation as defined in **General Guidelines (SL-1.0)**.

Sleep related hypoxemia (in lab full night PAP titration, only)

Sustained oxygen desaturation independent of respiratory events on prior facility-based study or during prior home sleep apnea testing with documentation on the sleep study report of one or more periods of sustained oxygen desaturation ≤88% lasting a minimum of 5 total minutes in the absence of apneas or hypopneas.

Documented OSA with unsuccessful APAP attempt (in lab full night PAP titration, only)

Therapy trial of at least 30 days duration and **EITHER** of the following:

- APAP machine download with AHI ≥5/hr with ongoing symptoms (of OSA)
- APAP use (≥4 hours per night on 70% of nights) with continued symptoms

Titration study (in lab full night PAP titration, only) is medically necessary with previously documented Central Sleep Apnea (CSA) defined as ALL of the following:

- Presence of **ONE OR MORE** of the following:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or non-restorative sleep
 - Awakening short of breath
 - Witnessed apneas
- Central apnea and/or central hypopnea index ≥ 5 per hour
- Central hypopnea/apneas are $>50\%$ of the total number of apneas and hypopneas

Titration study (in lab full night PAP titration, only) is medically necessary with previously documented Treatment Emergent Central Sleep Apnea defined as BOTH of the following:

- Diagnostic PSG demonstrates ≥ 5 respiratory events per hour of sleep
- PSG during use of positive airway pressure shows improvement of obstructive events and emergence or persistence of central apneas/hypopneas with **BOTH** of the following:
 - Central apnea and/or hypopnea index ≥ 5 per hour
 - Central hypopnea/apneas are $\geq 50\%$ of the total number of apneas and hypopneas

In-laboratory polysomnography - other suspected sleep disorders

In lab full night diagnostic sleep study **OR** in lab split night sleep study **OR** in lab full night PAP titration study is medically necessary for **ANY** of the following:

- **Complicated parasomnias** (potentially injurious or when there is stated concern for nocturnal seizure activity).
 - A complete description, assessment, and discussion of signs and symptomatology must be included.
 - Polysomnography is not medically necessary for typical disorders of arousal (e.g.: sleep terrors, confusional arousals, sleep walking, sleep-related eating disorder), nightmares, sleep talking, or bruxism in the absence of concern for safety, sleep disordered breathing, or nocturnal seizures.
- **Rapid Eye Movement (REM) Behavior Disorder (RBD)**: Characterized by the acting out of dreams that are vivid, intense, and violent.
 - Consideration of this diagnosis requires a clear description of dream-enacting behaviors, which consist of sleep-related vocalization in addition to complex motor

behaviors (such as punching or jumping from bed) correlating with sleep-related mentation.

- **Periodic limb movement disorder (PLMD)**, but not Restless Leg Syndrome (RLS)
 - Suspected PLMD is defined by periodic episodes of repetitive limb movements during sleep associated with insomnia or impaired daytime functioning (e.g. hypersomnia) not caused by another sleep disorder (such as untreated OSA, RLS, narcolepsy, REM sleep disorder), medical disorder, or mental disorder.
 - RLS is a subjective uncomfortable sensation experienced while awake in the evening that results in an urge to move the legs; it worsens during rest and improves during movement. RLS does not require a sleep study for diagnosis.
 - In the absence of comorbidities indicated in **General Guidelines (SL 1.0)** (site of care), individuals' who are undergoing initial diagnostic testing when there is a clinical suspicion of OSA should be evaluated for obstructive sleep apnea with an HSAT before a diagnosis of PLMD is considered.
- **Suspected narcolepsy or idiopathic hypersomnia** with MSLT

Note:

See PSG and Multiple Latency Testing (excessive sleepiness) Indications and Criteria (SL-2.3)

Split night study or two night study

Split night study

- Split night study (in lab split night sleep study) is a single-night PSG + PAP trial, and typically is medically necessary if **BOTH** of the following are met:
 - Apnea Hypopnea Index (AHI) is ≥ 15 /hr during ≥ 2 hours of recording time on the diagnostic PSG
 - ≥ 3 hours are available for PAP titration
- Split night study (in lab split night sleep study) can be achieved in the majority of cases in one night. This is the current recommended approach per the American Academy of Sleep Medicine (AASM) if the above criteria are met.

Two night study

- In some cases, a split night study cannot be completed because the above criteria is not met and sleep testing must be done in two nights.
 - The first night study was performed as a diagnostic study (in lab full night diagnostic study)

- Followed by second night PAP titration (in lab full night PAP titration)
- When PAP titration (in lab full night PAP titration) is subsequently requested after a completed PSG (full-night diagnostic of failed split-study) or HSAT meeting criteria for the diagnosis of obstructive sleep apnea (OSA) as defined in **General Guidelines (SL 1.0): Definitions**, the following information should be used to consider unattended APAP or attended CPAP titration:
 - Facility titration study (in lab full night PAP titration) is medically necessary when there is documentation of at least one of the suspected or known co-morbid diagnoses or clinical scenarios found in **General Guidelines (SL-1.0): Site of care** for sleep testing requests (home versus facility).
 - Immediate treatment with E0601 APAP, or if necessary, an E0601 APAP titration is medically necessary when there are no co-morbid diagnoses or clinical scenarios found in **General Guidelines (SL-1.0): Site of care** for sleep testing requests (home versus facility)
 - All new information from the PSG or HSAT including but not limited to frequency of central apneas, evidence of nocturnal hypoventilation independent of apnea/hypopnea, or abnormal sleep related behavior
 - Facility titration studies (in lab full night PAP titration) are also medically necessary in the following clinical scenarios:
 - Titrate non-invasive positive pressure ventilation (NIPPV) devices in individuals' with diurnal chronic alveolar hypoventilation (defined as hypercapnia, with awake arterial blood gas, end-tidal PCO₂ [ET PCO₂], or transcutaneous PCO₂ with PCO₂ >45 **OR** venous blood gas showing a PCO₂ ≥50 mmHg)
 - Titrate positive airway pressure or NIPPV for sleep-related hypoventilation (as defined in **General Guidelines (SL 1.0)**)

PSG and Multiple Sleep Latency Testing (excessive sleepiness)(SL-2.3)

SL.ST.203.A

v1.0.2025

MSLT - indications and criteria

ALL of the following must be met:

- In lab full night diagnostic sleep study followed by MSLT is being performed for suspected narcolepsy or idiopathic hypersomnia as evidenced by:
 - Excessive sleepiness (shown not due to other more common sleep disorders such as obstructive sleep apnea or insufficient sleep syndrome), **AND**
 - Recurrent daytime naps or lapses into sleep daily for at least 3 months
 - Additional symptoms may include:
 - Cataplexy- sudden loss of muscle tone occurring in association with intense emotion (laughing or crying), **OR**
 - Sleep paralysis, hypnagogic hallucinations, hypnopompic hallucinations, automatic behaviors, or disrupted major sleep episode
- Comprehensive sleep evaluation including performance of a relevant sleep questionnaire (ESS or Berlin) and consideration of alternative causes of sleepiness (e.g. medication/substance use or withdrawal, mental disorder, or medical disorder) **OR** comorbid sleep conditions (e.g. circadian rhythm sleep-wake disorder, insufficient sleep syndrome, or OSA)
 - The definition of Insufficient Sleep Syndrome requires that sleep time be shorter than expected for age on most days for at least 3 months. Minimum sleep duration for adults age ≥18 year is 7 hours, for teenagers age 13-17 years is 8 hours, and for school aged-children age 6-12 years is 9 hours.
- If there are symptoms or suspicion of OSA (or other types of sleep disordered breathing), diagnostic testing for these conditions must be performed first. See **General Guidelines (SL-1.0)** including section on **Site of care for sleep testing requests (home versus facility)**.
- If OSA (or other types of sleep disordered breathing) is present on diagnostic testing, therapy must be initiated prior to consideration of MSLT. If the individual is being treated with positive airway pressure (PAP) for obstructive sleep apnea, optimal efficacy and adherence has been achieved by demonstration of **ALL** of the following:
 - PAP download shows: AHI <5 on download

- Currently using PAP ≥ 4 hours per night on 70% of nights during a consecutive thirty (30) day period during the preceding 6 months.
- Therapy has resolved symptoms of increased upper airway resistance (i.e. eliminated snoring).
- If the individual is being treated with non-PAP therapy for obstructive sleep apnea, optimal efficacy and adherence has been achieved by demonstration of **ALL** of the following:
 - Self-reported adequate use
 - Efficacy confirmed on sleep testing where AHI < 5 /hour
 - Therapy has resolved symptoms of increased upper airway resistance (i.e., eliminated snoring)
- Is not requested to assess efficacy of PAP therapy for OSA (or other types of sleep disordered breathing).
- Multiple Sleep Latency Testing (MSLT) must immediately follow PSG (in lab full night diagnostic study).
 - It cannot follow a split night study or a PAP titration study in individual's for whom OSA (or other types of sleep disordered breathing) is not optimally treated based on ongoing symptoms (such as snoring and witnessed apneas despite compliant use of PAP) or lack of optimal efficacy and adherence (defined as AHI < 5 on PAP download [if available] and using PAP $\geq 70\%$ of the nights for an average of 4 hours or more per 24-hour period) based on downloaded data from the individual's PAP machine.

Note:

When the individual has known OSA (or other types of sleep disordered breathing) which is optimally treated with PAP therapy and has persistent excessive daytime sleepiness and symptoms concerning for narcolepsy or idiopathic hypersomnia, the preceding night's attended study, prior to the next-day MSLT, should be completed while the individual is using his/her PAP therapy at its optimal setting. Either CPT[®] 95810 or 95811 can be used to precede MSLT for individuals with known OSA controlled on PAP therapy.

- Reassessment of suspected narcolepsy or idiopathic hypersomnia with a repeat in lab full night diagnostic sleep study/MSLT is medically necessary if **EITHER** of the following:
 - Previous testing did not confirm the diagnosis but clinical suspicion is still present despite consideration of alternative causes of sleepiness (e.g. medication/substance use or withdrawal, mental disorder, or medical disorder) with adequate treatment of comorbid sleep conditions (e.g., circadian rhythm sleep-wake disorder, insufficient sleep syndrome, or OSA as described above)

- Due to a change in symptoms that might alter a previous diagnosis of Narcolepsy or Idiopathic Hypersomnia

Background and Supporting Information

The purpose of the PSG followed by MSLT is to measure an individual's physiological tendency to fall asleep in the absence of external alerting factors. It can be utilized to diagnose narcolepsy types 1 and 2 as well as idiopathic hypersomnia.

For individuals' with very mild OSA (AHI <10/hour, who are intolerant to therapy, consideration may be given to not use OSA therapy during the MSLT.

Per Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in adults: guidance from the American Academy of Sleep Medicine, *Journal of Clinical Sleep Medicine* 2021:

- The MSLT should be performed following an attended PSG which allows a minimum 7 hours of time in bed with at least 6 hours of sleep, with timing that corresponds with the individual's major sleep period. The test should not be performed after a night during which PAP pressures were adjusted (split-night or PAP titration study).
- "Individuals' on PAP/non-PAP therapies for sleep-disordered breathing should use them during the PSG and MSLT. The PAP settings and mask interface should match those used at home."

Per Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in children: guidance from the American Academy of Sleep Medicine, *Journal of Clinical Sleep Medicine* 2024:

- "Symptoms of narcolepsy can present differently in children and adolescents compared to adults. First, daytime sleepiness may be expressed as inattentive, hyperactive, emotionally labile, and/or impulsive behavior in children and adolescents. Second, children and adolescents with NT1 can present initially with cataplectic facies (a static form of cataplexy presenting with ptosis, jaw lowering, and tongue protrusion) before more classic emotionally triggered cataplexy develops."
- "MSLT may not be appropriate in children < 5 years of age due to lack of normative data in this age group. However, consideration may be given to performing MSLT in children < 5 years of age in special circumstances such as classic symptoms of narcolepsy with cataplexy, and/or inability to obtain CSF orexin testing. Additional consideration should be given when interpreting MSLT results in prepubertal children, because prepubertal children are less likely to fall asleep during the day than older adolescents. Habitual naps are common in younger children and should be taken into account when interpreting MSLT data. Furthermore, mean sleep latencies tend to decrease with advancing Tanner stages in pubescent children."

- The MSLT should be performed following an attended PSG which allows a minimum of 8 hours time in bed with at least 7 hours of sleep, ideally meeting age-based needs.
- "Any home PAP/non-PAP therapies for sleep-disordered breathing patients should be used during the PSG and considered for use during MSLT naps based on clinical judgment. The therapeutic modality, PAP settings, and/or mask interface should match those used at home."
- Moderate to severe OSA should be treated prior to PSG-MSLT evaluation.

Evidence Discussion

In light of the relatively uncommon nature of narcolepsy, as compared to more common sleep and medical disorders, guidelines are meticulously followed to exclude these common disorders before approval of testing for narcolepsy or idiopathic hypersomnia, treatment of which can be dangerous. To achieve a correct diagnosis of narcolepsy or idiopathic hypersomnia, the American Academy of Sleep Medicine (AASM) has published a protocol that requires the following:

- Adequate sleep documented by sleep diary or actigraphy for 2 weeks before testing
- Testing be performed when the individual is clinically stable and when treatment for existing sleep disorders are well-established and effective
- Medications that have alerting, sedating, and/or REM-sleep-modulating properties should be stopped at least 2 weeks before the Multiple Sleep Latency Test (MSLT)
- An overnight polysomnography (PSG) should be done on the night before the MSLT with optimal sleep apnea treatment (if applicable) for the purpose of documenting sufficient sleep, determining if a sleep-onset REM period (SOREMP) within 15 minutes of nighttime sleep onset occurred, and confirming the lack of any other sleep disorder that may be more common, such as sleep apnea. Split-night PSG should be avoided because it does not represent a full night of sleep uninterrupted by respiratory events and or PAP therapy adjustments

The test-retest reliability of the MSLT is low in narcolepsy type 2 and idiopathic hypersomnia (IH), with a confirmatory second MSLT in 18-47% of narcolepsy type 2 and 25% of idiopathic hypersomnia. Although a diagnosis of narcolepsy type 1 has a high rate of reproducibility on repeat MSLT, it can be missed on an initial MSLT and later confirmed in 20% of individuals' on a second MSLT who were not diagnosed during their first MSLT. Therefore repeat MSLT's can be done in the following circumstances:

- Narcolepsy and or idiopathic hypersomnia are still reasonably suspected but were not confirmed on the prior PSG/MSLT
- A previous diagnosis of narcolepsy or IH is now in question due to a change in symptoms

Maintenance of wakefulness testing (MWT)-indications and criteria (SL-2.4)

SL.ST.204.A

v1.0.2025

Maintenance of Wakefulness Testing-Indications

- Maintenance of Wakefulness Testing (MWT) using a 40-minute protocol is medically necessary when **ALL** of the following criteria are met:
 - The individual is stable with adherence to effective treatment for their sleep disorder
 - If the individual is being treated with positive airway pressure (PAP) for obstructive sleep apnea (or other types of sleep disordered breathing), optimal efficacy and adherence has been achieved by demonstration of **ALL** of the following:
 - PAP download shows: AHI <5 on download **AND**
 - Currently using PAP ≥4 hours per night on 70% of nights during a consecutive thirty (30) day period during the preceding 6 months **AND**
 - Therapy has resolved symptoms of increased upper airway resistance (i.e., eliminated snoring).
 - If the individual is on treatment with non-PAP therapy for obstructive sleep apnea, optimal efficacy and adherence has been achieved by demonstration of **ALL** of the following:
 - Self-reported adequate use **AND**
 - Efficacy confirmed on sleep testing with AHI <5/hour **AND**
 - Therapy has resolved symptoms of increased upper airway resistance (i.e., eliminated snoring)

Background and Supporting Information

Maintenance of Wakefulness Testing (MWT) measures the ability to stay awake for a defined period of time. Practice parameters on the clinical use of MWT were published by the American Academy of Sleep Medicine (AASM) in 2005 and in 2021. The MWT should be performed following the individual's major sleep period. The MWT should be conducted when an individual is clinically stable and when treatments of any known sleep disorders are well-established and effective. Unlike MSLT, the performance of overnight polysomnography the night prior to MWT is at the discretion of the sleep clinician. Per the AASM, the MWT 40-minute protocol is recommended. Clinical guidelines specify that MWT may be indicated to assess response to treatment in individuals' with sleep disorders associated with excessive daytime sleepiness.

MWT may also be useful to assess ability to maintain wakefulness when hypersomnia constitutes a public or personal safety concern. However, the utility of MWT is limited by the lack of large scale studies providing normative data for mean sleep latency on MWT. In addition, assessment of the daily ability to maintain wakefulness is complex and influenced by several variables not assessed during MWT such as long term treatment compliance, sleep duration and quality, circadian factors and shift work schedules.

The 2021 AASM practice parameters for adults indicate that:

- "Individual's on PAP/non-PAP therapies for sleep-disordered breathing should use them the night before (but not during) the MWT. If a PSG is performed, PAP settings and mask interface should match those used at home."
- "In individuals' with sleep-disordered breathing who are being evaluated for the effectiveness of therapy, the clinician should ensure the effectiveness (efficacy and adherence) based on review of downloaded data or self-reported use for non-PAP prior to testing."
- "The clinician should develop a plan regarding use of prescription medications, over the counter (OTC) agents, herbal remedies, and other substances. If the individual is chronically taking medications with alerting or sedating properties, they should be continued at a stable dose. Changes in medications should be avoided for two weeks prior to testing. The individual should be instructed to consult with the clinician before starting a prescription or OTC medication prior to the test."

Evidence Discussion

The Maintenance of Wakefulness Test (MWT) is a series of wake trials used to measure the ability to stay awake during soporific conditions. The American Academy of Sleep Medicine (AASM) recommends four 40-minute wake trials with sleep onset defined as the first epoch of sleep. The AASM protocol requires the study to be done when an individual is clinically stable and when treatments of any known sleep disorders are well-established and effective. In individuals' with sleep-disordered breathing on positive airway pressure (PAP), downloaded data should show efficacy and adherence.

Guidance from federal and state government agencies regarding fitness to drive with regard to sleep disorders are poorly detailed or are found only by gathering information from several different sources in the system. The U.S. Department of Transportation (DOT) and Federal Motor Carrier Safety Administration (FMCSA) recommend only repeat diagnostic testing for obstructive sleep apnea (OSA) if clinically indicated, but no recommendation on the use of the MWT. The Federal Aviation Administration (FAA) also does not require an MWT to evaluate for OSA treatment efficacy, but does include it as additional data when considering rescinding a prior diagnosis of narcolepsy or idiopathic hypersomnia.

The American Academy of Sleep Medicine (AASM) Practice Parameters recommends the MWT 40 minute protocol for when an individual's inability to remain awake constitute

a public or personal safety issue. Consideration of public safety in certain fields of employment where MWT may be used, such as commercial driving or flying, should be taken into account. The AASM also recommends the MWT to assess response to treatment in some situations involving individuals' with disorders of excessive sleepiness (e.g. medication response in individuals' with narcolepsy or idiopathic hypersomnia). However, the sleep clinician is advised not to rely on the MWT result as a single indicator of impairment or risk for accidents, but also to rely on clinical judgment including integration of clinical history and compliance with treatment.

Repeat sleep testing - home or attended sleep studies (SL-2.6)

SL.ST.206.A

v1.0.2025

Repeat Diagnostic Study

- Most risk factors for obstructive sleep apnea generally do not improve with the passage of time; therefore, follow-up HSAT or PSG is not routinely medically necessary for asymptomatic individuals' on PAP or non-PAP therapy. However, individuals' with persistent or recurrent symptoms despite adherence with OSA therapy may require repeat testing.
- Either home sleep apnea testing or in lab full night diagnostic sleep testing is medically necessary based on indications and comorbidities found in **General Guidelines (SL-1.0): Site of care for sleep testing requests (home versus facility)** when there is documentation of **ANY** of the following:
 - BMI decreases by 10% and there is a desire to discontinue PAP therapy and/or intolerance to PAP therapy and/or other planned change in management
 - BMI increases by 10% when test results would lead to a significant change in management
 - To reassess for the continued presence of OSA after **ANY** of the following:
 - Upper airway surgery for OSA both post-operatively after an appropriate period of healing (weeks to months) and repeat testing if there is concern for relapse after an initially successful surgery
 - At least 3 months after bariatric surgery when there is >10% weight loss
 - Individuals with a tracheostomy for OSA who are under consideration for decannulation (PSG or HSAT with trach capped)
 - OSA oral appliance trial
 - Positional therapy (not necessary if was proven effective on the diagnostic study)
 - Other non-PAP supportive interventions (e.g. nasal expiratory positive airway pressure, oral pressure therapy, eXciteOSA)
 - Repeat diagnostic testing is not medically necessary only to supply new PAP equipment or other new treatment modality (e.g. oral appliance, surgery) for sleep apnea in the absence of a significant change in clinical status that might alter sleep disordered breathing type or severity resulting in a change in management

- Results of previous medically necessary sleep test were inadequate and not diagnostic due to limited sleep time or other specified variables (report of prior sleep testing required).
 - Technically inadequate HSAT recordings lack a minimum of 4 hours of oximetry and flow data
- Results of previous medically necessary sleep test are unavailable and there is a documented effort by the requesting provider to obtain a copy of that test result.
 - See **General guidelines (SL-1.0)**

Evidence Discussion

Repeated routine sleep testing without positive impact on management of care, is not indicated. Aging as a sole indication for repeat diagnostic testing is not supported. In a subset of a large cohort of 427 randomly selected community-dwelling elderly subjects, age was found to have no statistically significant effect on RDI over an 18 year follow-up. RDI increase was associated with BMI increase and presence of self-reported high blood pressure. Supporting this association of RDI increase with BMI increase, and not age alone, was data from the Sleep Heart Health Study which showed a strong correlation between AHI and weight change.

When unnecessary repeat sleep testing is performed, a false negative result showing absence of sleep apnea may occur due to factors related to night to night variability.

This may result in loss of qualification for PAP therapy. Discontinuation of treatment would be harmful in this scenario, given the relationship between sleep apnea and neurocardiac morbidity and mortality. Significant time is saved in avoiding unnecessary testing and delay in re-initiation or continuation of treatment.

For some indications, a titration study is the best test, rather than a repeat diagnostic or split-night study. When a split night study is inappropriately performed instead of a titration study, the individual may be harmed during the study when a PAP machine is not being used and increases their risk of possible neurologic/cardiovascular adverse events. If a split night study is inappropriately performed, the individual would need to return to the laboratory for a full night titration, leading to a delay in care. Repeating a diagnostic study or diagnostic portion of a sleep study in lieu of a titration study creates a missed opportunity to obtain data about the titration of PAP therapy to make a decision on changes to therapy.

Repeat titration

- Most risk factors for obstructive sleep apnea generally do not improve with the passage of time; therefore, follow-up HSAT or PSG is not routinely medically necessary for asymptomatic individuals' on PAP or non-PAP therapy. However,

individuals' with persistent or recurrent symptoms despite adherence with OSA therapy may require repeat testing.

- A repeat titration study is not medically necessary to assess for the efficacy of PAP therapy in absence of recurrent or changed symptoms.
- Must demonstrate that recurrent or continued symptoms are not due to insufficient compliance (must be using PAP ≥ 4 hours per night on 70% of nights with continued symptoms).
- A repeat titration study is medically necessary if **ANY** of the following criteria is met:

OSA currently on CPAP

- Re-assessment of treatment results with in lab full night PAP titration **OR** unattended APAP based on indications and comorbidities found in **General Guidelines (SL-1.0): Site of care for sleep testing requests** currently using CPAP ≥ 4 hours on 70% of nights during a consecutive 30 day period anytime during the past 1 year is medically necessary when **ANY** of the following has occurred:
 - Substantial weight gain (10% of body weight) with return of symptoms
 - BMI decreases by 10% and there is intolerance of PAP pressure (compliant download not required)
 - Clinical response is insufficient due to ongoing symptoms
 - Symptoms return despite a good initial response to CPAP
 - Development of hypertension or worsening of hypertension despite a minimum of three months of adherent PAP usage as defined above
 - New onset decompensated heart failure or new stroke or TIA in an individual treated with PAP therapy
 - Newly discovered arrhythmia (such as atrial fibrillation) or bradycardia (such as atrioventricular block), particularly if the arrhythmia occurs during sleep, in an individual treated with PAP therapy
 - PAP machine download demonstrates elevated AHI with return of symptoms
 - A repeat titration study is not medically necessary only to supply new PAP equipment in the absence of a significant change in clinical status that might alter sleep disordered breathing type or severity resulting in a change in management
 - Results of previous PAP titration sleep test were inadequate and not diagnostic due to limited sleep time or other specified variables (report of prior sleep testing required)
 - Results of previous medically necessary sleep test are unavailable and there is a documented effort by the requesting provider to obtain a copy of that test result. See **General guidelines (SL-1.0)**

- CPAP/APAP is ineffective (as defined in the bullet points above) or is not tolerated and bi-level PAP titration utilizing HCPCS E0470 or E0471 is requested.

Sleep disordered breathing currently treated with bi-level PAP, APAP, ASV

- Re-assessment of treatment results with in lab full night PAP titration for an individual with known OSA currently using bilevel PAP, APAP, ASV ≥ 4 hours on 70% of nights during a consecutive 30 day period anytime during the past 1 year is medically necessary when **ANY** of the following has occurred:
 - Substantial weight gain (10% of body weight) with return of symptoms
 - BMI decreases by 10% and there is intolerance of PAP pressure (compliant download not required)
 - Clinical response is insufficient due to ongoing symptoms
 - Symptoms return despite a good initial response to CPAP
 - Development of hypertension or worsening of hypertension despite a minimum of three months of adherent PAP usage (as defined above).
 - New onset decompensated heart failure or new stroke or TIA in an individual treated with PAP therapy
 - Newly discovered arrhythmia (such as atrial fibrillation) or bradycardia (such as atrioventricular block), particularly if the arrhythmia occurs during sleep, in an individual treated with PAP therapy
 - PAP machine download demonstrates elevated AHI with return of symptoms.
 - A repeat titration study is not medically necessary only to supply new PAP equipment in the absence of a significant change in clinical status that might alter sleep disordered breathing type or severity resulting in a change in management
 - Results of previous PAP titration sleep test were inadequate and not diagnostic due to limited sleep time or other specified variables (report of prior sleep testing required)
 - Results of previous medically necessary sleep test are unavailable and there is a documented effort by the requesting provider to obtain a copy of that test result. See **General guidelines (SL-1.0)**

Evidence Discussion

The adverse outcomes of untreated sleep disordered breathing and benefits of treatment with positive airway pressure are well described in the literature. It is reasonable for a repeat titration/treatment study to be performed in certain clinical scenarios to optimize therapy.

Routine reassessment of asymptomatic individuals' with OSA on PAP therapy is not medically necessary. There should be sufficient subjective and objective reason to

suggest a retitration. Regular titration testing is unnecessary and possibly misleading and can result in inadequate titrations or the conclusion that Bilevel PAP is needed when CPAP therapy is sufficient.

In individuals' with known OSA, appropriate redirection to a full night of titration from a requested diagnostic or split night study can be beneficial in terms of allowing extra time during the sleep period to find the optimal settings for the PAP therapy. Performing the requested diagnostic study instead of an appropriate titration study causes harm by potentially necessitating a further titration study in the future and thereby delaying optimal care. Performing a split night study instead of a full night titration study, in the appropriate circumstances, can be harmful to the individual, due to limitation of the amount of time allotted for acquisition of titration data, increasing the chances for an ineffective or indeterminate titration. This leads to delays in optimal care. Lack of effective titration data leads to PAP therapy non-compliance, which is harmful, as noted above.

Diagnostic Testing pre- and post-hypoglossal nerve stimulator (HGNS) implantation (SL-2.8)

SL.ST.208.A
v1.0.2025

Indications

PSG (in lab full night diagnostic sleep study) is medically necessary for the following:

Pre-implantation

No prior sleep testing:

- Individuals with a high pre-test likelihood for moderate to severe obstructive sleep apnea who have not undergone prior sleep testing should undergo home sleep apnea testing, if appropriate per guidelines, and a PAP trial before consideration of facility testing for possible hypoglossal nerve stimulator implantation. Please see sections **General Guidelines (SL-1.0): Site of care for sleep testing requests (home versus facility)**

Prior diagnosis of OSA based on polysomnography or home sleep apnea testing:

- Pre-implant in lab full night diagnostic sleep study is medically necessary when:

ALL of the below are met:	AND	ONE of the below is met:
BMI \leq 40		Individuals have never had an in lab full night diagnostic sleep study (diagnosis based on HSAT only)
AHI or REI \leq 100		In lab full night diagnostic sleep study was performed > 24 months ago
Continued PAP intolerance after attempting to use it during a recent minimum one month trial including but not limited to mask fittings, different masks, pressure changes, or desensitization.		In lab full night diagnostic sleep study was performed \leq 24 months ago and there have been changes in weight or symptoms to suggest a clinically significant change in sleep study results

ALL of the below are met:	AND	ONE of the below is met:
Evaluation for HGNS should be performed by a board eligible or certified sleep physician or any provider in consultation with the sleep specialist.		

- For individuals age 13-18 years old with Down Syndrome, pre-implant full night diagnostic sleep study is medically necessary when:

ALL of the below are met:	AND	ONE of the below is met:
AHI or REI < 50		Individuals have never had an in lab full night diagnostic sleep study (diagnosis based on HSAT only)
Adenotonsillectomy is contraindicated or ineffective		In lab full night diagnostic sleep study was performed > 6 months ago
Continued PAP intolerance after attempting to use it during a recent minimum one month trial including but not limited to mask fittings, different masks, pressure changes, or desensitization		In lab full night diagnostic sleep study was performed ≤ 6 months ago and there have been changes in weight or symptoms to suggest a clinically significant change in sleep study results
Evaluation for HGNS should be performed by a board eligible or certified sleep physician or any provider in consultation with the sleep specialist		

Post-implantation:

- As per the clinical trial, polysomnography (in lab full night diagnostic sleep study) is medically necessary at least 2 months post-implantation for the purpose of titrating device parameters and determining therapeutic stimulation settings.
- For individuals age 13-18 years old with Down Syndrome, an additional acclimation polysomnography (in lab full night diagnostic sleep study) can be performed the evening of device activation.
- Following the initial post-implantation study, retesting (either HSAT or PSG-in lab full night diagnostic sleep study) is medically necessary if **ANY** of the following occurs:

- Clinical response is insufficient despite regular treatment with hypoglossal nerve stimulator
- Substantial weight gain with return of symptoms
- Results of previously medically necessary sleep test were inadequate due to limited sleep time or other variables (report of prior sleep testing is required)
- Following the initial post-implantation study, the choice of home sleep apnea testing or facility polysomnography for repeat testing will be based on indications and co-morbidities outlined in **General Guidelines (SL-1.0)**.

Background and Supporting Information

A hypoglossal nerve stimulator is a surgically implanted device that delivers stimulating electrical pulses to the hypoglossal nerve, which controls upper airway musculature. With a sensing lead, the device permits synchronization with ventilatory effort. The Stimulation Treatment for Apnea Reduction (STAR) trial was a prospective, multicenter trial of 126 participants with a body mass index (BMI) $<32 \text{ kg/m}^2$ moderate to severe obstructive sleep apnea (AHI 20-50), and difficulty tolerating/adhering to CPAP. Participants, who served as their own control, experienced a significant reduction in Apnea Hypopnea Index with hypoglossal nerve stimulation (68% decrease) and oxygen desaturation index (70% decrease) at 12 months, as well as a reduction in self-reported outcomes at 12 and 24 months. These improvements were maintained at 3 and 5 years. During the trial, a response to hypoglossal nerve stimulation at one year was defined as a reduction in baseline AHI by at least 50% with a final AHI of < 20 events per hour at one year (Sher's criteria for surgical success).

In 2017, the Food and Drug Administration revised the criteria to include individuals with AHI between 15 and 65. This was based on a prospective, single arm study conducted in Germany which utilized an inclusion criteria of AHI of 15 to 65 per hour and BMI $<35 \text{ kg/m}^2$. Significant reduction in AHI was achieved with median AHI decreasing from 28/h to 8.3/h at 6 months and sustained improvement at one year.

In 2020, the FDA added to its premarket approval individuals between the ages of 18 and 21 years old for whom adenotonsillectomy was not effective or contraindicated based on extrapolation from the STAR Trial and an ongoing pediatric Down Syndrome study. The BMI $\leq 32 \text{ kg/m}^2$ requirement did not apply, but the requirements of PAP failure/intolerance and lack of complete concentric collapse at the soft palate level on drug-induced sleep endoscopy (DISE) did apply to this new population. Very small studies of adults with OSA and tonsillar hypertrophy show mixed success after tonsillectomy (47.6 – 100% met Sher's criteria).

In 2023 the FDA expanded the label indications for HGNS therapy to include:

- Expansion of the upper limit baseline apnea-hypopnea index (AHI) from ≤ 65 to ≤ 100
- Expansion of the upper limit body mass index (BMI) from ≤ 32 to $\leq 40 \text{ kg/m}^2$

- Individuals with Down Syndrome between the ages of 13 and 18 years old with AHI >10 and <50 who qualify for the use of HGNS by the same indications as those between the ages of 18 and 21 years old. This was based on a 2022 trial of 42 individuals age 10 to < 22 years old with Down Syndrome who demonstrated safety and sustained efficacy at 12 months (65.9% met Sher's criteria).

The post-market approval ADHERE registry (5000 individuals enrolled since 10/2016) showed preliminary results of individuals implanted with higher AHI and BMI levels. 39 individuals with AHI > 65 and ≤ 100 showed a response rate of 66.7% per Sher's criteria. Similarly, a large 2022 retrospective study using ADHERE registry data analyzed the outcomes of 1963 subjects stratified into 5 different baseline AHI subgroups. There was no significant difference in treatment success (defined by Sher's criteria) between the subgroups. However, the conclusions are limited by small numbers, with only 33 subjects in the subgroup with AHI > 65 events/ hr.

Preliminary results from the ADHERE registry showed 216 individuals with BMI >32 kg/m² and ≤40kg/m² who had a response rate of 60.6% per Sher's criteria. Some non-congruent data exist regarding the optimal BMI cutoff for HGNS therapy. A retrospective case-control study of 153 individuals comparing the clinical outcome between cohorts with BMI >32 kg/m² and <32 kg/m² showed comparable results. Previous data from the ADHERE registry noted a 8.5% reduced odds of treatment success for each unit increase in BMI. More than half of individuals with BMI >32 kg/m² demonstrated AP collapse pattern on DISE. The correlation between higher AHI and BMI with an unfavorable pattern of airway collapse on the DISE procedure is modest at best. Therefore, there are individuals with BMI >32 kg/m² who may benefit from HGNS after a DISE showing AP collapse.

Clinical practice guidelines by the AASM and the American Academy of Otolaryngology-Head and Neck Surgery (AAO-HNS) specific to HGNS are not yet available. However, the German Society of Oto-Rhino-Laryngology, Head and Neck Surgery's 2021 Position Paper on HGNS recommends "besides suitable evaluation regarding any [HGNS], the reasons for PAP failure need to be documented during the screening process. It is, therefore, advisable to check for PAP optimization options and to evaluate other OSA treatment options in the individual decision-making process." In addition, sleep medicine expertise is recommended: "Ideally, individual selection, DISE, implantations, therapy adjustment and control are carried out in a department of otorhinolaryngology with profound sleep medicine knowledge or in a sleep medicine center under surveillance of the implanting surgeon."

A 2020 post-implant care pathway was produced by 5 high-volume HGNS centers in the US based on their experience. According to the 5-year STAR trial, residual disease burden may be reported anywhere from 10% (residual snoring) to 22% (residual AHI >15 or ESS >10) of HGNS individuals. Pathways were created for individuals with a combination of suboptimal adherence, suboptimal AHI, and/or suboptimal symptomatic

response. Assessment and management recommendations require expertise of both sleep and surgical specialists.

Evidence Discussion

Hypoglossal nerve stimulation (HGNS) was demonstrated as safe and effective by the Stimulation Therapy for Apnea Reduction Trial (STAR) and was initially approved by the Food and Drug Administration (FDA) for use in adults 22 years of age and older.

FDA expanded approval for individuals ages 18-21 years old, pediatric individuals with Down Syndrome ages 13-18 years, and for individuals with BMI from <32 to <40 upper apnea-hypopnea index (AHI) limit from < 65 to < 100.

Post-implantation polysomnogram (PSG) testing is recommended by the American Academy of Sleep Medicine (AASM) for titration of stimulation parameters to optimized outcomes for OSA. The AASM also recommends periodic clinical assessment since OSA may recur postoperatively. In individuals treated with PAP, repeat titration is indicated in the setting of significant weight gain with re-emergence of symptoms. This recommendation can be extrapolated to individuals treated with HGNS.

Diagnostic Testing pre- and post transvenous phrenic nerve stimulator (TPNS) implantation (SL-2.9)

SL.ST.0002.9.A

v1.0.2025

Indications

In lab full night diagnostic sleep study is medically necessary for the following:

Pre-implantation

Prior diagnosis of CSA based on PSG or HSAT done over 12 months ago or HSAT done within the past 12 months:

Polysomnography (in lab full night diagnostic sleep study) is medically necessary when **ALL** of the following are met:

- ≥ 18 years old
- Moderate to severe central sleep apnea seen on prior study as defined by central AHI ≥ 15 /hour
- CSA is not primarily attributed to opioid pain medications
- Absence of medical conditions that could transiently increase the severity of CSA on the PSG, including but not limited to:
 - Stroke or transient ischemic attack within the past 6 months
 - New diagnosis of heart failure less than 6 months ago
 - Left ventricular systolic dysfunction (EF $\leq 40\%$) not on optimal medical therapy (including a beta-blocker with **ONE** of the following: ACE inhibitor, angiotensin II receptor blocker or angiotensin receptor-neprilysin inhibitor) unless denoted contraindication to above
- Positive airway pressure failure or intolerance after attempting to use it, or is contraindicated
- Failed supplemental oxygen therapy
- Ordered by a board eligible or board certified sleep medicine physician or any provider in consultation with the sleep physician

Prior diagnosis of CSA based on PSG done within the past 12 months:

Repeat polysomnography (in lab full night diagnostic sleep study) is medically necessary when **ALL** of the following are met:

- There have been changes in weight, symptoms, medications, or neurological or cardiovascular status to suggest a clinically significant change in sleep study results.
- ≥ 18 years old
- Moderate to severe central sleep apnea seen on prior study as defined by central AHI ≥ 15 /hour
- CSA is not primarily attributed to opioid pain medications
- Absence of medical conditions that could transiently increase the severity of CSA on the PSG, including but not limited to:
 - Stroke or transient ischemic attack within the past 6 months
 - New diagnosis of heart failure less than 6 months ago
 - Left ventricular systolic dysfunction (EF $\leq 40\%$) not on optimal medical therapy (including a beta-blocker with **ONE** of the following: ACE inhibitor, angiotensin II receptor blocker or angiotensin receptor-neprilysin inhibitor) unless denoted contraindication to above
- Positive airway pressure failure or intolerance after attempting to use it, or is contraindicated
- Failed supplemental oxygen therapy
- Ordered by a board eligible or board certified sleep medicine physician or any provider in consultation with the sleep physician

Post-implantation

Polysomnography (in lab full night diagnostic sleep study) with interrogation and programming of the phrenic nerve stimulator system can be performed at least 2 months after implantation for the purpose of titrating device parameters and determining therapeutic and tolerable stimulation settings.

Following the initial post-implantation study, retesting with an in lab full night diagnostic sleep study can be performed if **ANY** of the following occurs:

- Clinical response is insufficient despite compliant use
- Results of previously medically necessary sleep test were inadequate due to limited sleep time or other variables (report of prior sleep testing required)

Background and Supporting Information

Central sleep apnea (CSA) was found to be an independent risk factor for atrial fibrillation (AF) in the Sleep Heart Health Study. CSA demonstrated double the risk for AF. There is a strong association between CSA and AF both in congestive heart failure (CHF) and idiopathic CSA. Possible mechanisms include CSA as a marker for cardiac dysfunction and CSA induced carbon dioxide fluctuations with arousals and oxygen desaturations leading to sympathetic over activation which then leads to arrhythmogenic

structural and electrical remodeling. In a prospective observational cohort study of 784 hospitalized individuals with systolic heart failure, CSA was found in 165 individuals (21%) and was found to be a predictor of 6 month cardiac readmission.

A meta-analysis of 19 randomized studies that included individuals with $AHI \geq 10$ /hour (predominantly CSA) and heart failure with reduced ejection fraction (HFrEF) $\leq 50\%$ showed short-term benefit of ASV, CPAP, and drug therapies on sleep, but long-term quality of life and cardiovascular benefit were lacking. Acetazolamide and theophylline require monitoring for use, and concerns exist with using these therapies in the HFrEF population. Transphrenic nerve stimulation (TPNS) demonstrated positive treatment outcomes on sleep and quality of life at 6 months with suggestion of improved cardiovascular outcomes. With ASV being contraindicated at this time for some heart failure individuals. TPNS is being considered for both these individuals and others who are not tolerating and or not experiencing benefit from non surgical therapies for CSA.

The Remede® System consists of a neurostimulator implanted into the right or left pectoral region and a combined stimulation and sensing lead implanted into the left pericardiophrenic vein or right brachiocephalic vein. There is an optional respiratory sensing lead placed in the azygos vein in older models where the stimulation lead did not sense respiration well. The stimulation lead transvenously stimulates only a single phrenic nerve, but both diaphragms move in response, resulting in a full breath.

The Pivotal Trial was a prospective, multicenter (31 hospital-based centers in Germany, Poland, USA) randomized trial that recruited treatment ($n = 73$) and control ($n = 78$) groups between 04/17/2013 and 05/28/2015. At baseline in both groups, the mean AHI was 46/hour, mean central apnea index (AI) was 24/hour, percent of sleep with O_2 saturation $\geq 90\%$ (T90%) was 10%, and Epworth Sleepiness Scale Score (ESS) was 8. 51% of the treatment group had an AHI reduction of 50% or greater at 6 months as compared to only 11% of the control group. At 5 year follow-up, pooled results of both treatment and control groups (activated 6 months later) showed sustained improvements in the AHI (24/hour), central AI (2/hour), T90% (3%), and ESS.

Analysis of subgroups showed similar sustained efficacy in PAP-treated versus PAP-naïve groups, 10 idiopathic central sleep apnea (16 participants which made up 11% of Pivotal Trial), and heart failure with either reduced or preserved ejection fraction (96 participants which made up 64% of the Pivotal Trial). Heart failure related hospitalization rates were 4.7% in the treatment group and 17.0% in the control group. There was no detectable difference in cardiovascular mortality in the first 6 months between the treatment and control groups. After 12 months of therapy, there was a statistically significant increase in left ventricular ejection fraction from 31.6% to 34.8% ($p = 0.0004$) as well as a statistically significant decrease in the Minnesota Living with Heart Failure score from 40.0 to 27.0 ($p = 0.005$).

The freedom from serious adverse events related to the implantation procedure, the system, or the delivered therapy was 91%. At 1, 3, and 5 years, 9%, 10%, and

14% of individuals had serious adverse events related to the implant procedure, device, or delivered therapy. These events were related to the implant site (impending erosion, hematoma, infection), lead failure/dislodgement/displacement, extra respiratory stimulation, concomitant device interaction, non-cardiac chest pain, and elevated transaminase. The two most frequent non-serious therapy-related complaints were abdominal discomfort near the diaphragm and sensation in areas remote from the diaphragm.

Based on sustained efficacy and low rate of serious adverse events, the Remede® System received FDA premarket approval on 10/06/2017 for the treatment of moderate to severe central sleep apnea (CSA) in adult individuals. Approval for adding MR Conditional labeling for 1.5T and 3T MRI scans to the implantable components was issued 03/28/2023.

The inclusion and exclusion criteria in the Pivotal Trial were extensive. Sleep study inclusion metrics were AHI ≥ 20 , central AI $\geq 50\%$ of all apneas with ≥ 30 central apnea events, and obstructive AI $\leq 20\%$ of the total AHI. In addition, there were numerous criteria for medical stability, including heart failure diagnosis being ≥ 6 months and systolic heart failure optimized with guideline-directed medical therapy (GDMT). Guideline-directed medical therapy (GDMT) is met when heart failure medications (e.g. beta-blockers, angiotensin-converting enzyme inhibitors, and others listed in the current guidelines) have been titrated up to trial defined target dosages as tolerated or are considered contraindicated or not medically necessary.

Stroke within 12 months of baseline testing was an exclusion criterion. Prospective studies and meta-analyses have shown a high prevalence of sleep apnea (both central and obstructive) after transient ischemic attacks (TIA's) and strokes without clear association with severity, neurological location, and vascular involvement. Central apneas have been shown to increase and decrease by 3 and 6 months post stroke/TIA. Therefore, stroke of any type and TIA within the prior 6 months was included in this guideline to ensure a PSG to qualify for the Remede® System was done when the CSA diagnosis was most stable.

CSA primarily attributed to opioids was an exclusion criterion. Although ASV is equally successful at reducing the CSA in opioid use and CHF, no randomized controlled trials have been done to assess the clinical benefits of opioid-induced CSA treatment such as reduction in morbidity and mortality. While a large cohort of 6002 CSA individuals at the VA showed all-cause CSA was associated with 1.5 times higher odds of cardiac hospital admission and chronic opioid users had 1.17 times increased odds, those opioid users with CSA did not. Increased morbidity and mortality in chronic opioid users has been demonstrated in multiple studies, but sleep disordered breathing was not assessed in these studies. A study of 50 methadone maintenance treatment individuals found increased daytime sleep propensity and reduced daytime function, but multiple regression analysis showed that the severity of CSA did not contribute to either of

these problems. Morbidity of untreated opioid induced CSA and clinical benefit from its treatment aside from AHI reduction has not been established in large studies.

The inclusion and exclusion criteria in the post-market reST Study were considerably simpler. The main inclusion criterion was moderate to severe central sleep apnea defined as AHI ≥ 15 /hour based on a PSG done within 12 months of the expected implant date and age ≥ 18 years old. The extensive exclusion criterion in the Pivotal Trial were limited to being able to do the study and not being or planning to become pregnant. Enrollment of up to 500 individuals in this multi-center, prospective, open label, non-randomized study started July 2019 and is estimated to be completed 6/2032. Safety and effectiveness data will be collected as well as heart failure specific outcomes in the heart failure subgroup.

ZOLL Medical's Patient Pathway Guide is based on the Pivotal Trial and reST Study protocols where the device is activated 1 month after implantation followed by office visits, PSG, or HSAT to assess efficacy and adjust therapy per each site's standard practice. PSG's and HSAT's for efficacy (not titration) at 6, 12, 18, and 24 months and HSAT at 36 and 60 months were specific to the trials. In ZOLL Medical's Patient Pathway Guide, 2-3 office visits occurring every 6-12 weeks to optimize therapy are done before usually PSG or HSAT is done to confirm efficacy and for final device optimization. HSAT's can report central apneas, but small studies show variable and low sensitivity. The American Academy of Sleep Medicine's 2017 Position Statement states that HSAT is an alternative to PSG for the diagnosis of OSA in uncomplicated individuals presenting with signs and symptoms that indicate an increased risk for moderate to severe OSA. Therefore, PSG and not HSAT for titration is included in this guideline.

Failure of nonsurgical therapies was not a requirement in either the Pivotal Trial or the reST study. However, the American Academy of Sleep Medicine (AASM) 2012 Practice Parameters on the Treatment of Central Sleep Apnea Syndromes (CSAS) in Adults recommended CPAP and nocturnal oxygen as standard treatment for CSAS related to CHF. After the SERVE-HF trial, an update in 2016 lowered the standard recommendation for adaptive servo-ventilation (ASV) to an option for CHF-associated CSAS in individuals with left ventricular ejection fraction $>45\%$ or mild CSAS. BPAP in spontaneous time (ST) mode is an option for CSAS related to CHF if there has been no response to adequate trials of CPAP, ASV, and oxygen therapies. Acetazolamide and theophylline are also only considered options for CSAS related to CHF and primary CSAS. Based on these practice parameters and a meta-analysis of CSA therapies, failure of mask based therapy and oxygen were included in this guideline, but not medication trials.

Experience with the hypoglossal nerve stimulator (HGNS) has shown that sleep medicine expertise is recommended for both individual selection and post implantation care. PAP failure despite optimization should be documented pre implantation.

Suboptimal adherence, AHI, and symptomatic response may occur post implantation and requires expertise of both sleep and surgical specialists.

Treatment of Sleep-related Breathing Disorders

Guideline	Page
Treatment - General information.....	55
Positive airway pressure devices (SL-4.2).....	60
Oral Appliances for the Treatment of Obstructive Sleep Apnea (SL-9).....	72

Treatment - General information

SL.TX.101.A

v1.0.2025

Treatment coding (SL-4.1.1)

The codes for treatment of obstructive sleep apnea include HCPCS and CPT®

The inclusion of any code in this table does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

Treatment codes

Treatment codes	HCPCS and CPT®
Continuous airway pressure (CPAP/APAP) device	E0601
Respiratory assist device, bi-level pressure (BPAP) capability, WITHOUT backup rate feature, used with noninvasive interface, e.g. nasal or facial mask (intermittent assist device with continuous positive airway pressure device)	E0470
Respiratory assist device, bi-level pressure (BPAP) capability (including ASV), WITH backup rate feature, used with noninvasive interface, e.g. nasal or facial mask (intermittent assist device with continuous positive airway pressure device)	E0471
Respiratory assist device, bi-level pressure (BPAP) capability, WITH backup rate feature, used with invasive interface, e.g. tracheostomy tube (intermittent assist device with continuous positive airway pressure device)	E0472
Humidifier, non-heated, used with positive airway pressure (CPAP/BPAP/APAP) device	E0561

Treatment codes	HCPCS and CPT®
Humidifier, heated, used with positive airway pressure (CPAP/BPAP/APAP) device	E0562
Tubing with heating element	A4604
Combination oral/nasal mask	A7027
Replacement oral cushion combo mask	A7028
Replacement nasal pillow comb mask	A7029
CPAP full face mask	A7030
Replacement facemask interface	A7031
Replacement nasal cushion	A7032
Replacement nasal pillows	A7033
Nasal interface (mask or cannula type) used with PAP device	A7034
Positive airway pressure headgear	A7035
Positive airway pressure chinstrap	A7036
Positive airway pressure tubing	A7037
Positive airway pressure filter	A7038
Filter, non-disposable w/ PAP	A7039
PAP oral interface	A7044
Replace exhalation port	A7045
Replacement, water chamber, PAP device	A7046

Treatment codes	HCPCS and CPT®
Monitoring feature/device, stand-alone or integrated, any type, includes all accessories, components and electronics, not otherwise classified (this code relates to Compliance and the data download of an individuals PAP therapy).	A9279
CPAP initiation and management (code is used to report the initiation and instruction when a patient begins therapy)	CPT 94660

PAP - General requirements (SL-4.1)

Valid sleep testing demonstrates a positive diagnosis of **ONE OR MORE** of the following sleep-related breathing disorders:

Obstructive sleep apnea (OSA) as measured by valid testing is defined as:

- The predominantly obstructive apnea-hypopnea index (AHI) or respiratory disturbance index (RDI) on a PSG, or the apnea-hypopnea index (AHI) or respiratory event index (REI) on a HSAT is ≥ 15 events per hour; **OR**
- The predominantly obstructive apnea-hypopnea index (AHI) or respiratory disturbance index (RDI) on a PSG, or the apnea-hypopnea index (AHI) or respiratory event index (REI) on a HSAT is ≥ 5 and < 15 events per hour with documentation of **ANY** of the following:
 - Symptoms of sleepiness, fatigue, insomnia, or other symptoms leading to impaired sleep-related quality of life
 - Report of awakening with breath holding, gasping, or choking
 - Bed partner or other observer reports habitual snoring, breathing interruptions, or both during sleep
 - Hypertension, a mood disorder, cognitive dysfunction, coronary artery disease, congestive heart failure, atrial fibrillation, type 2 diabetes mellitus, or stroke

Central sleep apnea (CSA) is defined as ALL of the following:

- Presence of **ONE OR MORE** of the following:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or non-restorative sleep
 - Awakening short of breath

- Witnessed apneas
- Central apnea and/or central hypopnea index ≥ 5 per hour
- Central hypopnea/apneas are $>50\%$ of the total number of apneas and hypopneas

Central sleep apnea (CSA) with Cheyne-Stokes Breathing is defined as ALL of the following:

- Presence of **ONE OR MORE** of the following:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or non-restorative sleep
 - Awakening short of breath
 - Witnessed apneas
 - Known atrial fibrillation/flutter, congestive heart failure, or a neurological disorder
- PSG must show **ALL** of the following:
 - Central apnea and/or central hypopnea index ≥ 5 per hour
 - Central hypopnea/apneas are $\geq 50\%$ of the total number of apneas and hypopneas
 - Pattern of breathing meets criteria for Cheyne-Stokes Breathing (episodes of ≥ 3 consecutive central apneas and/or central hypopneas separated by a crescendo and decrescendo change in breathing amplitude with a cycle length of ≥ 40 seconds)

Treatment Emergent Central Sleep Apnea is defined as ALL of the following:

- Presence of **ONE OR MORE** of the following thought to be attributable to the central events:
 - Sleepiness
 - Difficulty initiating or maintaining sleep, frequent awakenings or nonrestorative sleep
 - Awakening short of breath
 - Witnessed apneas
- Diagnostic PSG demonstrates ≥ 5 predominantly obstructive respiratory events per hour of sleep (obstructive or mixed apneas, hypopneas, respiratory effort related arousals [RERAs]).
- PSG during use of positive airway pressure without a backup rate shows improvement of obstructive events and emergence or persistence of central apneas/hypopneas with **BOTH** of the following:
 - Central apnea and/or hypopnea index ≥ 5 per hour

- Central hypopnea/apneas are $\geq 50\%$ of the total number of apneas and hypopneas

Sleep-related hypoventilation is defined as when EITHER of the following occur during sleep:

- Increase in arterial PCO_2 , transcutaneous PCO_2 , or end-tidal PCO_2 to a value >55 mmHg for ≥ 10 minutes
- There is a ≥ 10 mmHg increase in arterial PCO_2 , transcutaneous PCO_2 , or end-tidal PCO_2 during sleep (compared to awake supine value) to a value >50 mmHg for ≥ 10 minutes

Results from a sleep study are used to determine the type of sleep apnea, the severity of the breathing disorder, and the most appropriate form of treatment. Depending on these factors, a variety of PAP devices, and location of titration of therapy, can be considered.

Positive airway pressure is the treatment of choice for the various forms of sleep apnea. Positive airway pressure (PAP) is produced by a flow generator and applied to the airway through nasal, oral, or oronasal mask interfaces

Current practice recommendations for CSA with CHF (SL-4.1.2)

- CPAP: Standard
- BPAP (including ST): Option if CPAP ineffective
- ASV:
 - OPTION if EF $>45\%$ or mild central sleep apnea syndrome
 - STANDARD AGAINST if EF $\leq 45\%$ with moderate/severe central sleep apnea syndrome

Positive airway pressure devices (SL-4.2)

SL.TX.102.A

v1.0.2025

Auto-adjusting positive airway pressure (APAP) in unattended setting (SL-4.2.1)

Initial Assessment of PAP pressure with APAP

Please see sections referring to initiation of E0601 and authorization of purchase for E0601 in **Continuous Positive Airway Pressure (CPAP) or Auto-Adjusting Positive Airway Pressure (APAP) Therapy (SL 4.2.2)**

Repeat Assessment of PAP pressure with APAP

E0601 APAP Treatment (for the purpose of determining appropriate PAP pressure for individual's currently on fixed pressure treatment with CPAP) is medically necessary for **ALL** of the following:

- A positive diagnosis of OSA, as measured by HSAT or PSG as defined in **General Guidelines (SL-1.0)**
- Attempted compliance with fixed CPAP (≥ 4 hours per night on 70% of nights) has not adequately treated signs and symptoms of OSA.
- Persistent symptoms or unimproved AHI/RDI in individual currently on fixed CPAP therapy, when the individual and/or their caregiver has received the following from the treating physician or supplier of the PAP device:
 - Instruction in the proper use and care of the equipment
 - Mask re-fitting or adjustment if necessary
 - Education for proper use of PAP accessories

Continuous positive airway pressure (CPAP) or Auto-adjusting positive airway pressure (APAP) therapy (SL-4.2.2)

Initiation of HCPCS E0601 PAP therapy and establishing compliance (**ALL** of the following):

- A positive diagnosis of OSA or central sleep apnea, as measured by HSAT or PSG as defined in **General Guidelines (SL-1.0) and Auto-Adjusting positive airway pressure (APAP) in unattended setting (SL-4.2.1)**
- The individual and/or their caregiver have received instruction from the treating physician and supplier of the CPAP device and accessories in the proper use and care of the equipment
- A compliance support plan between the treating physician and DME supplier has been established

Authorization for equipment purchase:

- PAP device must be used ≥ 4 hours per night on 70% of nights during a consecutive thirty (30) day period anytime during the first three (3) months of initial usage
- Individual has regular follow-up to evaluate symptoms

Repeat initiation of APAP/CPAP HCPCS E0601 device after failing an initial trial period is medically necessary when:

- A clinical evaluation by a provider has occurred to determine the etiology of the failure to respond to PAP therapy with a support plan to address these causes.

Extension of establishing compliance with E0601 (**ALL** of the following):

- Individual history includes **ONE** of the following:
 - Failure to resolve symptoms or unimproved AHI during initial compliance period, **OR**
 - Inconsistent usage of device related to improper fit, lack of education, intolerance of PAP therapy, or device malfunction
- Individual has received from the ordering physician or supplier of the PAP device in the past 30 days (**ALL** of the following):
 - Instruction in the proper use and care of the equipment
 - Mask refitting or adjustment if necessary
 - Education for proper use of PAP accessories

Replacement APAP/CPAP HCPCS E0601 device (**ALL** of the following):

- Continued resolution of symptoms and improved AHI on therapy
- Device had been consistently used ≥ 4 hours per night on 70% of nights
- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

Bi-level positive airway pressure- spontaneous mode (SL-4.2.3)

Initiation of HCPCS E0470 PAP therapy and establishing compliance (ALL):

- **ONE** of the following medical conditions must be documented in the individual's record:
 - Obstructive sleep apnea when **BOTH** of the following:
 - Diagnosis of OSA as defined in General Guidelines (SL-1.0)
 - CPAP (HCPCS E0601) has been tried and proven either ineffective or not tolerated, based on a therapeutic trial conducted in either a facility or a home setting
 - Central sleep apnea diagnosis (either primary central sleep apnea or central sleep apnea with Cheyne-Stokes Breathing) when **BOTH** of the following:
 - Valid testing meeting criteria for central sleep apnea as defined in General Guidelines (SL-1.0)
 - Significant improvement of the central events with the use of HCPCS E0470 device on the settings that will be prescribed for initial use at home.
 - Treatment-Emergent Central Sleep Apnea when **BOTH** of the following:
 - Valid testing meeting criteria for treatment emergent central sleep apnea as defined in General Guidelines (SL-1.0)
 - Significant improvement of the central events with the use of HCPCS E0470 device on the settings that will be prescribed for initial use at home.
 - Neuromuscular disease or restrictive thoracic disorder when **BOTH** of the following:
 - Neuromuscular disease (e.g. amyotrophic lateral sclerosis) or a severe thoracic cage abnormality (e.g. severe kyphoscoliosis, post-thoracoplasty), **AND**
 - **ONE** of the following:
 - Symptoms (orthopnea, dyspnea, morning headache, daytime sleepiness, unrefreshing sleep) with vital capacity <80%
 - An arterial blood gas PaCO₂, done while awake and breathing the individual's prescribed FiO₂, is ≥45 mmHg
 - End-tidal CO₂, transcutaneous CO₂, or venous blood gas ≥50 mmHg
 - Sleep oximetry, or sleep testing, demonstrates oxygen saturation ≤88% for ≥5 minutes of nocturnal recording time or ≤90% for ≥5% of the night
 - Vital Capacity (Forced Vital Capacity or Slow Vital Capacity) ≤50% of predicted
 - Maximal inspiratory pressure ≤60 cm of H₂O
 - Sniff nasal inspiratory pressure ≤40 cm of water

- Severe COPD when **BOTH** of the following:
 - An arterial blood gas PaCO_2 is ≥ 52 mm Hg done while awake and breathing the individual's prescribed FiO_2
 - OSA and CPAP treatment have been considered and ruled out (formal sleep testing not required)
- Obesity hypoventilation syndrome (OHS) as defined in **General Guidelines (SL-1.0): Site of Care** when **ANY** of the following:
 - $\text{BMI} \geq 45$
 - $\text{BMI} \geq 30 \text{ kg/m}_2$ plus **EITHER** awake arterial blood gas (ABG), end-tidal PCO_2 (EtPCO_2), or transcutaneous PCO_2 (TcPCO_2) with $\text{PCO}_2 \geq 45$ **OR** venous blood gas (VBG) showing a $\text{PCO}_2 \geq 50$ mmHg
 - $\text{BMI} \geq 30 \text{ kg/m}_2$ plus serum bicarbonate ≥ 27 with stated concern for obesity hypoventilation syndrome (OHS)
- Other hypoventilation syndrome (defined awake arterial blood gas, end-tidal PCO_2 (EtPCO_2), or transcutaneous PCO_2 with $\text{PCO}_2 \geq 45$ **OR** venous blood gas showing a $\text{PCO}_2 \geq 50$ mmHg) when it is due to **ONE** of the following:
 - Hypoventilation due to central respiratory drive depression (associated with medication, substance use, or other medical conditions).
 - Hypoventilation due to respiratory system failure other than COPD or neuromuscular disease/thoracic cage abnormalities (for example end-stage interstitial lung disease)
- Sleep Related hypoventilation as defined in **General Guidelines (SL-1.0)**

Continued HCPCS E0470 therapy after initial 3 months:

If indication is sleep disordered breathing categorized as OSA, central sleep apnea or treatment emergent central sleep apnea when **BOTH** of the following:

- PAP device must be used ≥ 4 hours per night on 70% of nights during a consecutive thirty (30) day period anytime during the first three (3) months of initial usage
- Individual has regular follow-up to evaluate symptoms

For **ALL** other indications:

- Individual has been re-evaluated by treating practitioner
- Documentation of continued use, benefit, and medical need

Repeat initiation of Bilevel PAP HCPCS E0470 device after failing an initial trial period is medically necessary when:

- A clinical evaluation by a provider has occurred to determine the etiology of the failure to respond to PAP therapy with a support plan to address these causes

Replacement Bilevel PAP HCPCS E0470 device

If indication is sleep disordered breathing categorized as OSA, central sleep apnea or treatment emergent central sleep apnea (**ALL** of the following):

- Continued resolution of symptoms and improved AHI/RDI on therapy
- Device had been consistently used ≥ 4 hours per night on 70% of nights
- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

For **ALL** other indications:

- Continued resolution of symptoms
- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

Bi-level positive airway pressure: spontaneous/timed mode (SL-4.2.4)

Initiation of HCPCS E0471 PAP Therapy and establishing compliance (**ALL**):

- **ONE** of the following medical conditions must be documented in the individual's record:
 - Central sleep apnea diagnosis (either primary central sleep apnea or central sleep apnea with Cheyne-Stokes Breathing) when **BOTH** of the following:
 - Valid testing meeting criteria for central sleep apnea as defined in **PAP General Requirements (SL-4.1)** and **General Guidelines (SL-1.0)**
 - Significant improvement of the central events with the use of HCPCS E0471 device on the settings that will be prescribed for initial use at home
 - See **Adaptive Servo Ventilation (ASV) Therapy (SL-4.2.6)** for guidelines specific to adaptive servo ventilation (ASV)
 - Treatment-Emergent Central Sleep Apnea when
 - Valid testing meeting criteria for treatment-emergent central sleep apnea as defined in **General Guidelines (SL-1.0)**
 - Significant improvement of the central events with the use of HCPCS E0471 device on the settings that will be prescribed for initial use at home
 - Neuromuscular disease or restrictive thoracic disorder:

- Neuromuscular disease (e.g. amyotrophic lateral sclerosis) or a severe thoracic cage abnormality (e.g. severe kyphoscoliosis, post-thoracoplasty) and **ONE** of the following:
 - Symptoms (orthopnea, dyspnea, morning headache, daytime sleepiness, unrefreshing sleep) with vital capacity <80%
 - An arterial blood gas PaCO₂, done while awake and breathing the individual's prescribed FiO₂, is ≥45 mm Hg
 - End-tidal CO₂, transcutaneous CO₂, or venous blood gas ≥50
 - Sleep oximetry or sleep testing demonstrates oxygen saturation ≤88% for ≥5 minutes of nocturnal recording time or ≤90% for ≥5% of the night
 - Vital Capacity (Forced Vital Capacity or Slow Vital Capacity) ≤50% of predicted
 - Maximal inspiratory pressure ≤60 cm of H₂O
 - Sniff nasal inspiratory pressure ≤40 cm of water
- Severe COPD when **BOTH** of the following:
 - An arterial blood gas PaCO₂ is ≥52 mm Hg done while awake and breathing the individual's prescribed FiO₂
 - OSA and CPAP treatment have been considered and ruled out (formal sleep testing not required)
- Obesity hypoventilation syndrome as defined in **General Guidelines (SL-1.0): Site of Care** when **ANY**:
 - Individual was recently hospitalized with acute-on-chronic respiratory failure with persistent awake hypoventilation at the time of discharge
 - Individual has been diagnosed with OSA that is not severe (AHI or RDI <30)
 - Individual has been diagnosed with severe OSA (AHI, RDI ≥30) and has ongoing hypoventilation despite 3 months of compliant CPAP or bilevel HCPCS E0470 use
- Other hypoventilation syndrome (defined awake arterial blood gas, end-tidal PCO₂ (ET PCO₂), or transcutaneous PCO₂ with PCO₂ ≥45 OR venous blood gas showing a PCO₂ ≥50 mmHg) when it is due to **ONE** of the following:
 - Hypoventilation due to central respiratory drive depression (associated with medication, substance use, or other medical conditions)
 - Hypoventilation due to respiratory system failure other than COPD or neuromuscular disease/thoracic cage abnormalities (for example end-stage interstitial lung disease).
- Sleep-related hypoventilation as defined in **General Guidelines (SL-1.0)**

Continued HCPCS E0471 therapy after initial 3 months:

If the indication is sleep disordered breathing categorized as OSA, central sleep apnea or treatment emergent central sleep apnea:

- PAP device must be used ≥ 4 hours per night on 70% of nights during a consecutive thirty (30) day period anytime during the first three (3) months of initial usage
- Individual has regular follow-up to evaluate symptoms

For **ALL** other indications:

- Individual has been re-evaluated by treating practitioner
- Documentation of continued use, benefit, and medical need

Replacement HCPCS E0471 device

If indication is sleep disordered breathing categorized as OSA, central sleep apnea or treatment emergent central sleep apnea (**ALL** of the following):

- Continued resolution of symptoms and improved AHI on therapy
- Device had been consistently used ≥ 4 hours per night on 70% of nights
- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

For **ALL** other indications:

- Continued resolution of symptoms
- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

Heated and non-heated humidifier (SL-4.2.5)**Initial set-up (heated HCPCS E0562 and non-heated HCPCS E0561)**

ALL of the following:

- When requested by treating physician and PAP device (HCPCS E0470/471 or E0601) has been approved
- No previous humidifier has been provided

Replacement heated or non-heated humidifier (HCPCS E0562 or E0561) device

ALL of the following:

- Continued resolution of symptoms and improved AHI on therapy
- Device had been consistently used ≥ 4 hours per night on 70% of nights
- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

Adaptive servo ventilation (ASV) therapy (SL-4.2.6)

Initiation of ASV Therapy (HCPCS E0471) and establishing compliance

- **ONE** of the following medical conditions must be documented in the individual's record:
 - Central sleep apnea (either primary central sleep apnea or central sleep apnea with Cheyne-Stokes Breathing):
 - Diagnosis of central sleep apnea as defined in **General Guidelines (SL-1.0) and PAP-General Requirements (SL-4.1)**
 - Mild CSA, **OR** Moderate to Severe CSA with EF $>45\%$
 - Treatment Emergent Central Sleep Apnea
 - Diagnosis of treatment emergent central sleep apnea as defined in **General Guidelines (SL-1.0)**
 - Central Sleep Apnea Syndrome due to opioid or substance use:
 - CPAP has been shown to be ineffective following a reasonable treatment attempt/trial, **AND**
 - Opioid therapy cannot be reduced or discontinued, **AND**
 - Central hypopnea/apneas are $>50\%$ of total, **AND**
 - Central apnea index ≥ 5 per hour.

Continued ASV therapy HCPCS E0471 after initial 3 months:

ALL of the following:

- PAP device must be used ≥ 4 hours per night on 70% of nights during a consecutive thirty (30) day period anytime during the first three (3) months of initial usage
- Individual has regular follow-up to evaluate symptoms

Replacement ASV HCPCS E0471 device

ALL of the following:

- Continued resolution of symptoms and improved AHI on therapy
- Device had been consistently used ≥ 4 hours per night on 70% of nights

- Device is not operating
- DME supplier has physically evaluated the device and determined that it is unable to be repaired
- Device to be replaced is no longer covered under a warranty

Continuous positive airway pressure ventilation (CPAP), initiation, and management (SL-4.2.7)

- Physician face-to-face service addressing PAP usage (CPT® 94660):
 - Physician application or adjustment of mask or pressure titration or PAP related service; **AND**
 - Service cannot be adequately provided by a certified or registered respiratory therapist, licensed clinician, or sleep technologist when within scope of practice per state regulations; **AND**
 - Another evaluation and management service is not performed

Continuous PAP, Bilevel PAP, or automatic PAP Loaner Rental (SL-4.2.8)

CPAP, APAP, or BPAP loaner rental for up to 30 days is considered medically necessary when there is a description of the device malfunction and documentation that equipment has been sent for repair/assessment

Positive airway pressure – accessories and supplies (SL-4.3)

PAP masks and parts

Combination oral/nasal mask, used with PAP, each (HCPCS A7027):

- Frequency: 1 per 3 months.
- No other PAP mask ordered (i.e., HCPCS A7030, A7034, or A7044)
- Provided data meets compliance criteria (device used ≥4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Oral cushion used with combination oral/nasal mask, replacement only (HCPCS A7028):

- Only compatible with HCPCS A7027 mask.
- Frequency: 2 per month.

- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Nasal pillows used with combination oral/nasal mask, replacement only, pair (HCPCS A7029):

- Only compatible with HCPCS A7027 mask.
- Frequency: 2 per month.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Full face mask used with PAP, each (HCPCS A7030):

- Frequency: 1 per 3 months.
- No other PAP mask ordered (i.e., HCPCS A7027, A7034, or A7044).
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Full face mask interface replacement, each (HCPCS A7031):

- Only compatible with HCPCS A7030 mask.
- Frequency: 2 per month.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Nasal interface (mask or cannula type) used with PAP, each (HCPCS A7034):

- Frequency: 1 per 3 months.
- No other PAP mask ordered (i.e., HCPCS A7027, A7030, or A7044)
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Cushion for use on nasal mask interface, replacement only, each (HCPCS A7032):

- Only compatible with HCPCS A7034 mask.
- Frequency: 2 per month.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Nasal pillow for use on nasal cannula type interface, replacement only, pair (HCPCS A7033):

- Only compatible with HCPCS A7034 mask.
- Frequency: 2 per month.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Oral interface used with PAP, each (HCPCS A7044):

- Frequency: 1 per 6 months.
- No other PAP mask ordered (i.e., HCPCS A7027, A7030, or A7034).
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Headgear used with PAP, each (HCPCS A7035):

- Frequency: 1 per 6 months.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Chinstrap used with PAP, each (HCPCS A7036):

- Frequency: 1 per 6 months.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Positive airway pressure tubing

Tubing with integrated heating element for use with PAP devices, each (HCPCS A4604):

- Frequency: 1 per 3 months.
- No other PAP tubing ordered (i.e., HCPCS A7037).
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Tubing used with PAP devices, each (HCPCS A7037):

- Frequency: 1 per 3 months.
- No other PAP tubing ordered (i.e., HCPCS A4604).
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Positive airway pressure device filters

Filter, disposable, used with PAP devices (HCPCS A7038):

- Frequency: 2 per 1 month.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Filter, non-disposable, used with PAP devices (HCPCS A7039):

- Frequency: 1 per 6 months.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Miscellaneous positive airway pressure supplies

Exhalation port with or without swivel used with accessories for positive airway devices, replacement only (HCPCS A7045):

- Frequency: 1 per 6 months.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Water chamber for humidifier, used with positive airway pressure device, replacement, each (HCPCS A7046):

- Frequency: 1 per 6 months.
- Provided data meets compliance criteria (device used ≥ 4 hours per night on 70% of nights), with 30 continuous days of compliance data provided from the time period within 6 months prior to the date of request

Oral Appliances for the Treatment of Obstructive Sleep Apnea (SL-9)

SL.TX.108.A

v1.0.2025

Coding

The inclusion of any code in this table does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

Treatment codes for oral appliances

Treatment Description	HCPCS
Oral device/appliance used to reduce upper airway collapsibility, adjustable or non-adjustable, prefabricated, includes fitting and adjustment	E0485
Oral device/appliance used to reduce upper airway collapsibility, adjustable or non-adjustable, custom fabricated, includes fitting and adjustment	E0486
Non-covered item or service (Used for oral appliances that do not incorporate all of the criteria as set forth in the Policy Article; tongue-retaining or tongue-positioning devices; and devices that are used only to treat snoring without a diagnosis of obstructive sleep apnea)	A9270
Oral device/appliance used to reduce upper airway collapsibility, without fixed mechanical hinge, custom fabricated, includes fitting and adjustment	K1027

Custom-fit oral appliances (SL-9.1)

Custom-fit oral appliances - indications (SL-9.1.2)

Custom fit oral appliances are medically necessary when **ALL** of the following criteria are met:

- A positive diagnosis of obstructive sleep apnea on a covered sleep study as demonstrated by **ONE** of the following:
 - AHI, RDI, or REI ≥ 5 and < 15 events per hour over the duration of the sleep test and documentation of:
 - Excessive daytime sleepiness, impaired cognition, mood disorders, or insomnia, or
 - Hypertension, ischemic heart disease, or history of stroke;
 - AHI, RDI, or REI ≥ 15 per hour over the duration of the sleep test
- Documentation of:
 - Intolerance or lack of benefit after a minimum of a one-month trial of PAP, **OR**
 - PAP is contraindicated for the individual as documented by the treating physician, **OR**
 - Individual prefers alternative treatment to CPAP (after a discussion of treatment options with the treating physician) **AND** AHI, RDI, or REI is < 30 .
- The device is ordered by the treating physician following a face to face visit and review of sleep study results
- A qualified licensed dentist (DDS and DMD) provides a custom device and follow-up to assess for dental-related side effects.

Note:

Oral devices to prevent temporomandibular joint (TMJ) disorders are considered experimental, investigational, or unproven (EIU).

Replacement custom fit oral appliances (SL-9.1.3)

Custom fit oral appliances can be replaced when **ALL** of the following criteria are met:

- Device is being used consistently with continued resolution of symptoms
- The device is ordered by the treating physician following a face to face visit
- A qualified licensed dentist (DDS and DMD) provides a custom device and follow-up to assess for dental-related side effects.
- **ONE** of the following applies
 - Device has been lost or irreparably damaged due to a specific accident, natural disaster or breakdown of device from regular use
 - Device is greater than 5 years old

Background and Supporting Information

- Continuous positive airway pressure is the gold standard for treatment of obstructive sleep apnea. Oral appliances are an alternative treatment option for individuals who are intolerant to PAP therapy or who prefer an alternative to CPAP. Subjective adherence and side effect profile are improved with oral appliances compared to

CPAP. However, CPAP results in a greater reduction in respiratory events (AHI, RDI or REI) and greater improvement in oxygen saturation. Oral appliances significantly reduce apnea hypopnea index regardless of severity of obstructive sleep apnea, although individuals with moderate to severe OSA are more likely to achieve their target AHI with CPAP compared to the oral appliance. Both oral appliances and CPAP improve excessive daytime sleepiness, quality of life, and cognitive performance.

- The AASM task force indicates that use of oral appliances in individuals with severe obstructive sleep apnea should be reserved for clinical scenarios where CPAP is not tolerated or does not provide benefit.
- The most common oral appliance utilized for the treatment of obstructive sleep apnea is the mandibular advancement device. There was insufficient evidence for the AASM task force to assess the efficacy of tongue retaining devices, which are also less well tolerated. Custom-made mandibular advancement devices are more effective for symptom improvement, compliance and tolerance compared to ready-made appliances.

Pediatric Oral Appliances (SL-9.2)

- Oral appliances may be considered medically necessary in the treatment of children with craniofacial anomalies with signs and symptoms of OSA
- Oral appliances are considered EIU for the treatment of OSA in children not meeting the above criteria

Pediatric Guidelines

Guideline	Page
Pediatric Sleep Guidelines (SL-3).....	76

Pediatric Sleep Guidelines (SL-3)

SL.ST.103.A

v1.0.2025

Polysomnography in pediatrics (SL-3.1)

General information

Use of home/portable sleep studies for the diagnosis of OSA in children (17 years and younger) is considered experimental, investigational, or unproven at this time. Limited portable studies, or studies in the home, are not sufficient to exclude OSA in a child with suggestive symptoms, nor can they reliably assess the severity of the disorder due to lack of CO₂ monitoring and arousal monitoring among other factors.

Pediatric definitions

Pediatric definitions for apneas and hypopneas differ compared with adults

Pediatric apnea Drop in peak signal excursion by $\geq 90\%$ of pre-event baseline with an oronasal thermal sensor or alternative apnea sensor (diagnostic) or PAP device flow (titration study) for at least the duration of 2 breaths during the baseline portion of the study (obstructive or mixed events). **Note:** Duration criteria differ for central events.

Pediatric hypopnea The peak signal excursion drops by $>30\%$ of the pre-event baseline using nasal pressure or alternative hypopnea sensor (diagnostic study) or PAP device flow (titration study) for ≥ 2 breaths.

There is a $>3\%$ oxygen desaturation from pre-event baseline or the event is associated with an arousal.

Improper uses of polysomnography in pediatrics

- The peer-reviewed medical literature **does not** support the following:
 - Repeat polysomnography in the follow-up of individuals with obstructive sleep apnea treated with CPAP when symptoms attributable to sleep apnea have resolved
 - Polysomnography in children for **ANY** of the following:

- Uncomplicated typical disorders of arousal (e.g.: sleep terrors, sleep walking), nightmares, sleepwalking, or bruxism in the absence of concern for safety, sleep disordered breathing, or nocturnal seizures
- Routine evaluation of adenotonsillar hypertrophy alone without other clinical signs or symptoms suggestive of obstructive sleep disordered breathing
- Routine follow-up for children with mild OSA whose symptoms have resolved post-adenotonsillectomy

Proper uses of polysomnography in pediatrics

Initial polysomnography for diagnosis

In lab full night diagnostic sleep study (**See Polysomnography (facility-based-PSG) - coding (SL-1.4.2)** for age specific codes) is appropriate for children (17 years of age and younger) for the diagnosis of the following conditions:

- Sleep related breathing disorders, such as obstructive sleep apnea, upper airway resistance syndrome or concern for sleep disordered breathing as evidenced by symptoms which may include:
 - Snoring
 - Restless or disturbed sleep
 - Behavioral disturbance, or learning disorders including deterioration in academic performance, hyperactivity, or attention deficit disorder
 - Unexplained enuresis
 - Frequent awakenings
 - Witnessed apnea
 - Labored breathing during sleep
 - Headaches on awakening
 - Hypertension
 - Sleep enuresis in children older than 5 years old
 - Daytime sleepiness, tiredness, fatigue, or mood disturbances unexplained by other conditions or etiologies
 - Polycythemia unexplained by other conditions or etiologies
 - Cor pulmonale unexplained by other conditions or etiologies
 - Documentation of **ONE** of the following, with provider concern that finding is related to obstructive sleep apnea:
 - Failure to thrive or growth impairment
 - Underweight or Overweight
 - Tonsillar hypertrophy with symptoms of OSA (any of the symptoms listed above)
 - Adenoidal facies with symptoms of OSA (any of the symptoms listed above)

- While limited data exists on the performance of in lab split-night testing (**See Polysomnography (facility-based-PSG) - coding (SL-1.4.2) for age specific codes**) in the pediatric population, recent data has demonstrated the feasibility of split night testing in children. Split night testing may be appropriate in the pediatric population at the discretion of the treating provider.

Initial polysomnography for positive airway pressure (PAP) titration

In lab full night PAP titration (**See Polysomnography (facility-based-PSG) - coding (SL-1.4.2) for age specific codes**) in children with obstructive sleep apnea syndrome is considered medically necessary in the following circumstances:

- Diagnosis of obstructive sleep apnea
 - Polysomnographic normal standards differ between children and adults. Diagnosis of pediatric obstructive sleep apnea is demonstrated by both of the following:
 - The presence of **ONE OR MORE** of the following
 - Snoring
 - Labored, paradoxical or obstructed breathing during the child's sleep
 - Sleepiness, hyperactivity, behavioral problems, or learning problems
 - PSG demonstrates **ONE OR MORE** of the following:
 - One or more obstructive apneas, mixed apneas, or hypopneas per hour of sleep
 - A pattern of obstructive hypoventilation defined as at least 25% of total sleep time with hypercapnia ($\text{PaCO}_2 > 50 \text{ mm Hg}$) in association with one or more of the following: snoring, flattening of the inspiratory nasal pressure waveform, paradoxical thoracoabdominal motion

Repeat polysomnography for diagnosis

- Repeat in lab full night diagnostic sleep study (**See Polysomnography (facility-based-PSG) - coding (SL-1.4.2) for age specific codes**) to assess for OSA is medically necessary in **ANY** of the following circumstances:
 - If obesity was a major contributing factor and significant weight loss has been achieved, repeat testing may be indicated to determine the need for continued therapy
 - A child with previously diagnosed obstructive sleep apnea treated with non PAP therapy (e.g. medication management, watchful waiting, oral appliance, rapid maxillary expansion) who has ongoing symptoms (such as snoring) despite treatment
 - Initial polysomnography is inadequate or non-diagnostic and the accompanying caregiver reports that the child's sleep and breathing patterns during the testing were not representative of the child's sleep at home

- Residual obstructive sleep apnea post-adenotonsillectomy when **ONE** of the following criteria is met:
 - Residual symptoms of OSA are present with mild OSA preoperatively
 - Moderate to severe OSA
 - Obesity
 - Craniofacial abnormalities that obstruct the upper airway
 - Neurological disorders such as Down Syndrome, Prader-Willi, and meningocele

Special scenarios

In lab full night diagnostic sleep study and in lab full night PAP titration (**See Polysomnography (facility-based-PSG) - coding (SL-1.4.2) for age specific codes**) are considered medically necessary in **ANY** of the following circumstances:

- A pediatric diagnostic sleep study can be performed as a screening for OSA in children with Down syndrome even without symptoms of obstructive sleep apnea
- Congenital central alveolar hypoventilation syndrome or sleep related hypoventilation due to neuromuscular disorders or chest wall deformities
- Atypical, frequent, or potentially injurious parasomnias
- Differentiate a parasomnia from sleep-related epilepsy when the initial clinical evaluation and standard EEG are inconclusive
- A pediatric positive airway pressure titration study can be performed without a baseline sleep study for sleep-related hypoventilation under the following circumstances:
 - Documented neuromuscular disease such as Duchenne muscular dystrophy or spinal muscular atrophy
 - Individuals being discharged from the hospital determined to have sleep-related hypoventilation during hospitalization
- Polysomnography when there is clinical evidence of a sleep related breathing disorder in infants who have experienced an Apparent Life-Threatening Event (ALTE) or a Brief Resolved Unexplained Event (BRUE)
- Primary central sleep apnea of infancy (when not explained by another sleep disorder, medical disorder, or medications)
- PSG as part of the evaluation prior to decannulation for children treated with tracheostomy for sleep related breathing disorders
- Periodic evaluation with polysomnography to adjust ventilator settings
- Suspected nocturnal seizure activity

- Suspected periodic limb movement disorder (when other medical disorders have been ruled out)

Repeat polysomnography for positive airway pressure (PAP) titration

Repeat in lab full night PAP titration (**See Polysomnography (facility-based-PSG) - coding (SL-1.4.2) for age specific codes**) is medically necessary to periodically re-evaluate the appropriateness of continuous positive airway pressure (CPAP) setting based on the child's growth pattern or the presence of recurrent symptoms while on PAP.

Note:

In lab full night diagnostic sleep study should be performed in conjunction with an MSLT for diagnosis of narcolepsy or idiopathic hypersomnia. See **PSG and Multiple Sleep Latency testing (excessive sleepiness)(SL-2.3)**

CPAP in pediatrics (SL-3.2)

- CPAP is medically necessary when **ALL** of the following criteria are met:
 - OSA diagnosis has been established by PSG; **AND**
 - Adenotonsillectomy has been unsuccessful or is determined to be clinically inappropriate, or when definitive surgery is indicated but must await complete dental and facial development

Program Exclusions

Guideline	Page
Sleep Apnea Treatment Program Exclusions (SL-5).....	82

Sleep Apnea Treatment Program Exclusions (SL-5)

SL.TXS.105.A

v1.0.2025

Experimental, investigational, or unproven (SL-5.1)

- Certain therapies may be considered experimental, investigational, or unproven if there is **ANY** of the following:
 - A paucity of supporting evidence in the peer reviewed literature
 - The evidence has not matured to exhibit improved health parameters
 - The therapy lacks a collective opinion of support

Note:

The list below is not comprehensive

- The effectiveness of the following therapies has not been established in the treatment of OSA; these therapies, as well as other therapies not addressed in these guidelines, may be considered experimental, investigational, or unproven:
 - Bongo Rx
 - ULTepap
 - iNAP
 - eXciteOSA (HCPCS E0492 and E0493)
 - Somnera
 - MATRx oral appliance test
 - Electronic positional obstructive sleep apnea treatment with sensor (HCPCS E0530)
- SleepTesting exclusions:
 - Actigraphy (CPT[®] 95803)
 - Actigraph devices, worn on the wrist, record movement and utilize rest activity patterns to estimate sleep parameters.
 - While actigraphy is performed as part of certain home sleep apnea testing devices, actigraphy performed as a stand-alone study is considered not medically necessary.

Durable medical equipment device and supply exclusions (SL-5.2)

- CPT® 94799 – Unlisted pulmonary service or procedure
 - Due to the presence of more specific codes, medical necessity for this code cannot be established
- HCPCS E1399 – Miscellaneous durable medical equipment items, components, and accessories
 - Due to the presence of more specific codes for PAP equipment, and supplies needing scheduled replacement, E1399 cannot be approved for routinely replaced PAP machine and parts. E1399 may be approved for parts required in CPAP repair when the DME supplier specifies what needs to be repaired and what replacement parts (ie, blower motor) are needed. Note that K0739 and K0740 which refer to labor costs and rental costs during the repair period do not require approval

PAP-NAP (SL-2.7)**CPT® 95807-52**

PAP-NAP, a daytime abbreviated cardiorespiratory sleep study, was developed as a means of improving adherence to positive airway pressure in patients with sleep disordered breathing and co-morbid insomnia and psychiatric disorders. A pilot study performed in 2008 demonstrated improvement in PAP adherence compared to historical controls in patients with insomnia and diagnosed and/or symptoms of psychiatric disorders. However, no subsequent controlled studies have been published. Therefore, CPT® 95807 or 95807-52 for the purposes of performing PAP-NAP is not covered procedure

Note:

Result of previous studies should be submitted for review prior to authorization of additional studies

References

Guideline	Page
General References.....	85
Pediatric References.....	95
Questionnaires (SL-8).....	97

General References

SL.AD.100.A

v1.0.2025

1. Acelajado MC, Hughes ZH, Oparil S, Calhoun DA. Treatment of Resistant and Refractory Hypertension. *Circ Res*. 2019;124(7):1061-1070. doi:10.1161/CIRCRESAHA.118.312156
2. Ahmad A, Ahmad R, Meteb M, et al. The relationship between opioid use and obstructive sleep apnea: A systematic review and meta-analysis. *Sleep Med Rev*. 2021;58:101441. doi:10.1016/j.smrv.2021.101441.
3. Akintomide GS, Rickards H. Narcolepsy: a review. *Neuropsychiatric disease and treatment*. 2011;7:507-518. doi:10.2147/NDT.S23624
4. American Academy of Sleep Medicine. International Classification of Sleep Disorders. 3rd ed, text revision. Darien, IL: American Academy of Sleep Medicine; 2023.
5. Agrawal R, Sharafkhaneh A, Gottlieb DJ, et al. Mortality Patterns Associated with Central Sleep Apnea among Veterans: A Large, Retrospective, Longitudinal Report. *Ann Am Thorac Soc*. 2023 Mar;20(3):450-455. doi:10.1513/AnnalsATS.202207-648OC.
6. Alrubasy WA, Abuawwad MT, Taha MJJ, et al. Hypoglossal nerve stimulation for obstructive sleep apnea in adults: An updated systematic review and meta-analysis. *Respir Med*. Published online October 12, 2024. doi:10.1016/j.rmed.2024.107826
7. Ancoli-Israel S, Gehrman P, Kripke DF, et al. Long-term follow-up of sleep disordered breathing in older adults. *Sleep Med*. 2001;2(6):511-516. doi:10.1016/s1389-9457(00)00096-4
8. Arellano-Maric MP, Hamm C, Duiverman ML, et al. Obesity hypoventilation syndrome treated with non-invasive ventilation: Is a switch to CPAP therapy feasible? *Respirology*. 2020 Apr;25(4):435-442. doi: 10.1111/resp.13704.
9. Arzt M, Floras JS, Logan AG, et al. Suppression of Central Sleep Apnea by Continuous Positive Airway Pressure and Transplant-Free Survival in Heart Failure: A Post Hoc Analysis of the Canadian Continuous Positive Airway Pressure for Patients with Central Sleep Apnea and Heart Failure Trial (CANPAP). *Circulation*. 2007;115(25):3173-3180. doi:10.1161/circulationaha.106.683482.
10. Arzt M, Young T, Peppard PE, et al. Dissociation of obstructive sleep apnea from hypersomnolence and obesity in patients with stroke. *Stroke*. 2010;41(3):e129-e134. doi:10.1161/STROKEAHA.109.566463
11. Askland K, Wright L, Wozniak DR, et al. Educational, supportive and behavioural interventions to improve usage of continuous positive airway pressure machines in adults with obstructive sleep apnoea. *Cochrane Database Syst Rev*. 2020;4(4):CD007736. Published 2020 Apr 7. doi:10.1002/14651858.CD007736.pub3
12. Aurora RN, Casey KR, Kristo D, et al. Practice Parameters for the Surgical Modifications of the Upper Airway for Obstructive Sleep Apnea in Adults. *Sleep*. 2010;33(10):1408-1413. doi:10.1093/sleep/33.10.1408.
13. Aurora RN, Bista SR, Casey KR, et al. Updated Adaptive Servo-Ventilation Recommendations for the 2012 AASM Guideline: "The Treatment of Central Sleep Apnea Syndromes in Adults: Practice Parameters with an Evidence-Based Literature Review and Meta-Analyses." *J Clin Sleep Med*. 2016;12(05):757-761. doi:10.5664/jcsm.5812.
14. Aurora RN, Chowdhuri S, Ramar K, et al. The treatment of central sleep apnea syndromes in adults: practice parameters with an evidence-based literature review and meta-analyses. *Sleep*. 2012;35(1):17-40. Published 2012 Jan 1. doi:10.5665/sleep.1580
15. Morgenthaler TI, Auerbach S, Casey KR, et al. Position Paper for the Treatment of Nightmare Disorder in Adults: An American Academy of Sleep Medicine Position Paper. *J Clin Sleep Med*. 2018;14(6):1041-1055. doi:10.5664/jcsm.7178,
16. Aurora RN, Collop NA, Jacobowitz O, et al. Quality Measures for the Care of Adult Patients with Obstructive Sleep Apnea. *J Clin Sleep Med*. January 2015. doi:10.5664/jcsm.4556.
17. Aurora RN, Kristo DA, Bista SR, et al. The Treatment of Restless Legs Syndrome and Periodic Limb Movement Disorder in Adults—An Update for 2012: Practice Parameters with an Evidence-Based Systematic Review and Meta-Analyses. *Sleep*. 2012;35(8):1039-1062. doi:10.5665/sleep.1988.
18. Aurora RN, Kristo DA, Bista SR, et al. Update to the AASM Clinical Practice Guideline: "The Treatment of Restless Legs Syndrome and Periodic Limb Movement Disorder in Adults—an Update for 2012: Practice

- Parameters with an Evidence-Based Systematic Review and Meta-Analyses." *Sleep*. 2012;35(8):1037-1037. doi:10.5665/sleep.1986.
19. Aurora RN, Zak RS, Auerbach SH, et al. Best practice guide for the treatment of nightmare disorder in adults. *J Clin Sleep Med*. 2010;6(4):389#401.
 20. Baptista PM, Costantino A, Moffa A, Rinaldi V, Casale M. Hypoglossal Nerve Stimulation in the Treatment of Obstructive Sleep Apnea: Patient Selection and New Perspectives. *Nat Sci Sleep*. 2020 Feb 13;12:151-159. doi: 10.2147/NSS.S221542.
 21. Bassetti C, Aldrich MS, Chervin RD, Quint D. Sleep apnea in patients with transient ischemic attack and stroke: a prospective study of 59 patients. *Neurology*. 1996;47(5):1167-1173. doi:10.1212/wnl.47.5.1167
 22. Benbadis SR. Association between the Epworth Sleepiness Scale and the Multiple Sleep Latency Test in a Clinical Population. *Ann Intern Med*. 1999;130(41):289-292. Doi:10.7326/0003-4819-130-4-199902160-00014.
 23. Bernhardt J, Hayward KS, Kwakkel G, et al. Agreed definitions and a shared vision for new standards in stroke recovery research: The Stroke Recovery and Rehabilitation Roundtable taskforce. *Int J Stroke*. 2017;12(5):444-450. doi:10.1177/1747493017711816
 24. Berry RB, Hill G, Thompson L, McLaurin V. Portable monitoring and autotitration versus polysomnography for the diagnosis and treatment of sleep apnea. *Sleep*. 2008;31(10):1423#1431.
 25. Bosschieter PFN, de Vries N, Mehra R, et al. Similar effect of hypoglossal nerve stimulation for obstructive sleep apnea in 5 disease severity categories. *J Clin Sleep Med*. 2022;18(6):1657–1665 .
 26. Bosschieter P, DeVries N, Vanderveken O. Comparison of upper airway stimulation outcomes in star and adhere sub-populations: narrow ahi range vs broad ahi range. *Sleep*, Volume 46, Supplement 1, 2023.
 27. Berry RB, Troester MM, Quan SF, et al. for the American Academy of Sleep Medicine. The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications, Version 3, Darien, IL: American Academy of Sleep Medicine; 2023.
 28. Berry RB, Chediak A, Brown LK, et al. Best clinical practices for the sleep center adjustment of noninvasive positive pressure ventilation (NPPV) in stable chronic alveolar hypoventilation syndromes. *J Clin Sleep Med*. 2010;6(5):491-509.
 29. Brookes CCD, Boyd SB. Controversies in Obstructive Sleep Apnea Surgery. *Oral and Maxillofacial Surgery Clinics of North America*. 2017;29(4):503-513. doi:10.1016/j.coms.2017.07.005.
 30. Bull M and the Committee on genetics. Clinical Report – Health Supervision for Children with Down Syndrome. *Pediatrics* 2011;128:393–406.
 31. Caples SM, Anderson WM, Calero K, Howell M, Hashmi SD. Use of polysomnography and home sleep apnea tests for the longitudinal management of obstructive sleep apnea in adults: an American Academy of Sleep Medicine clinical guidance statement. *J Clin Sleep Med*. 2021;17(6):1287-1293. doi:10.5664/jcsm.9240
 32. CGS Administrators, LLC. Hypoglossal Nerve Stimulation for the Treatment of Obstructive Sleep Apnea Local Coverage Determination (LCD) L38307. Nashville, TN: CGS Administrators, LLC;4/1/20. Available at:www.cgsmedicare.com.
 33. CGS Administrators, LLC. Respiratory Assist Devices Local Coverage Determination (LCD) L33800 Nashville, TN: CGS Administrators, LLC;08/08/2021. Available at:www.cgsmedicare.com.
 34. Centers for Medicare & Medicaid Services' (CMS') Healthcare Common Procedure Coding System (HCPCS) Level II Final Coding, Benefit Category and Payment Determinations (Appendix A: HCPCS Level II Temporary ("K" codes) Coding Revision on pg 110 and 111).
 35. Chang JL, Goldberg AN, Alt JA, et al. International Consensus Statement on Obstructive Sleep Apnea. *Int Forum Allergy Rhinol*. 2023 Jul;13(7):1061-1482. doi: 10.1002/alr.23079.
 36. Chervin, R.D. (2011). Chapter 59 Use of Clinical Tools and Tests in Sleep Medicine. In M.H. Kryger, T. Roth, & W.C. Dement (Eds.), *Principles and practice of sleep medicine*, 5th ed. (pp. 675-676). Elsevier Saunders.
 37. Choi JH, Cho SH, Kim S-N, Suh JD, Cho JH. Predicting Outcomes after Uvulopalatopharyngoplasty for Adult Obstructive Sleep Apnea. *Otolaryngol Head Neck Surg*. 2016;155(6):904-913. doi:10.1177/0194599816661481.
 38. Chung F, Yegneswaran B, Liao P, et al. STOP Questionnaire. *Anesthesiology*. 2008;108(5):812-821. doi:10.1097/aln.0b013e31816d83e4.
 39. Collop NA, Anderson WM, Boehlecke B, et al. Clinical Guidelines for the Use of Unattended Portable Monitors in the Diagnosis of Obstructive Sleep Apnea in Adult Patients. *J Clin Sleep Med*. 2007;03(07):737-747. doi:10.5664/jcsm.27032.

40. Coelho, Frenando MS. Benefit of repeat multiple sleep latency testing in confirming a possible narcolepsy diagnosis. *Journal of Clinical Neurophysiology* 28(4):p 412-414, August 2011. | DOI: 10.1097/WNP.0b013e31822734a3.
41. Costanzo MR, Augostini R, Goldberg LR, Ponikowski P, Stellbrink C, Javaheri S. Design of the remedē System Pivotal Trial: A Prospective, Randomized Study in the Use of Respiratory Rhythm Management to Treat Central Sleep Apnea. *J Card Fail.* 2015;21(11):892-902. doi:10.1016/j.cardfail.2015.08.344
42. Costanzo MR, Ponikowski P, Javaheri S, et al. Transvenous neurostimulation for central sleep apnoea: a randomised controlled trial. *Lancet.* 2016;388(10048):974-982. doi:10.1016/S0140-6736(16)30961-8
43. Costanzo MR, Ponikowski P, Coats A, et al. Phrenic nerve stimulation to treat patients with central sleep apnoea and heart failure. *Eur J Heart Fail.* 2018;20(12):1746-1754. doi:10.1002/ehf.1312
44. Costanzo MR, Javaheri S, Ponikowski P, et al. Transvenous Phrenic Nerve Stimulation for Treatment of Central Sleep Apnea: Five-Year Safety and Efficacy Outcomes. *Nat Sci Sleep.* 2021;13:515-526. Published 2021 Apr 29. doi:10.2147/NSS.S300713
45. Costanzo MR, Ponikowski P, Javaheri S, et al. Sustained 12 Month Benefit of Phrenic Nerve Stimulation for Central Sleep Apnea. *Am J Cardiol.* 2018;121(11):1400-1408. doi:10.1016/j.amjcard.2018.02.022
46. Cowie MR, Woehrle H, Wegscheider K, et al. Adaptive Servo-Ventilation for Central Sleep Apnea in Systolic Heart Failure. *N Engl J Med.* 2015;373(12):1095-1105. doi:10.1056/nejmoa1506459.
47. Consensus Conference Panel, Watson NF, Badr MS, et al. Recommended Amount of Sleep for a Healthy Adult: A Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society. *J Clin Sleep Med.* 2015;11(6):591-592. Published 2015 Jun 15. doi:10.5664/jcsm.4758
48. Consensus Conference Panel, Watson NF, Badr MS, et al. Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society on the Recommended Amount of Sleep for a Healthy Adult: Methodology and Discussion. *J Clin Sleep Med.* 2015;11(8):931-952. Published 2015 Aug 15. doi:10.5664/jcsm.4950
49. Drakatos P, Leschziner G. Cataplexy with normal sleep studies and normal csf hypocretin: an explanation? *J Clin Sleep Med* 2016;12(3):449–450. doi: 10.5664/jcsm.5604.
50. El-Solh AA, Rudraraju A, Pasrija D, Bui H. Pharmacotherapy of residual excessive sleepiness among continuous positive airway pressure (CPAP) treated patients with sleep apnea. *Expert Opin Pharmacother.* 2022;23(4):507-516. doi:10.1080/14656566.2022.2029408
51. Epstein LJ, Kristo D, Strollo PJ Jr, et al. Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *J Clin Sleep Med.* 2009;5(3):263-276.
52. FDA Approval letter 4/14/20 (age 18-21 added): https://www.accessdata.fda.gov/cdrh_docs/pdf13/P130008S039A.pdf.
53. FDA Approval letter 3/20/23 (Adol Down): <https://www.fda.gov/medical-devices/recently-approved-devices/inspire-upper-airway-stimulation-p130008s089>.
54. FDA Approval letter 6/8/23 (adult BMI/AHI expansion): https://www.accessdata.fda.gov/cdrh_docs/pdf13/P130008S090A.pdf.
55. Filiatrault M-L, Chauny J-M, Daoust R, Roy M-P, Denis R, Lavigne G. Medium Increased Risk for Central Sleep Apnea but Not Obstructive Sleep Apnea in Long-Term Opioid Users: A Systematic Review and Meta-Analysis. *J Clin Sleep Med.* 2016;12(04):617-625. doi:10.5664/jcsm.5704.
56. Final MRB Task 16-01 Letter Report from MCSAC and MRB | FMCSA. Dot.gov. Published 2017. <https://www.fmcsa.dot.gov/advisory-committees/mrb/final-mrb-task-16-01-letter-report-mcsac-and-mrb>
57. First Coast Service Options, INC. Hypoglossal Nerve Stimulation for the Treatment of Obstructive Sleep Apnea Local Coverage Determination (LCD) L38398. Jacksonville, FL:First Coast Service Options, Inc.;03/15/20. Available at:<https://medicare.fcso.com>.
58. Fox H, Oldenburg O, Javaheri S, et al. Long-term efficacy and safety of phrenic nerve stimulation for the treatment of central sleep apnea. *Sleep.* 2019;42(11):zsz158. doi:10.1093/sleep/zsz158
59. Fraga da Rosa JC, et al. Diagnostic accuracy of oximetry for obstructive sleep apnea: a study on older adults in a home setting. *Clinics (Sao Paulo).* 2021; 76: e3056. doi: 10.6061/clinics/2021/e3056.
60. Freeman CG, Durgham R, Ren E, et al. The Impact of Hypoglossal Nerve Stimulation on Secondary Health Outcomes. *Ear Nose Throat J.* Published online February 29, 2024. doi:10.1177/01455613241235538
61. Fudim M, Shahid I, Emani S, et al. Evaluation and Treatment of Central Sleep Apnea in Patients with Heart Failure. *Curr Probl Cardiol.* 2022 Dec;47(12):101364. doi:10.1016/j.cpcardiol.2022.101364.

62. Galiè N, Humbert M, Vachiery J-L, et al. 2015 ESC/ERS Guidelines for the diagnosis and treatment of pulmonary hypertension. *Eu Heart J*. 2015;37(1):67-119. doi:10.1093/eurheartj/ehv317.
63. Gay PC, Owens RL; ONMAP Technical Expert Panel. Executive Summary: Optimal Noninvasive Ventilation Medicare Access Promotion (ONMAP) [published online ahead of print, 2021 Jul 28]. *Chest*. 2021;S0012-3692(21)01481-1. doi:10.1016/j.chest.2021.05.074.
64. Georges M, Nguyen-Baranoff D, Griffon L, Foignot C, Bonniaud P, Camus P, Pepin JL, Rabec C. Usefulness of transcutaneous PCO₂ to assess nocturnal hypoventilation in restrictive lung disorders. *Respirology*. 2016 Oct;21(7):1300-6. doi: 10.1111/resp.12812.
65. Gillespie MB, Soose RJ, Woodson BT, et al. Upper Airway Stimulation for Obstructive Sleep Apnea: Patient-Reported Outcomes after 48 Months of Follow-up. *Otolaryngol Head Neck Surg*. 2017;156(4):765-771. doi:10.1177/0194599817691491.
66. Goldberg LR, Fox H, Stellbrink C, et al. Design of the remedē System Therapy (rēST) study: A prospective non-randomized post-market study collecting clinical data on safety and effectiveness of the remedē system for the treatment of central sleep apnea. *Sleep Med*. 2022;100:238-243. doi:10.1016/j.sleep.2022.08.026
67. Harmon EK, Stafford P, Ibrahim S, et al. Atrial fibrillation is associated with central sleep apnea in clinic patients undergoing diagnostic polysomnography. *J Arrhythm*. 2020;36(6):991-996. Published 2020 Sep 3. doi:10.1002/joa3.12427
68. Heiser C, Steffen A, Boon M, et al. Post-approval upper airway stimulation predictors of treatment effectiveness in the ADHERE registry. *Eur Respir J* 2019; 53: 1801405 [https://doi.org/10.1183/13993003.01405-2018].
69. Heiser C, Maurer JT, Hofauer B, Sommer JU, Seitz A, Steffen A. Outcomes of Upper Airway Stimulation for Obstructive Sleep Apnea in a Multicenter German Postmarket Study. *Otolaryngol Head Neck Surg*. 2017;156(2):378-384. doi:10.1177/0194599816683378
70. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines Circulation. 2022;145(18):e895-e1032. doi:10.1161/CIR.0000000000001063
71. Hill NS, Criner J, Branson RD, et al; ONMAP Technical Expert Panel. Optimal NIV Medicare Access Promotion: Patients with COPD: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest* 2021;160(5):e389-e397. DOI: 10.1016/j.chest.2021.06.082.
72. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's updated sleep duration recommendations: final report. *Sleep Health*. 2015;1(4):233-243. doi:10.1016/j.sleh.2015.10.004
73. Hirshkowitz M, Whiton K, Albert SM, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. 2015;1(1):40-43. doi:10.1016/j.sleh.2014.12.010
74. Hoepfer MM, Bogaard HJ, Condliffe R, et al. Definitions and Diagnosis of Pulmonary Hypertension. *J Am Coll Cardiol*. 2013;62(25). doi:10.1016/j.jacc.2013.10.032.
75. Huang YS, Guilleminault C, Lin CH, Chen CH, Chin WC, Chen TS. Multiple sleep latency test in narcolepsy type 1 and narcolepsy type 2: a 5-year follow-up study. *J Sleep Res*. 2018;27(5):e12700. https://doi.org/10.1111/jsr.12700.
76. Huhtakangas JK, Saaresranta T, Bloigu R, Huhtakangas J. The Evolution of Sleep Apnea Six Months After Acute Ischemic Stroke and Thrombolysis [published correction appears in J Clin Sleep Med. 2020 May 15;16(5):837-838. doi: 10.5664/jcsm.8428]. *J Clin Sleep Med*. 2018;14(12):2005-2011. Published 2018 Dec 15. doi:10.5664/jcsm.7524
77. Jacobs SS, Krishnan JA, Lederer DJ, GhazipuraM, Hossain T, Tan AM, et al. Home oxygen therapy for adults with chronic lung disease: an official American Thoracic Society clinical practice guideline. *Am J Respir Crit Care Med* 2020;202:e121–e141. doi: 10.1164/rccm.202009-3608ST.
78. Jacobs SS, Krishnan JA, Lederer DJ, GhazipuraM, Hossain T, Tan AM, et al. Erratum: home oxygen therapy for adults with chronic lung disease: an official American Thoracic Society clinical practice guideline. *Am J Respir Crit Care Med* 2020;203(8):1045-1046. doi: 10.1164/rccm.v203erratum7.
79. Javaheri S, Smith J, Chung E. The Prevalence and Natural History of Complex Sleep Apnea. *J Clin Sleep Med*. 2009;05(03):205-211. doi:10.5664/jcsm.27486.
80. Javaheri S. Update on Persistent Excessive Daytime Sleepiness in OSA. *Chest*. 2020;158(2):776-786. doi:10.1016/j.chest.2020.02.036
81. Javaheri S, Harris N, Howard J, Chung E. Adaptive servoventilation for treatment of opioid-associated central sleep apnea. *J Clin Sleep Med*. 2014;10(6):637-643. Published 2014 Jun 15. doi:10.5664/jcsm.3788

82. Javaheri S, McKane S. Transvenous phrenic nerve stimulation to treat idiopathic central sleep apnea. *J Clin Sleep Med*. 2020;16(12):2099-2107. doi:10.5664/jcsm.8802
83. Jensen MH, Dalggaard F, Rude Laub R, et al; DAN-APNO investigators. Prevalence of sleep apnea in unselected patients with atrial fibrillation by a home-monitoring device: The DAN-APNO study. *Int J Cardiol Heart Vasc*. 2023 May 18;47:101219. doi: 10.1016/j.ijcha.2023.101219.
84. Joglar JA, Chung MK, Armbruster AL, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2023 Nov 30. doi:10.1161/CIR.0000000000001193.
85. Johal A, Haria P, Manek S, Joury E, Riha R. Ready-Made Versus Custom-Made Mandibular Repositioning Devices in Sleep Apnea: A Randomized Clinical Trial. *J Clin Sleep Med*. 2017;13(02):175-182. doi:10.5664/jcsm.6440.
86. Johns MW. A New Method for Measuring Daytime Sleepiness: The Epworth Sleepiness Scale. *Sleep*. 1991;14(6):540-545. doi:10.1093/sleep/14.6.540.
87. Johnson KG, Sullivan SS, Nti A, Rastegar V, Gurubhagavatula I. The impact of the COVID-19 pandemic on sleep medicine practices. *J Clin Sleep Med*. 2021;17(1):79-87. doi:10.5664/jcsm.8830.
88. Kapur VK, Auckley DH, Chowdhuri S, et al. Clinical Practice Guideline for Diagnostic Testing for Adult Obstructive Sleep Apnea: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2017;13(03):479-504. doi:10.5664/jcsm.6506.
89. Kendzerska T, Gomes T, Malhotra A, et al. Association of obstructive sleep apnea and opioids use on adverse health outcomes: A population study of health administrative data. *PLoS One*. 2022 Jun 28;17(6):e0269112. doi:10.1371/journal.pone.0269112.
90. Kihara Y, Seino Y, Momomura S; SAVIOR-C investigators. Adaptive Servo-Ventilation for Central Sleep Apnea in Heart Failure. *N Engl J Med*. 2016;374(7):687-688. doi:10.1056/NEJMc1515007
91. Korenstein D, Chimonas S, Barrow B, Keyhani S, Troy A, Lipitz-Snyderman A. Development of a Conceptual Map of Negative Consequences for Patients of Overuse of Medical Tests and Treatments. *JAMA Intern Med*. 2018;178(10):1401-1407. doi:10.1001/jamainternmed.2018.3573
92. Krahn LE, Arand DL, Avidan AY, et al. Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in adults: guidance from the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2021 Dec 1;17(12):2489-2498. doi:10.5664/jcsm.9620. Erratum in: *J Clin Sleep Med*. 2022 Aug 1;18(8):2089.
93. Krakow B, Ulibarri V, Melendrez D, et al. A Daytime, Abbreviated Cardio-Respiratory Sleep Study (CPT 95807–52) To Acclimate Insomnia Patients with Sleep Disordered Breathing to Positive Airway Pressure (PAP-NAP). *J Clin Sleep Med*. 2008;04(03):212-222. doi:10.5664/jcsm.27183.
94. Kushida CA, Chediak A, Berry RB, et al. Clinical Guidelines for the Manual Titration of Positive Airway Pressure in Patients with Obstructive Sleep Apnea. *J Clin Sleep Med*. 2008;04(02):157-171. doi:10.5664/jcsm.27133.
95. Khayat R, Abraham W, Patt B, et al. Central sleep apnea is a predictor of cardiac readmission in hospitalized patients with systolic heart failure. *J Card Fail*. 2012;18(7):534-540. doi:10.1016/j.cardfail.2012.05.003
96. LCD - Positive Airway Pressure (PAP) Devices for the Treatment of Obstructive Sleep Apnea (L33718). Cms.gov. Published 2023. Accessed November 15, 2024. <https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?LCDId=33718&DocID=L33718>
97. Leschziner G. Narcolepsy: a clinical review. *Pract Neurol*. 2014;14(5):323-331. doi:10.1136/practneurol-2014-000837
98. Levendowski D, Steward D, Woodson BT, Olmstead R, Popovic D, Westbrook P. The impact of obstructive sleep apnea variability measured in-lab versus in-home on sample size calculations. *Int Arch Med*. 2009;2(1):2. Published 2009 Jan 2. doi:10.1186/1755-7682-2-2
99. Li L, Umbach DM, Li Y, et al. Sleep apnoea and hypoventilation in patients with five major types of muscular dystrophy. *BMJ Open Respir Res*. 2023 Apr;10(1):e001506. doi:10.1136/bmjresp-2022-001506.
100. Lin Y, Wu Y, Lin Q, et al. Objective Sleep Duration and All-Cause Mortality Among People With Obstructive Sleep Apnea. *JAMA Netw Open*. 2023;6(12):e2346085. Published 2023 Dec 1. doi:10.1001/jamanetworkopen.2023.46085
101. Littner MR, Kushida C, Wise M, et al. Practice parameters for clinical use of the multiple sleep latency test and the maintenance of wakefulness test. *Sleep*. 2005;28(1):113-121. doi:10.1093/sleep/28.1.113

102. Liu SY, Riley RW, Yu MS. Surgical Algorithm for Obstructive Sleep Apnea: An Update. *Clin Exp Otorhinolaryngol*. 2020;13(3):215-224. doi:10.21053/ceo.2020.01053
103. Liu X, Lam DC, Chan KPF, Chan HY, Ip MS, Lau KK. Prevalence and Determinants of Sleep Apnea in Patients with Stroke: A Meta-Analysis. *J Stroke Cerebrovasc Dis*. 2021;30(12):106129. doi:10.1016/j.jstrokecerebrovasdis.2021.106129
104. Lopez R, Doukkali A, Barateau L, et al.. Test-retest reliability of the Multiple Sleep Latency Test in central disorders of hypersomnolence. *Sleep*. 2017;40(12):zsx164. DOI: 10.1093/sleep/zsx164.
105. Macavei VM, Spurling KJ, Loft J, Makker HK. Diagnostic predictors of obesity-hypoventilation syndrome in patients suspected of having sleep disordered breathing. *J Clin Sleep Med*. 2013;9(9):879-884. doi:10.5664/jcsm.2986
106. Malhotra RK, Kirsch DB, Kristo DA, et al; American Academy of Sleep Medicine Board of Directors. Polysomnography for Obstructive Sleep Apnea Should Include Arousal-Based Scoring: An American Academy of Sleep Medicine Position Statement. *J Clin Sleep Med*. 2018 Jul 15;14(7):1245-1247. doi: 10.5664/jcsm.7234.
107. Malhotra A, Bertisch S, Wellman A. Complex sleep apnea: it isn't really a disease. *J Clin Sleep Med*. 2008;4(5):406-408.
108. Marklund M, Carlberg B, Forsgren L, et al. Oral Appliance Therapy in Patients With Daytime Sleepiness and Snoring or Mild to Moderate Sleep Apnea: A Randomized Clinical Trial. *JAMA Intern Med*. 2015 Aug;175(8):1278-85. doi:10.1001/jamainternmed.2015.2051.
109. Martin JL, Hakim AD. Wrist Actigraphy. *Chest*. 2011;139(6):1514-1527. doi:10.1378/chest.10-1872.
110. Martínez-García MA, Capote F, Campos-Rodríguez F, et al. Effect of CPAP on blood pressure in patients with obstructive sleep apnea and resistant hypertension: the HIPARCO randomized clinical trial. *JAMA*. 2013;310(22):2407-2415. doi:10.1001/jama.2013.281250
111. Maski K, Trotti LM, Kotagal S, et al. Treatment of central disorders of hypersomnolence: an American Academy of Sleep Medicine clinical practice guideline. *J Clin Sleep Med*. 2021;17(9):1881-1893. doi:10.5664/jcsm.9328
112. Maski KP, Amos LB, Carter JC, Koch EE, Kazmi U, Rosen CL. Recommended protocols for the Multiple Sleep Latency Test and Maintenance of Wakefulness Test in children: guidance from the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2024;20(4):631-641. doi:10.5664/jcsm.10974
113. May AM, Gharibeh T, Wang L, et al. CPAP Adherence Predictors in a Randomized Trial of Moderate-to-Severe OSA Enriched With Women and Minorities. *Chest*. 2018 Sep;154(3):567-578. doi: 10.1016/j.chest.2018.04.010.
114. McEvoy RD, Antic NA, Heeley E, et al. CPAP for Prevention of Cardiovascular Events in Obstructive Sleep Apnea. *N Engl J Med*. 2016;375(10):919-931. doi:10.1056/NEJMoa1606599
115. Medical Examiner's Handbook 2024 Edition. FMCSA. Dot.gov. Published 2024. <https://www.fmcsa.dot.gov/regulations/medical/driver-medical-requirements/medical-examiners-handbook-2024-edition>
116. Meira, L., Zeller, M.V., Eusébio, E.S., Clara, E.S., Viana, P., & Drummond, M. (2017). Maintenance of Wakefulness Test in clinical practice. *ERJ Open Research*, 3, 5.
117. Miller CM, Khanna A, Strohl KP. Assessment and policy for commercial driver license referrals. *J Clin Sleep Med*. 2007;3(4):417-423
118. Mokhlesi B, Masa JF, Brozek JL, et al. Evaluation and Management of Obesity Hypoventilation Syndrome. An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med*. 2019;200(3). doi:10.1164/rccm.201905-1071.
119. Mokhlesi B, Won CH, Make BJ, et al; ONMAP Technical Expert Panel. Optimal NIV Medicare Access Promotion: Patients With Hypoventilation Syndromes: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. 2021 Nov;160(5):e377-e387. doi:10.1016/j.chest.2021.06.083 .
120. Montesi SB, Edwards BA, Malhotra A, Bakker JP. The effect of continuous positive airway pressure treatment on blood pressure: a systematic review and meta-analysis of randomized controlled trials. *J Clin Sleep Med*. 2012;8(5):587-596. Published 2012 Oct 15. doi:10.5664/jcsm.2170
121. Morgenthaler TI, Malhotra A, Berry RB, et al. ONMAP Technical Expert Panel. Optimal NIV Medicare Access Promotion: Patients With Central Sleep Apnea: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. 2021 Nov;160(5):e419-e425. doi:10.1016/j.chest.2021.07.039.
122. Moser RJ 3rd, Rajagopal KR. Obstructive sleep apnea in adults with tonsillar hypertrophy. *Arch Intern Med*. 1987;147(7):1265-1267.

123. Murphy PB, Rehal S, Arbane G, et al. Effect of Home Noninvasive Ventilation With Oxygen Therapy vs Oxygen Therapy Alone on Hospital Readmission or Death After an Acute COPD Exacerbation: A Randomized Clinical Trial. *JAMA*. 2017 Jun 6;317(21):2177-2186. doi:10.1001/jama.2017.4451
124. Narcolepsylink.com. Published 2017. <https://www.narcolepsylink.com/pediatric-patients/screening/>
125. National Government Solutions, Inc. Hypoglossal Nerve Stimulation for the Treatment of Obstructive Sleep Apnea Local Coverage Determination (LCD) L38387. Indianapolis, IN: National Government Solutions, Inc. Available at: www.ngsmedicare.com.
126. Netzer NC, Stoohs RA, Netzer CM, et al. Using the Berlin Questionnaire to Identify Patients at Risk for the Sleep Apnea Syndrome. *Ann Intern Med*. 1999;131(7):485-491. Doi: 10.7326/0003-4819-131-7-199910050-00002.
127. Nord R, Fitzpatrick T, DeShazo JP, Reiter ER. Comparison of readmission and complication rates between traditional sleep surgery and hypoglossal nerve stimulation. *Laryngoscope Invest Otolaryngol*. 2022;7(5):1659-1666. Published 2022 Sep 22. doi:10.1002/lto2.883
128. Noridian Healthcare Solutions, LLC. Hypoglossal Nerve Stimulation for the Treatment of Obstructive Sleep Apnea Local Coverage Determination (LCD) L38310. Fargo, ND: Noridian Healthcare Solutions, LLC. Available at: www.noridianmedicare.com.
129. Novitas Solutions, Inc. Hypoglossal Nerve Stimulation for the Treatment of Obstructive Sleep Apnea Local Coverage Determination (LCD) L38385. Mechanicsburg, PA: Novitas Solutions, Inc. Available at: www.novitas-solutions.com.
130. Parra O, Arboix A, Bechich S, et al. Time course of sleep-related breathing disorders in first-ever stroke or transient ischemic attack. *Am J Respir Crit Care Med*. 2000;161(2 Pt 1):375-380. doi:10.1164/ajrccm.161.2.9903139
131. Patel NP, Ahmed M, Rosen I. Split-Night Polysomnography. *Chest*. 2007;132(5):1664-1671. doi:10.1378/chest.06-1801.
132. Patient Pathway. remede® System. Published November 15, 2023. Accessed November 15, 2024. <https://remede.zoll.com/clinicians/patient-pathway/>
133. Patil SP, Ayappa IA, Caples SM, Kimoff RJ, Patel SR, Harrod CG. Treatment of Adult Obstructive Sleep Apnea With Positive Airway Pressure: An American Academy of Sleep Medicine Systematic Review, Meta-Analysis, and GRADE Assessment. *J Clin Sleep Med*. 2019;15(2):301-334. Published 2019 Feb 15. doi:10.5664/jcsm.7638
134. Patil SP, Ayappa IA, Caples SM, et al. Treatment of Adult Obstructive Sleep Apnea with Positive Airway Pressure: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2019;15(02):335-343. doi:10.5664/jcsm.7640.
135. Patil SP, Collop NA, Chediak AD, et al; ONMAP Technical Expert Panel. Optimal NIV Medicare Access Promotion: Patients With OSA: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. 2021 Nov;160(5):e409-e417. doi: 10.1016/j.chest.2021.06.084.
136. Pavwoski P, Shelgikar AV. Treatment options for obstructive sleep apnea. *Neurology: Clinical Practice*. 2016;7(1):77-85. doi:10.1212/cpj.0000000000000320.
137. Peppard PE, Young T, Palta M, Dempsey J, Skatrud J. Longitudinal study of moderate weight change and sleep-disordered breathing. *JAMA*. 2000;284(23):3015-3021. doi:10.1001/jama.284.23.3015
138. Piccirillo F, Crispino SP, Buzzelli L, Segreti A, Incalzi RA, Grigioni F. A State-of-the-Art Review on Sleep Apnea Syndrome and Heart Failure. *Am J Cardiol*. 2023 May 15;195:57-69. doi: 10.1016/j.amjcard.2023.02.020.
139. Philip P, Chaufton C, Taillard J, et al. Maintenance of Wakefulness Test scores and driving performance in sleep disorder patients and controls. *Int J Psychophysiol*. 2013;89(2):195-202. doi:10.1016/j.ijpsycho.2013.05.013
140. Philip P, Chaufton C, Taillard J, et al. Modafinil improves real driving performance in patients with hypersomnia: a randomized double-blind placebo-controlled crossover clinical trial. *Sleep*. 2014;37(3):483-487. Published 2014 Mar 1. doi:10.5665/sleep.3480
141. Philip P, Sagaspe P, Taillard J, et al. Maintenance of Wakefulness Test, obstructive sleep apnea syndrome, and driving risk. *Ann Neurol*. 2008;64(4):410-416. doi:10.1002/ana.21448
142. Pillar G, Berall M, Berry R, et al. Detecting central sleep apnea in adult patients using WatchPAT-a multicenter validation study. *Sleep Breath*. 2020;24(1):387-398. doi:10.1007/s11325-019-01904-5

143. Pizza F, Contardi S, Mondini S, Trentin L, Cirignotta F. Daytime sleepiness and driving performance in patients with obstructive sleep apnea: comparison of the MSLT, the MWT, and a simulated driving task. *Sleep*. 2009;32(3):382-391. doi:10.1093/sleep/32.3.382
144. Prasad B, Usmani S, Steffen AD, et al. Short-Term Variability in Apnea-Hypopnea Index during Extended Home Portable Monitoring. *J Clin Sleep Med*. 2016;12(6):855-863. Published 2016 Jun 15. doi:10.5664/jcsm.5886
145. Premarket Approval (PMA). www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P130008S039
146. Punjabi NM, Patil S, Crainiceanu C, Aurora RN. Variability and Misclassification of Sleep Apnea Severity Based on Multi-Night Testing. *Chest*. 2020;158(1):365-373. doi:10.1016/j.chest.2020.01.039
147. Ramar K, Dort LC, Katz SG, et al. Clinical Practice Guideline for the Treatment of Obstructive Sleep Apnea and Snoring with Oral Appliance Therapy: An Update for 2015. *J Clin Sleep Med*. 2015;11(7):773-827. doi:10.5664/jcsm.4858.
148. Ramar K, Ramar P, Morgenthaler TI. Adaptive servoventilation in patients with central or complex sleep apnea related to chronic opioid use and congestive heart failure. *J Clin Sleep Med*. 2012;8(5):569-576. Published 2012 Oct 15. doi:10.5664/jcsm.2160
149. Rizzo D, Libman E, Baltzan M, Fichten C, Bailes S. Impact of the COVID-19 pandemic on obstructive sleep apnea: recommendations for symptom management. *J Clin Sleep Med*. 2021;17(3):429-434. doi:10.5664/jcsm.8922.
150. Roeder M, Bradicich M, Schwarz EI, et al. Night-to-night variability of respiratory events in obstructive sleep apnoea: a systematic review and meta-analysis. *Thorax*. 2020;75(12):1095-1102. doi:10.1136/thoraxjnl-2020-214544
151. Rosen IM, Kirsch DB, Chervin RD, et al. Clinical Use of a Home Sleep Apnea Test: An American Academy of Sleep Medicine Position Statement. *J Clin Sleep Med*. 2017;13(10):1205-1207. doi:10.5664/jcsm.6774.
152. Rosen CL, Auckley D, Benca R, et al. A Multisite Randomized Trial of Portable Sleep Studies and Positive Airway Pressure Autotitration versus Laboratory-Based Polysomnography for the Diagnosis and Treatment of Obstructive Sleep Apnea: The Home PAP Study. *Sleep*. 2012;35(6):757-767. doi:10.5665/sleep.1870.
153. Ruoff C, Pizza F, Trotti LM, et al. The MSLT is repeatable in narcolepsy type 1 but not narcolepsy type 2: a retrospective patient study. *J Clin Sleep Med*. 2018;14(1):65-74. <https://doi.org/10.5664/jcsm.6882>.
154. Sagaspe P, Taillard J, Chaumet G, et al. Maintenance of wakefulness test as a predictor of driving performance in patients with untreated obstructive sleep apnea. *Sleep*. 2007;30(3):327-330.
155. Sagaspe P, Micoulaud-Franchi JA, Coste O, et al. Maintenance of Wakefulness Test, real and simulated driving in patients with narcolepsy/hypersomnia. *Sleep Med*. 2019;55:1-5. doi:10.1016/j.sleep.2018.02.009
156. Saletu MT, Kotzian ST, Schwarzsinger A, Haider S, Spatt J, Saletu B. Home Sleep Apnea Testing is a Feasible and Accurate Method to Diagnose Obstructive Sleep Apnea in Stroke Patients During In-Hospital Rehabilitation. *J Clin Sleep Med*. 2018;14(9):1495-1501. Published 2018 Sep 15. doi:10.5664/jcsm.7322
157. Sanchez AM, Germany R, Lozier MR, Schweitzer MD, Koseifi S, Anand R. Central sleep apnea and atrial fibrillation: A review on pathophysiological mechanisms and therapeutic implications. *Int J Cardiol Heart Vasc*. 2020 May 22;30:100527. doi: 10.1016/j.ijcha.2020.100527.
158. Sands SA, Owens RL. Congestive heart failure and central sleep apnea. *Crit Care Clin*. 2015 Jul;31(3):473-95. doi: 10.1016/j.ccc.2015.03.005.
159. Sarber KM, Chang KW, Ishman SL, Epperson MV, Dhanda Patil R. Hypoglossal Nerve Stimulator Outcomes for Patients Outside the U.S. FDA Recommendations. *Laryngoscope*. 2020 Apr;130(4):866-872. doi: 10.1002/lary.28175. Epub 2019 Jul 10. PMID: 31291013.
160. Sateia MJ. International Classification of Sleep Disorders-Third Edition. *Chest*. 2014;146(5):1387-1394. doi:10.1378/chest.14-0970.
161. Schwartz AR, Goldberg LR, McKane S, Morgenthaler TI. Transvenous phrenic nerve stimulation improves central sleep apnea, sleep quality, and quality of life regardless of prior positive airway pressure treatment. *Sleep Breath*. 2021;25(4):2053-2063. doi:10.1007/s11325-021-02335-x
162. Schwarz EI, Scherff F, Haile SR, Steier J, Kohler M. Effect of Treatment of Central Sleep Apnea/Cheyne-Stokes Respiration on Left Ventricular Ejection Fraction in Heart Failure: A Network Meta-Analysis. *J Clin Sleep Med*. 2019;15(12):1817-1825. doi:10.5664/jcsm.8092
163. Schwab RJ, Badr SM, Epstein LJ, et al. An official American Thoracic Society statement: continuous positive airway pressure adherence tracking systems. The optimal monitoring strategies and outcome measures in adults. *Am J Respir Crit Care Med*. 2013;188(5):613-620. doi:10.1164/rccm.201307-1282ST.

164. Shah NM, Shrimanker S, Kaltsakas G. Defining obesity hypoventilation syndrome. *Breathe* (Sheff). 2021;17(3):210089. doi:10.1183/20734735.0089-2021
165. Shapira-Daniels A, Mohanty S, Contreras-Valdes FM, Tieu H, Thomas RJ, Natale A, Anter E. Prevalence of Undiagnosed Sleep Apnea in Patients With Atrial Fibrillation and its Impact on Therapy. *JACC Clin Electrophysiol*. 2020 Nov;6(12):1499-1506. doi: 10.1016/j.jacep.2020.05.030.
166. Sheldon SH, Ferber R, Kryger MH. Diagnosis of Obstructive Sleep Apnea. In: *Principles and Practice of Pediatric Sleep Medicine*. 2nd ed. London: Elsevier Saunders; 2014:221-230.
167. Shott S, Amin R, Chini B et al. Obstructive Sleep Apnea Should all Children with Down Syndrome Be Tested? *Arch Otolaryngol Head Neck Surg*. 2006;132:432-436
168. Sjöblom HM, Nahkuri M, Suomela M, Jero J, Piitulainen JM. Treatment of sleep apnoea with tonsillectomy: a retrospective analysis using long-term follow-up data. *Eur Arch Otorhinolaryngol*. 2022;279(7):3727-3732. doi:10.1007/s00405-022-07350-6
169. Simonneau G, Montani D, Celermajer DS, et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J* 2019; 53: 1801913 doi:10.1183/13993003.01913-2018.
170. Skomro RP, Gjevre J, Reid J, et al. Outcomes of Home-Based Diagnosis and Treatment of obstructive Sleep Apnea. *Chest*. 2010;138(2):257-263. doi:10.1378/chest.09-0577.
171. Selim BJ, Ramar K. Management of Sleep Apnea Syndromes in Heart Failure. *Sleep Medicine Clinics*. 2017;12(1):107-121. doi:10.1016/j.jsmc.2016.10.004.
172. Sleep Medicine Codes. American Academy of Sleep Medicine – Association for Sleep Clinicians and Researchers. <https://aasm.org/clinical-resources/coding-reimbursement/sleep-medicine-codes/>
173. Smith MT, McCrae CS, Cheung J, et al. Use of Actigraphy for the Evaluation of Sleep Disorders and Circadian Rhythm Sleep-Wake Disorders: An American Academy of Sleep Medicine Clinical Practice Guideline. *J Clin Sleep Med*. 2018;14(07):1231-1237. doi:10.5664/jcsm.7230.
174. Soose RJ, Woodson BT, Gillespie MB, et al. Upper Airway Stimulation for Obstructive Sleep Apnea: Self-Reported Outcomes at 24 Months. *J Clin Sleep Med*. 2016;12(01):43-48. doi:10.5664/jcsm.5390.
175. Soose RJ, Faber K, Greenberg H, Boon M, Woodson T, Strollo P. Post-implant care pathway: lessons learned and recommendations after 5 years of clinical implementation of hypoglossal nerve stimulation therapy. *Sleep*. 2021;44(Supplement_1):S4-S10. doi:10.1093/sleep/zsaa279
176. Spencer J, Patel M, Mehta N, et al. Special Consideration Regarding the Assessment and Management of Patients Being Treated with Mandibular Advancement Oral Appliance Therapy for Snoring and Obstructive Sleep Apnea. *Cranio*. 2013;31(1):10-13. doi:10.1179/crn.2013.002.
177. Standards of Practice Committee of the American Academy of Sleep Medicine. Practice parameters for clinical use of the multiple sleep latency test and the maintenance of wakefulness test. *Sleep* 2005;28(1):113-121.
178. Steffen A, Heiser C, Galetke W, et al. Hypoglossal nerve stimulation for obstructive sleep apnea: updated position paper of the German Society of Oto-Rhino-Laryngology, Head and Neck Surgery. *Eur Arch Otorhinolaryngol*. 2022;279(1):61-66. doi:10.1007/s00405-021-06902-6
179. Stevens D, Martins RT, Mukherjee S, Vakulin A. Post-Stroke Sleep-Disordered Breathing-Pathophysiology and Therapy Options. *Front Surg*. 2018;5:9. Published 2018 Feb 26. doi:10.3389/fsurg.2018.00009
180. Strohl MM, Yamauchi M, Peng Z, Strohl KP. Insights since FDA Approval of Hypoglossal Nerve Stimulation for the Treatment of Obstructive Sleep Apnea. *Curr Sleep Med Rep*. 2017 Sep;3(3):133-141. doi: 10.1007/s40675-017-0088-x.
181. Strollo PJ, Soose RJ, Maurer JT, et al. Upper-Airway Stimulation for Obstructive Sleep Apnea. *N Engl J Med*. 2014;370(2):139-149. doi:10.1056/nejmoa1308659.
182. Sutherland K, Vanderveken OM, Tsuda H, et al. Oral Appliance Treatment for Obstructive Sleep Apnea: An Update. *J Clin Sleep Med*. 2014;10(02):215-227. doi:10.5664/jcsm.3460.
183. Tanaka N, Okada M, Tanaka K, Onishi T, Hirao Y, Harada S, Kawahira M, Koyama Y, Fujii K, Watanabe H, Okamura A, Sakata Y, Iwakura K. Sleep apnea severity in patients undergoing atrial fibrillation ablation: Home sleep apnea-test and polysomnography comparison. *J Arrhythm*. 2023 Jun 1;39(4):523-530. doi: 10.1002/joa3.12869.
184. Thaler E, Schwab R, Maurer J, et al. Results of the ADHERE upper airway stimulation registry and predictors of therapy efficacy. *Laryngoscope*. 2020 May;130(5):1333-1338. doi: 10.1002/lary.28286.
185. The Long Term Oxygen Treatment Trial Research Group. A randomized trial of long-term oxygen for COPD with moderate desaturation. *N Engl J Med* 2016;375:1617-27. DOI: 10.1056/NEJMoa1604344.

186. Trotti LM, Staab BA, Rye DB. Test-retest reliability of the multiple sleep latency test in narcolepsy without cataplexy and idiopathic hypersomnia. *J Clin Sleep Med*. 2013;9(8):789–795. <https://doi.org/10.5664/jcsm.2922>.
187. Trotti LM. Twice is nice? Test-retest reliability of the Multiple Sleep Latency Test in the central disorders of hypersomnolence. *J Clin Sleep Med*. 2020;16(suppl_1):17S–18S. doi: 10.5664/jcsm.8884.
188. Tung P, Levitzky YS, Wang R, et al. Obstructive and Central Sleep Apnea and the Risk of Incident Atrial Fibrillation in a Community Cohort of Men and Women. *J Am Heart Assoc*. 2017;6(7):e004500. Published 2017 Jul 1. doi:10.1161/JAHA.116.004500
189. Tschopp S, Wimmer W, Caversaccio M, Borner U, Tschopp K. Night-to-night variability in obstructive sleep apnea using peripheral arterial tonometry: a case for multiple night testing. *J Clin Sleep Med*. 2021;17(9):1751–1758. doi:10.5664/jcsm.9300
190. Unger T, Borghi C, Charchar F, et al. 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2020;75(6):1334–1357. doi:10.1161/HYPERTENSIONAHA.120.15026
191. Verse T, Kroker BA, Pirsig W, Brosch S. Tonsillectomy as a treatment of obstructive sleep apnea in adults with tonsillar hypertrophy. *Laryngoscope*. 2000;110(9):1556–1559. doi:10.1097/00005537-200009000-00029
192. Voigt J, Emani S, Gupta S, Germany R, Khayat R. Meta-Analysis Comparing Outcomes of Therapies for Patients With Central Sleep Apnea and Heart Failure With Reduced Ejection Fraction [published correction appears in *Am J Cardiol*. 2020 Nov 1;134:162. doi: 10.1016/j.amjcard.2020.08.020]. *Am J Cardiol*. 2020;127:73–83. doi:10.1016/j.amjcard.2020.04.011
193. Wang D, Yee BJ, Grunstein RR, Chung F. Chronic Opioid Use and Central Sleep Apnea, Where Are We Now and Where To Go? A State of the Art Review. *Anesth Analg*. 2021;132(5):1244–1253. doi:10.1213/ANE.0000000000005378
194. Weinreich G, Armitstead J, Töpfer V, Wang YM, Wang Y, Teschler H. Validation of ApneaLink as screening device for Cheyne-Stokes respiration. *Sleep*. 2009;32(4):553–557. doi:10.1093/sleep/32.4.553
195. Wolfe LF, Benditt JO, Aboussouan L, et al. ONMAP Technical Expert Panel. Optimal NIV Medicare Access Promotion: Patients With Thoracic Restrictive Disorders: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. 2021 Nov;160(5):e399–e408. doi: 10.1016/j.chest.2021.05.075.
196. Woodson BT, Soose RJ, Gillespie MB, et al. Three-Year Outcomes of Cranial Nerve Stimulation for Obstructive Sleep Apnea. *Otolaryngol Head Neck Surg*. 2016;154(1):181–188. doi:10.1177/0194599815616618.
197. Woodson BT, Strohl KP, Soose RJ, et al. Upper Airway Stimulation for Obstructive Sleep Apnea: 5-Year Outcomes. *Otolaryngol Head Neck Surg*. 2018;159(1):194–202. doi:10.1177/0194599818762383.
198. Wolfe LF, Benditt JO, Aboussouan L, Hess DR, Coleman JM 3rd; ONMAP Technical Expert Panel. Optimal NIV Medicare Access Promotion: Patients With Thoracic Restrictive Disorders: A Technical Expert Panel Report From the American College of Chest Physicians, the American Association for Respiratory Care, the American Academy of Sleep Medicine, and the American Thoracic Society. *Chest*. 2021;160(5):e399–e408. doi:10.1016/j.chest.2021.05.07
199. Yeghiazarians Y, Jneid H, Tietjens JR, et al. Obstructive Sleep Apnea and cardiovascular Disease. *Circulation*. 2021;144:e56–e67. DOI: 10.1161/CIR.0000000000000988.
200. Yeh HK, et al. Summary for clinicians: clinical practice guidelines on home oxygen therapy for adults with chronic lung disease. *Ann Am Thorac Soc* 2021;18(9): 1444–1449.
201. Young T, Peppard PE, Gottlieb DJ. Epidemiology of obstructive sleep apnea: a population health perspective. *Am J Respir Crit Care Med*. 2002;165(9):1217–1239. doi:10.1164/rccm.2109080
202. Young T, Finn L, Peppard PE, et al. Sleep disordered breathing and mortality: eighteen-year follow-up of the Wisconsin sleep cohort. *Sleep*. 2008;31(8):1071–1078.
203. Young T, Palta M, Dempsey J, Peppard PE, Nieto FJ, Hla KM. Burden of sleep apnea: rationale, design, and major findings of the Wisconsin Sleep Cohort study. *WMJ*. 2009;108(5):246–249
204. Zorick F, Roehrs T, Wittig R, Lamphere J, Sickelsteel J, Roth T. Sleep-wake abnormalities in narcolepsy. *Sleep*. 1986;9(1 Pt 2):189–193. doi:10.1093/sleep/9.1.189

Pediatric References

SLPD
v1.0.2025

1. American Academy of Sleep Medicine. International Classification of Sleep Disorders. 3rd ed, text revision. Darien, IL: American Academy of Sleep Medicine; 2023.
2. Amos L, Afolabi-Brown O, Gault D, et al. Age and weight considerations for the use of continuous positive airway pressure therapy in pediatric populations: an American Academy of Sleep Medicine position statement. *J Clin Sleep Med*. 2022 Aug 1;18(8):2041-2043. doi: 10.5664/jcsm.10098.
3. Aurora RN, Lamm CI, Zak RS, et al. Practice parameters for the non-respiratory indications for polysomnography and multiple sleep latency testing for children. *Sleep*. 2012;35(11):1467-1473. Published 2012 Nov 1. doi:10.5665/sleep.2190.
4. American Thoracic Society. Standards and indications for cardiopulmonary sleep studies in children. *American Journal of Respiratory and Critical Care Medicine*. 1996;153(2):866-878. doi:10.1164/ajrccm.153.2.8564147.
5. American Thoracic Society. Cardiorespiratory Sleep Studies in Children. *American Journal of Respiratory and Critical Care Medicine*. 1999;160(4):1381-1387. doi:10.1164/ajrccm.160.4.16041.
6. Aurora RN, Collop NA, Jacobowitz O, et al. Quality Measures for the Care of Adult Patients with Obstructive Sleep Apnea. *Journal of Clinical Sleep Medicine*. January 2015. doi:10.5664/jcsm.4556.
7. Aurora RN, Zak RS, Karippot A, et al. Practice Parameters for the Respiratory Indications for Polysomnography in Children. *Sleep*. 2011;34(3):379-388. doi:10.1093/sleep/34.3.379.
8. Borrelli M, Corcione A, Cimbalo C, et al. Diagnosis of Paediatric Obstructive Sleep-Disordered Breathing beyond Polysomnography. *Children (Basel)*. 2023 Aug 1;10(8):1331. doi:10.3390/children10081331.
9. Carvalho FR, Lentini-Oliveira DA, Prado LB, Prado GF, Carvalho LB. Oral appliances and functional orthopaedic appliances for obstructive sleep apnoea in children. *Cochrane Database of Systematic Reviews*. 2016. doi:10.1002/14651858.cd005520.pub3.
10. Ehsan Z, Ishman SL. Pediatric Obstructive Sleep Apnea. *Otolaryngologic Clinics of North America*. 2016;49(6):1449-1464. doi:10.1016/j.otc.2016.07.001.
11. El-Kersh K, Cavallazzi R, Senthilvel E. Outcomes of adenotonsillectomy in severe pediatric obstructive sleep apnea. *Ear, nose, & throat journal*. www.ncbi.nlm.nih.gov/pubmed/29236274. Published December 2017.
12. Gergund CA, et al. Feasibility of split night polysomnography in children to diagnose and treat sleep related breathing disorders. *Sleep Medicine*. 2022;96:107-112.
13. Huang Y-S, Guilleminault C, Lee L-A, et al. Treatment Outcomes of Adenotonsillectomy for Children with Obstructive Sleep Apnea: A Prospective Longitudinal Study. *Sleep*. 2014;37(1):71-76. doi:10.5665/sleep.3310.
14. Huang Y-S, Guilleminault C. Pediatric Obstructive Sleep Apnea: Where Do We Stand? Sleep-Related Breathing Disorders Advances. *Oto-Rhino-Laryngology*. 2017:136-144. doi:10.1159/000470885.
15. Kang M, Mo F, Witmans M, Santiago V, Tablizo MA. Trends in Diagnosing Obstructive Sleep Apnea in Pediatrics. *Children (Basel)*. 2022 Feb 24;9(3):306. doi:10.3390/children9030306.
16. Kirk V, Baughn J, D'Andrea L, Friedman N, Galion A, Garetz S, Hassan F, Wrede J, Harrod CG, Malhotra RK. American Academy of Sleep Medicine Position Paper for the Use of a Home Sleep Apnea Test for the Diagnosis of OSA in Children. *J Clin Sleep Med*. 2017 Oct 15;13(10):1199-1203. doi:10.5664/jcsm.6772.
17. Lee CH, Hsu WC, Chang WH, Lin MT, Kang KT. Polysomnographic findings after adenotonsillectomy for obstructive sleep apnoea in obese and non-obese children: a systematic review and meta-analysis. *Clin Otolaryngol*. 2016 Oct;41(5):498-510. doi:10.1111/coa.12549.
18. Marcus CL, Brooks LJ, Draper KA, et al. Diagnosis and Management of Childhood Obstructive Sleep Apnea Syndrome. *Pediatrics*. 2012;130(3):576-584. doi:10.1542/peds.2012-1671.
19. Marciuc D, Morarasu S, Morarasu BC, et al. Dental Appliances for the Treatment of Obstructive Sleep Apnea in Children: A Systematic Review and Meta-Analysis. *Medicina (Kaunas)*. 2023 Aug 10;59(8):1447. doi:10.3390/medicina59081447.
20. Marcus CL, Moore RH, Rosen CL, et al. A Randomized Trial of Adenotonsillectomy for Childhood Sleep Apnea. *New England Journal of Medicine*. 2013;368(25):2366-2376. doi:10.1056/nejmoa1215881.

21. Marcus CL, Brooks LJ, Draper KA, et al. Diagnosis and Management of Childhood Obstructive Sleep Apnea Syndrome. *Pediatrics*. 2012;130(3):576-584. doi:10.1542/peds.2012-1671.
22. Mitchell RB, Archer SM, Ishman SL, et al. Clinical Practice Guideline: Tonsillectomy in Children (Update)-Executive Summary. *Otolaryngol Head Neck Surg*. 2019 Feb;160(2):187-205. doi:10.1177/0194599818807917.
23. Mussi N, Forestiero R, Zambelli G, et al. The First-Line Approach in Children with Obstructive Sleep Apnea Syndrome (OSA). *J Clin Med*. 2023 Nov 14;12(22):7092. doi:10.3390/jcm12227092.
24. Pavwoski P, Shelgikar AV. Treatment options for obstructive sleep apnea. *Neurology: Clinical Practice*. 2016;7(1):77-85. doi:10.1212/cpj.0000000000000320.
25. Paruthi S. Telemedicine in Pediatric Sleep. *Sleep Med Clin*. 2020 Sep;15(3S):e1-e7. doi:10.1016/j.jsmc.2020.07.003.
26. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Recommended Amount of Sleep for Pediatric Populations: A Consensus Statement of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2016;12(6):785-786. Published 2016 Jun 15. doi:10.5664/jcsm.5866
27. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Consensus Statement of the American Academy of Sleep Medicine on the Recommended Amount of Sleep for Healthy Children: Methodology and Discussion. *J Clin Sleep Med*. 2016;12(11):1549-1561. Published 2016 Nov 15. doi:10.5664/jcsm.6288
28. Piumelli et al. Apparent Life-Threatening Events (ALTE): Italian guidelines. *Italian Journal of Pediatrics* (2017) 43:111. doi:10.1186/s13052-017-0429-x-x.
29. Roland PS, Rosenfeld RM, Brooks LJ, et al. Clinical Practice Guideline: Polysomnography for Sleep-Disordered Breathing Prior to Tonsillectomy in Children. *Otolaryngology-Head and Neck Surgery*. 2011;145(1_suppl). doi:10.1177/0194599811409837.
30. Savin S, Mezzofranco L, Gracco A, Bruno G, De Stefani A. Management of the Pediatric Patient with Suspected Diagnosis of Obstructive Sleep Apnea Syndrome. *Children (Basel)*. 2023 Jul 14;10(7):1225. doi:10.3390/children10071225.
31. Schechter MS. Technical Report: Diagnosis and Management of Childhood Obstructive Sleep Apnea Syndrome. *Pediatrics*. 2002;109(4). doi:10.1542/peds.109.4.e69.
32. Selvadurai S, Voutsas G, Massicotte C, et al. Positional obstructive sleep apnea in an obese pediatric population. *J Clin Sleep Med*. 2020 Aug 15;16(8):1295-1301. doi: 10.5664/jcsm.8496.
33. Stowe RC and Afolabi-Brown O. Pediatric polysomnography—A review of indications, technical aspects, and interpretation. *Paediatric Respiratory Reviews*. 2020(34):9-17.
34. Tieder JS, Bonkowsky JL, Etzel RA, et al. Brief Resolved Unexplained Events (Formerly Apparent Life Threatening Events) and Evaluation of Lower-Risk Infants. *Pediatrics*. 2016;137(5):e20160590.
35. Todd CA, Bareiss AK, Mccoul ED, Rodriguez KH. Adenotonsillectomy for Obstructive Sleep Apnea and Quality of Life: Systematic Review and Meta-analysis. *Otolaryngology-Head and Neck Surgery*. 2017;157(5):767-773. doi:10.1177/0194599817717480.
36. Watach AJ, Xanthopoulos MS, Afolabi-Brown O, et al. Positive airway pressure adherence in pediatric obstructive sleep apnea: A systematic scoping review. *Sleep Med Rev*. 2020 Jun;51:101273. doi: 10.1016/j.smrv.2020.101273.
37. Van de Perck E, Van Hoorenbeeck K, Verhulst S, et al. Effect of body weight on upper airway findings and treatment outcome in children with obstructive sleep apnea. *Sleep Med*. 2021 Mar;79:19-28. doi: 10.1016/j.sleep.2020.12.028.
38. Yu PK, Stenerson M, Ishman SL, et al. Evaluation of Upper Airway Stimulation for Adolescents With Down Syndrome and Obstructive Sleep Apnea. *JAMA Otolaryngol Head Neck Surg*. 2022;148(6):522-528. doi:10.1001/jamaoto.2022.0455

Questionnaires (SL-8)

SL.AD.102.A
v1.0.2025

Epworth Sleepiness Scale (SL-8.1)

The Epworth Sleepiness Scale is comprised of eight questions, with a maximum score of 24. A score > 10 indicates moderate to high probability of excessive daytime sleepiness. Use the following scale to choose the most appropriate number for each situation:

- 0

Would never doze or sleep.
- 1

Slight chance of dozing or sleeping.
- 2

Moderate chance of dozing or sleeping.
- 3

High chance of dozing or sleeping.

Situation	Chance of dozing or sleeping
Sitting and reading	
Watching TV	
Sitting inactive in a public place	
Being a passenger in a motor vehicle for an hour or more	
Lying down in the afternoon	
Sitting and talking to someone	
Sitting quietly after lunch (no alcohol)	
Stopped for a few minutes in traffic while driving	
Total Epworth Score (add up the points)	

The Berlin Questionnaire (SL-8.2)

The Berlin Questionnaire is comprised of 3 categories and ten questions. Two or more categories with a positive score indicate high probability of OSA.

Patient BMI - _____	
Category 1	Category 2
1. Do you snore? a. Yes b. No c. Don't know <i>If you snore:.....</i>	6. How often do you feel tired or fatigued after your sleep a. Nearly every day b. 3-4 times a week c. 1-2 times a week d. 1-2 times a month e. Never or nearly never
2. Your snoring is: a. Slightly louder than breathing b. As loud as talking c. Louder than talking d. Very loud-can be heard in adjacent rooms	7. During your waking time, do you feel tired, fatigued, or not up to par? a. Nearly every day b. 3-4 times a week c. 1-2 times a week d. 1-2 times a month e. Never or nearly never
3. How often do you snore? a. Almost every day b. 3-4 times a week c. 1-2 times a week d. 1-2 times a month e. Never or almost never	8. Have you ever nodded off or fallen asleep while driving a vehicle a. Yes b. No <i>If yes:</i>
4. Does your snoring bother other people? a. Yes b. No c. Don't know	9. How often does this occur? a. Nearly every day b. 3-4 times a week c. 1-2 times a week d. 1-2 times a month e. Never or nearly never
5. Has anyone noticed that you quit breathing during your sleep? a. Nearly every day b. 3-4 times a week c. 1-2 times a week d. 1-2 times a month e. Never or nearly never	Category 3 a. Do you have high blood pressure? b. Yes c. No Don't know

Berlin Questionnaire scoring

Category 1: Items 1-5

- Item 1: if **Yes**, assign 1 point
- Item 2: if **c** or **d**, assign 1 point
- Item 3: if **a** or **b**, assign 1 point
- Item 4: if **a**, assign 1 point
- Item 5: if **a** or **b**, assign 2 points

- Add points. Category 1 is positive if the total score is 2 or more points.

Category 2: Items 6, 7, 8 (item 9 should be noted separately)

- Item 6: if **a** or **b**, assign 1 point
- Item 7: if **a** or **b**, assign 1 point
- Item 8: if **a**, assign 1 point
- Add points. Category 2 is positive if the total score is 2 or more points.

Category 3

Category 3 is positive if the answer to Item 10 is Yes **OR** if the BMI of the patient is greater than 30 kg/m².

High risk

2 or more categories where the score is positive.

Low risk

1 or no categories where the score is positive.

STOP BANG Questionnaire (SL-8.3)

The **STOP Bang** Questionnaire has eight yes/no questions. A **yes** answer on three or more questions indicates high probability of OSA.

Snoring

1. Do you snore loudly (louder than talking or loud enough to be heard through closed doors)?

Tired

2. Do you often feel tired, fatigued, or sleepy during the daytime?

Observed

3. Has anyone observed you stop breathing during your sleep?

Blood pressure

4. Do you have or are you being treated for high blood pressure?

BMI

5. BMI higher than 35 kg/m²

Age

6. Age over 50 years old?

Neck circumference

7. Neck circumference 16 inches/40 cm or greater?

Gender

8. Gender male?

Intermediate to High risk of OSA

Answering **yes** to three or more items. Please see note below for score of three or greater

Low risk of OSA

Answering **yes** to less than three items.

Note:

High Risk is determined by either one of the following

- Answering “yes” to two or more of four STOP questions + any one BANG question except age
- Answering “yes” to 5 or more questions

Insomnia Severity Index (SL-8.4)

The Insomnia Severity Index has seven questions. The seven answers are added up to get a total score. When you have your total score, look at the *Guidelines for Scoring/ Interpretation* at the bottom of the Insomnia Severity Index page to see where your sleep difficulty fits.

For each question, please **circle** the number that best describes your answer.

Please rate the CURRENT (i.e. LAST 2 WEEKS) SEVERITY of your insomnia problem(s).

Insomnia problem	None	Mild	Moderate	Severe	Very severe
1. Difficulty falling asleep	0	1	2	3	4
2. Difficulty staying asleep	0	1	2	3	4
3. Problem waking up too early	0	1	2	3	4

4. How SATISFIED/DISSATISFIED are you with your CURRENT sleep pattern?

Very satisfied	Satisfied	Moderately satisfied	Dissatisfied	Very dissatisfied
0	1	2	3	4

5. How NOTICEABLE to others do you think your sleep problem is in terms of impairing the quality of your life?

Not at all noticeable	A little	Somewhat	Much	Very much noticeable
0	1	2	3	4

6. How WORRIED/DISTRESSED are you about your current sleep problem?

Not at all worried	A little	Somewhat	Much	Very much worried
0	1	2	3	4

7. To what extent do you consider your sleep problem to INTERFERE with your daily functioning (e.g. daytime fatigue, mood, ability to function at work/daily chores, concentration, memory, mood, etc.) CURRENTLY?

Not at all interfering	A little	Somewhat	Much	Very much interfering
0	1	2	3	4

Guidelines for scoring and interpretation

Add the scores for all seven items (questions 1 + 2 + 3 + 4 + 5 +6 + 7) = _____ your total score.

Total score categories

The score categories are as follows:

0-7	No clinically significant insomnia
8-14	Sub threshold insomnia
15-21	Clinical insomnia (moderate severity)
22-28	Clinical insomnia (severe)