

Pneumatic Compression Devices

DME.SW.101.A

v2.0.2024

HCPCS Codes

E0650, E0651, E0652, E0655, E0660, E0665, E0666, E0667, E0668, E0669, E0670, E0671, E0672, E0673, E0675 (only for PAD), E0676 (only for DVT prevention)

Pneumatic Compression Devices - Criteria

Basic Pneumatic Compression Device

HCPCS codes (E0650, E0651)

- A basic pneumatic compression device is indicated when **one** of the following applies:
 - The individual has documented, symptomatic lymphedema and **both**:
 - Has documented skin changes or measurements have been taken over time
 - Has failed 4 or more weeks of conservative treatments (ie compression garment or bandage, exercise and limb elevation, manual lymphatic drainage)
 - The individual has venous insufficiency with venous leg ulcers and **either**:
 - 6 months of compression treatment have failed
 - The individual cannot tolerate compression treatment

Pneumatic Compression Device with Calibrated Gradient Pressure

HCPCS codes (E0652)

A pneumatic compression device with calibrated gradient pressure is indicated when **all** of the following apply:

- The individual has lymphedema extending to the chest, trunk and/or abdomen
- The individual has documented, symptomatic lymphedema and **both** of the following:
 - Has documented skin changes or measurements have been taken over time
 - Has failed 4 or more weeks of conservative treatments (i.e., compression garment or bandage, exercise and limb elevation, manual lymphatic drainage)

Pneumatic Compression Device

HCPCS codes (E0676)

A pneumatic compression device may be indicated when **both** of the following apply:

- The pump will be used to prevent deep venous thrombosis in an immobile individual

- There is a clinical reason that anticoagulation alone would not be effective

References

1. Chang CJ, Cormier JN. Lymphedema interventions: exercise, surgery, and compression devices. *Semin Oncol Nurs.* 2013;29(1):28-40. doi:10.1016/j.soncn.2012.11.005 .
2. Pavon JM, Adam SS, Razouki ZA, et al. Effectiveness of Intermittent Pneumatic Compression Devices for Venous Thromboembolism Prophylaxis in High-Risk Surgical Patients: A Systematic Review. *J Arthroplasty.* 2016;31(2):524-532. doi:10.1016/j.arth.2015.09.043.
3. Sharfman ZT, Campbell JC, Mirocha JM, Spitzer AI. Balancing Thromboprophylaxis and Bleeding in Total Joint Arthroplasty: Impact of Eliminating Enoxaparin and Predonation and Implementing Pneumatic Compression and Tranexamic Acid. *J Arthroplasty.* 2016;31(6):1307-1312. doi:10.1016/j.arth.2015.11.046.
4. Karaca-Mandic P, Hirsch AT, Rockson SG, Ridner SH. The Cutaneous, Net Clinical, and Health Economic Benefits of Advanced Pneumatic Compression Devices in Patients With Lymphedema. *JAMA Dermatol.* 2015;151(11):1187-1193. doi:10.1001/jamadermatol.2015.1895.
5. Schaverien MV, Moeller JA, Cleveland SD. Nonoperative Treatment of Lymphedema. *Semin Plast Surg.* 2018;32(1):17-21. doi:10.1055/s-0038-1635119.
6. Feldman JL, Stout NL, Wanchai A, Stewart BR, Cormier JN, Armer JM. Intermittent pneumatic compression therapy: a systematic review. *Lymphology* 2012;45(1):13-25.
7. Vojackova N, Fialova J, Hercogova J. Management of lymphedema. *Dermatologic Therapy* 2012;25(4):352-357.
8. Nelson EA, Hillman A, Thomas K. Intermittent pneumatic compression for treating venous leg ulcers. *Cochrane Database Syst Rev.* 2014;(5):CD001899. Published 2014 May 12. doi:10.1002/14651858.CD001899.pub4.
9. Executive Committee of the International Society of Lymphology. The diagnosis and treatment of peripheral lymphedema: 2020 Consensus Document of the International Society of Lymphology. *Lymphology.* 2020;53(1):3-19.
10. Phillips JJ, Gordon SJ. Intermittent Pneumatic Compression Dosage for Adults and Children with Lymphedema: A Systematic Review. *Lymphat Res Biol.* 2019;17(1):2-18. doi:10.1089/lrb.2018.0034.
11. Muluk SC, Hirsch AT, Taffe EC. Pneumatic compression device treatment of lower extremity lymphedema elicits improved limb volume and patient-reported outcomes. *Eur J Vasc Endovasc Surg.* 2013;46(4):480-487. doi:10.1016/j.ejvs.2013.07.012.
12. Dolibog P, Franek A, Taradaj J, et al. A comparative clinical study on five types of compression therapy in patients with venous leg ulcers. *Int J Med Sci.* 2013;11(1):34-43. Published 2013 Dec 14. doi:10.7150/ijms.7548.
13. Kolluri R. Management of venous ulcers. *Tech Vasc Interv Radiol.* 2014;17(2):132-138. doi:10.1053/j.tvir.2014.02.012.