

CIGNA MEDICAL COVERAGE POLICIES - MUSCULOSKELETAL

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Effective Date: March 07, 2026



EviCore
By EVERNORTH

Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by EviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

These guidelines include procedures EviCore does not review for Cigna. Please refer to the [Cigna CPT code list](#) for the current list of high-tech imaging procedures that EviCore reviews for Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five-digit codes, nomenclature and other data are copyright 2025 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

© Copyright 2025 EviCore healthcare

Table of Contents

Guideline	Page
Definitions.....	3
General Guidelines.....	8
Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body.....	10
Diagnostic Arthroscopy.....	12
Arthroscopic Debridement (Chondroplasty) or Loose Body Removal.....	15
Synovectomy.....	18
Meniscectomy or Meniscal Repair.....	21
Meniscal Allograft Transplantation.....	25
Anterior Cruciate Ligament (ACL) Reconstruction and Repair.....	28
Anterolateral Ligament (ALL) Reconstruction/Lateral Extra-Articular Tenodesis (LEAT).....	31
Posterior Cruciate Ligament (PCL) Reconstruction.....	34
Medial/Lateral Collateral Ligament (MCL/LCL) Repair/Reconstruction.....	37
Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT).....	40
Osteochondral Allograft/Autograft Transplantation Systems (OATS)/Mosaicplasty.....	43
Abrasion Arthroplasty/Subchondral Drilling/Microfracturing.....	47
Procedures for Patellofemoral Conditions.....	50
High Tibial Osteotomy.....	55
Lysis of Adhesions.....	58
Manipulation Under Anesthesia (MUA).....	61
Procedures Not Addressed Elsewhere.....	64
Codes (CMM-312).....	66
References (CMM-312).....	71

Definitions

Guideline	Page
Definitions.....	4

Definitions

CMM.JT.DF.312

v2.0.2025

Arthrofibrosis

a condition of the appendicular skeletal system that has resulted from disease, injury, or surgery, and results in pain and restricted range of motion due to internal scarring of the joint with consequent stiffness.

Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT)

a cell-based cartilage repair surgical technique that utilizes an individual's own cells in an effort to repair damage to articular cartilage with the goal of improving joint function and reducing pain. The procedure involves the collection and culture of articular cartilage cells (i.e., chondrocytes) that are then implanted into the cartilage defect with the intent that the cultured cells will contribute to the regeneration and repair of the articular surface.

- Hybrid Autologous Chondrocyte Implantation (ACI): ACI is combined with other surgical repair techniques of cartilage defects (e.g., osteochondral autograft transfer).

Kellgren-Lawrence Grading System

a radiographic grading system describing osteoarthritic changes to the tibial-femoral joint of the knee. When used, the radiographic findings on plain X-rays are typically reported within one of the following categories:

- Grade I: doubtful narrowing of joint space and possible osteophytic lipping
- Grade II: definite osteophytes and possible narrowing of joint space
- Grade III: moderate multiple osteophytes, definite narrowing of joint space, some sclerosis, and possible deformity of bone contour
- Grade IV: large osteophytes, marked narrowing of joint space, severe sclerosis, and definite deformity of bone contour

Kissing Lesion

an articular cartilage defect on opposing joint surfaces of the knee and that are in contact between either the patella and distal femur or the distal femur and tibia (e.g., bipolar lesion).

Lateral Extra-Articular Tenodesis (LEAT)

techniques that include a heterogeneous group of procedures beyond just reconstruction of the Anterolateral Ligament (ALL): modified Lemaire technique; Marcacci technique; Losee tenodesis; modified iliotibial band tenodesis; and, MacIntosh-modified Coker-Arnold procedure.

MACI® Implant

Until recently, Carticel® (Vericel Corporation, Cambridge, MA [formerly Genzyme Biosurgery]) was the only technology that received FDA approval for the culturing of chondrocytes. MACI® Implant (Matrix Induced Autologous Chondrocyte Implant) received approval from the U.S. Food and Drug Administration December 2016 as an autologous cellularized scaffold indicated for repair of single or multiple symptomatic, full-thickness cartilage defects of the knee with or without bone involvement in adults. MACI® Implant is utilized as part of an ACI procedure in which cartilage cells are removed during arthroscopy, and shipped to a laboratory, where the cells are cultured over a period of several weeks. The cells are seeded on a porcine collagen membrane, and once the culturing process is complete, the cells seeded on the membrane are returned to the surgeon for implantation during the procedure. The membrane is placed into the defect and over several months the cells create a matrix that is intended to cover the articular surface of the knee. The safety and effectiveness of MACI® Implant in joints other than the knee has not been established.

Modified Outerbridge Classification

a system developed for judging articular cartilage injury to the knee. This system allows delineation of varying areas of chondral pathology, based on the qualitative appearance of the cartilage surface as viewed on MRI, and can assist in identifying those injuries that are suitable for repair techniques. The characterization of cartilage in this system is as follows:

- Grade I: softening with swelling
- Grade II: fragmentation and fissuring $<1\text{cm}^2$
- Grade III: fragmentation and fissuring $>1\text{cm}^2$
- Grade IV: subchondral bone exposed

Mosaicplasty (or osteochondral cylinder transplantation)

a surgical technique that consists of harvesting cylindrical bone-cartilage grafts and transplanting them into focal chondral or osteochondral defects in the knee. After excision of the chondral lesion, an abrasion arthroplasty is performed to refresh the base of the defect. The grafting procedure involves collecting grafts from the posterior aspect of the distal femoral articular surfaces (medial condyle, lateral condyle, or trochlea) and implanting the grafts in a mosaic-like pattern that will contribute to regeneration and repair the articular surface. A recipient tunnel is created and sized with a drill bit slightly larger than the length of the graft. The harvested graft is placed in the tunnel by a press-fit method. All subsequent grafts are inserted in a similar pattern.

Non-Surgical Management (with regard to the treatment of lower extremity joint pain)

any provider-directed non-surgical treatment, which has been demonstrated in the scientific literature as efficacious and/or is considered reasonable care in the treatment of lower extremity joint pain. The types of treatment involved can include, but are not limited to, the following: relative rest/ activity modification; weight loss; supervised physiotherapy modalities and therapeutic exercises; prescription and non-prescription medications; assistive devices; and/or, intra-articular injections.

Osteochondral Allograft Transplantation (OATS)

a procedure that is similar to mosaicplasty, involving the use of a larger, single plug that usually fills an entire defect. It is often performed to graft chondral defects that are also associated with anterior cruciate ligament (ACL) tears. This method allows arthroscopic access to both the ACL and the chondral defect for the performance of a repair and the grafting procedure.

Outerbridge Classification

a system that has been developed for judging articular cartilage injury to the knee. This system allows delineation of varying areas of chondral pathology, based on the qualitative appearance of the cartilage surface as viewed by direct visualization intraoperatively, and can assist in identifying those injuries that are suitable for repair techniques. The characterization of cartilage in this system is as follows:

- Grade I: softening with swelling
- Grade II: fragmentation and fissuring $<1\text{cm}^2$
- Grade III: fragmentation and fissuring $>1\text{cm}^2$
- Grade IV: subchondral bone exposed

Subchondral Drilling or Microfracturing

a surgical procedure that is performed after the calcified cartilage is debrided and the surgeon creates tiny fractures in the adjacent bones (using an awl). Blood and bone marrow (which contains stem cells) seep out of the fractures, creating a blood clot that releases cartilage-building cells. The microfractures are treated as an injury by the body, which is why the surgery results in new, replacement cartilage. Studies have shown that microfracturing techniques do not fill the chondral defect fully and the repair material that forms is fibrocartilage. Fibrocartilage is not as mechanically sound as the original hyaline cartilage; it is much denser and is not able to withstand the demands of everyday activities as well as hyaline cartilage. Therefore, fibrocartilage is at a higher risk of breaking down. The procedure is less effective in treating older individuals, overweight individuals, or in larger cartilage lesions. Furthermore, chances are high that after only one or two years, symptoms start to return as the fibrocartilage wears away, forcing the individual to reengage in articular cartilage repair.

General Guidelines

Guideline	Page
General Guidelines.....	9

General Guidelines

CMM.JT.GG.312

v2.0.2025

Application of Guideline

- The determination of medical necessity for the performance of knee surgery is always made on a case-by-case basis.
- For advanced-imaging indications related to knee conditions refer to **MS-25: Knee**.

Health Equity Considerations

Health equity is the highest level of health for all individuals; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which individuals are born, grow, live, work, and age. Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include the following: safe housing, transportation, and neighborhoods; racism, discrimination, and violence; education, job opportunities, and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body

Guideline	Page
Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body.....	11

Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body

CMM.JT.IN.312

v2.0.2025

Arthroscopic or open knee surgery may be considered **medically necessary** when surgery is being performed for fracture, tumor, infection, or foreign body that has led to, or will likely lead to, progressive destruction.

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Diagnostic Arthroscopy

Guideline	Page
Diagnostic Arthroscopy Indications.....	13
Diagnostic Arthroscopy Non-Indications.....	14

Diagnostic Arthroscopy Indications

CMM.JT.IN.312

v2.0.2025

Diagnostic arthroscopy is considered **medically necessary** as a stand-alone procedure when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - absence of Kellgren-Lawrence Grade II or greater findings on plain radiographs
 - MRI or CT arthrogram is inconclusive for internal derangement/pathology
- Physical exam demonstrates ANY of the following findings:
 - limited range of motion
 - evidence of joint swelling/effusion
 - joint line tenderness
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment for at least six (6) months duration
- Failure of provider-directed non-surgical management for at least three (3) months duration

Diagnostic Arthroscopy Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Diagnostic arthroscopy is considered **not medically necessary** for ANY other indication or condition.
- Based on lack of scientific evidence of efficacy and safety, "in-office" diagnostic arthroscopy (e.g., Mi-Eye™, VisionScope®) is considered **not medically necessary**

Arthroscopic Debridement (Chondroplasty) or Loose Body Removal

Guideline	Page
Arthroscopic Debridement (Chondroplasty) or Loose Body Removal Indications.....	16
Arthroscopic Debridement (Chondroplasty) or Loose Body Removal Non-Indications.....	17

Arthroscopic Debridement (Chondroplasty) or Loose Body Removal Indications

CMM.JT.IN.312

v2.0.2025

Arthroscopic debridement (chondroplasty) or loose body removal are considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - absence of Kellgren-Lawrence Grade II or greater findings on plain radiographs
 - **Criteria exception:** The absence of Kellgren-Lawrence grade II or greater findings is not required for loose body removal if there is the presence of an acutely locked knee on physical exam.
 - presence of EITHER of the following findings:
 - MRI or CT arthrogram shows articular cartilage degeneration with ANY of the following additional findings:
 - loose body within the joint
 - unstable articular cartilage flaps
 - meniscal tear that extends to the articular surface (not simply degenerative changes, [i.e., fraying]) in conjunction with articular cartilage degeneration within the same compartment
 - impinging osteophytes that would be reasonably expected to result in mechanical symptoms and loss of knee function
 - orthogonal radiograph shows a loose body within the tibiofemoral or patellofemoral joint space
- Symptoms include BOTH of the following:
 - function-limiting knee pain and/or loss of knee function
 - presence of ANY of the following mechanical symptoms:
 - knee range of motion is "blocked" due to pain
 - giving way, subjective weakness, or buckling
 - painful locking, clicking, catching, or popping during weight-bearing activities
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for the presence of painful locking, clicking, catching, or popping during weight-bearing activities when these symptoms are attributed to an intra-articular loose body or foreign body.

Arthroscopic Debridement (Chondroplasty) or Loose Body Removal Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Arthroscopic debridement (chondroplasty) and loose body removal are considered **not medically necessary** for ANY other indication or condition.

Synovectomy

Guideline	Page
Synovectomy Indications.....	19
Synovectomy Non-Indications.....	20

Synovectomy Indications

CMM.JT.IN.312

v2.0.2025

Synovectomy (limited [e.g., plica or shelf resection]); as a stand-alone procedure ; or, as a major procedure with two (2) or more compartments [e.g., medial and lateral]) is considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - MRI or CT arthrogram shows evidence of synovitis or plica
 - **Criteria exception:** Advanced imaging is not required for the **clinical diagnosis** of patellar clunk syndrome following knee replacement surgery.
 - absence of Kellgren-Lawrence Grade IV findings on plain radiographs
- Presence of ANY of the following conditions:
 - Plica Syndrome
 - inflammatory arthritis (i.e., rheumatoid arthritis, gout, pseudogout, psoriatic arthritis)
 - Pigmented Villonodular Synovitis (PVNS)
 - synovial chondromatosis
 - Lyme synovitis
 - Hemophilia
 - Hemochromatosis
 - non-specific synovitis (e.g., proliferative synovitis, post-operative synovitis as a sequela from a knee replacement, patellar clunk syndrome, cyclops lesion, etc.)
 - recurrent hemarthrosis (e.g., secondary to sickle cell anemia, bleeding diathesis, etc.)
- Physical exam demonstrates ANY of the following findings:
 - limited range of motion
 - evidence of joint swelling/effusion
 - joint line tenderness or plica tenderness
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

Synovectomy Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Synovectomy is considered **not medically necessary** for ANY other indication or condition.

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Meniscectomy or Meniscal Repair

Guideline	Page
Meniscectomy or Meniscal Repair Indications.....	22
Meniscectomy or Meniscal Repair Non-Indications.....	24

Meniscectomy or Meniscal Repair Indications

CMM.JT.IN.312

v2.0.2025

Meniscal Tear

Meniscectomy (partial or total) or meniscal repair is considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - MRI or CT arthrogram shows a meniscal tear that extends to the articular surface (not simply degenerative changes, [i.e., fraying]) **and** correlates with the individual's reported symptoms and physical exam findings
 - absence of Kellgren-Lawrence Grade II or greater findings on plain radiographs
 - **Criteria exception:** The absence of Kellgren-Lawrence grade II or greater findings is not required if there is the presence of a meniscal tear and a locked knee on physical exam.
- Physical exam demonstrates at least TWO of the following findings:
 - limited range of motion
 - evidence of joint swelling/effusion
 - joint line tenderness
 - positive McMurray's Test
 - positive Thessaly Test
 - positive Apley's Compression Test
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required if EITHER of the following conditions are present:
 - an acute traumatic anterior, posterior, medial, or lateral meniscal root tear/ avulsion confirmed on MRI
 - meniscal tear with a locked knee on physical exam

Discoid Lateral Meniscus

Meniscectomy/saucerization for discoid lateral meniscus is considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - MRI confirms the presence of a discoid lateral meniscus
 - absence of Kellgren-Lawrence Grade II or greater findings on plain radiographs
- Physical exam demonstrates at least TWO of the following findings:
 - limited range of motion
 - evidence of joint swelling/effusion
 - joint line tenderness
 - positive McMurray's Test
 - positive Thessaly Test
 - positive Apley's Compression Test
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

Meniscectomy or Meniscal Repair Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Meniscectomy (partial or total) or meniscal repair is considered **not medically necessary** for ANY other indication or condition.

Meniscal Allograft Transplantation

Guideline	Page
Meniscal Allograft Transplantation Indications.....	26
Meniscal Allograft Transplantation Non-Indications.....	27

Meniscal Allograft Transplantation Indications

CMM.JT.IN.312

v2.0.2025

Meniscal allograft transplantation is considered **medically necessary** when ALL of the following criteria have been met:

- History of ANY of the following conditions affecting the meniscus:
 - prior significant trauma resulting in an irreparable meniscal tear
 - has undergone a meniscectomy where at least 50% of the meniscus has been removed
- Physical exam demonstrates ANY of the following findings:
 - limited range of motion
 - evidence of joint swelling/effusion
 - joint line tenderness
- Body Mass Index (BMI) ≤ 35
- Age is ≤ 49 years
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

Meniscal Allograft Transplantation Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Meniscal allograft transplantation is considered **not medically necessary** for ANY other indication, condition, or when EITHER of the following are present:
 - standing radiographs show Kellgren-Lawrence Grade III or IV findings.
 - MRI shows Modified Outerbridge Classification Grade III or IV articular cartilage degeneration in the affected compartment.

Anterior Cruciate Ligament (ACL) Reconstruction and Repair

Guideline	Page
Anterior Cruciate Ligament (ACL) Reconstruction Indications.....	29
Anterior Cruciate Ligament (ACL) Reconstruction and Repair Non-Indications.....	30

Anterior Cruciate Ligament (ACL) Reconstruction Indications

CMM.JT.IN.312

v2.0.2025

Anterior cruciate ligament (ACL) reconstruction (with allograft or autograft) is considered **medically necessary** when ALL of the following criteria have been met:

- MRI, CT arthrogram, or arthroscopy shows a tear/disruption or significant laxity of the anterior cruciate ligament (ACL)
- Physical exam demonstrates ANY of the following findings:
 - positive Lachman's Test
 - positive Anterior Drawer Test
 - positive Pivot Shift Test
- Symptoms include BOTH of the following:
 - function-limiting knee pain and/or loss of knee function which interferes with ANY of the following:
 - ability to carry out age-appropriate activities of daily living
 - demands of employment
 - need to return to activities that require cutting, pivoting, and/or agility in which ACL need to return to activities that require cutting, pivoting, and/or agility in which
 - reports of knee instability which is noted as subjective weakness or buckling.
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required if there is an acute injury setting and joint instability has been documented with ANY of the following additional conditions:
 - need to return to activities that require cutting, pivoting, and/or agility in which ACL insufficiency may predispose to further instability episodes that may result in new articular or meniscal cartilage injuries
 - a confirmed ACL tear **and** a repairable meniscus tear
 - concomitant ligament injuries (i.e., multi-ligamentous knee injury) that require reconstruction to provide stability

Anterior Cruciate Ligament (ACL) Reconstruction and Repair Non- Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Anterior cruciate ligament (ACL) reconstruction is considered **not medically necessary** for ANY other indication or condition.
- Anterior cruciate ligament (ACL) repair is considered **not medically necessary**.

Anterolateral Ligament (ALL) Reconstruction/ Lateral Extra-Articular Tenodesis (LEAT)

Guideline	Page
Anterolateral Ligament (ALL) Reconstruction/Lateral Extra-Articular Tenodesis (LEAT) Indications.....	32
Anterolateral Ligament (ALL) Reconstruction/Lateral Extra-Articular Tenodesis (LEAT) Non-Indications.....	33

Anterolateral Ligament (ALL) Reconstruction/Lateral Extra-Articular Tenodesis (LEAT) Indications

CMM.JT.IN.312

v2.0.2025

Anterolateral ligament (ALL) reconstruction or lateral extra-articular tenodesis (LEAT) is considered **medically necessary** when ALL of the following criteria have been met:

- Anterolateral ligament (ALL) reconstruction or lateral extra-articular tenodesis (LEAT) is required to augment the anterior cruciate ligament (ACL) reconstruction
- MRI, CT arthrogram, or arthroscopy shows a tear/disruption or significant laxity of the anterior cruciate ligament (ACL)
- Physical exam demonstrates ANY of the following findings:
 - positive Lachman's Test
 - positive Anterior Drawer Test
 - positive Pivot Shift Test
- Symptoms include BOTH of the following:
 - function-limiting knee pain and/or loss of knee function which interferes with ANY of the following:
 - ability to carry out age-appropriate activities of daily living
 - demands of employment
 - need to return to activities that require cutting, pivoting, and/or agility in which ACL insufficiency may predispose to further instability episodes that may result in new articular or meniscal cartilage injuries
 - reports of knee instability which is noted as giving way, subjective weakness, or buckling
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required if there is an acute injury setting **and** joint instability has been documented with ANY of the following additional conditions:
 - need to return to activities that require cutting, pivoting, and/or agility in which ACL insufficiency may predispose to further instability episodes that may result in new articular or meniscal cartilage injuries
 - a confirmed ACL tear **and** a repairable meniscus tear
 - concomitant ligament injuries (i.e., multi-ligamentous knee injury) that require reconstruction to provide stability

Anterolateral Ligament (ALL) Reconstruction/Lateral Extra-Articular Tenodesis (LEAT) Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Anterolateral ligament (ALL) reconstruction is considered **not medically necessary** for ANY other indication or condition.
- Lateral extra-articular tenodesis (LEAT) is considered **not medically necessary** for ANY other indication or condition.

Posterior Cruciate Ligament (PCL) Reconstruction

Guideline	Page
Posterior Cruciate Ligament (PCL) Reconstruction Indications.....	35
Posterior Cruciate Ligament (PCL) Reconstruction Non-Indications.....	36

Posterior Cruciate Ligament (PCL) Reconstruction Indications

CMM.JT.IN.312

v2.0.2025

Posterior cruciate ligament (PCL) reconstruction (with allograft or autograft) is considered **medically necessary** when ALL of the following criteria have been met:

- MRI, CT arthrogram, or arthroscopy shows a tear/disruption or significant laxity of the posterior cruciate ligament (PCL)
- Presence of ANY of the following findings:
 - stress radiographs show $\geq 8\text{mm}$ of increased posterior translation
 - Physical exam demonstrates ANY of the following findings:
 - positive Posterior Drawer Sign
 - positive Posterior Sag Sign or Tibial Drop Back Test
 - positive Quadriceps Active Test
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required if there is an acute injury setting **and** joint instability has been documented with EITHER of the following additional conditions
 - need to return to activities that require cutting, pivoting, and/or agility in which PCL insufficiency may predispose to further instability episodes that may result in new articular or meniscal cartilage injuries
 - concomitant ligament injuries (i.e., multi-ligamentous knee injury) that require reconstruction to provide stability

Posterior Cruciate Ligament (PCL) Reconstruction Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Posterior cruciate ligament (PCL) reconstruction is considered **not medically necessary** for ANY other indication or condition.

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Medial/Lateral Collateral Ligament (MCL/LCL) Repair/Reconstruction

Guideline	Page
Medial/Lateral Collateral Ligament (MCL/LCL) Repair/Reconstruction Indications.....	38
Medial/Lateral Collateral Ligament (MCL/LCL) Repair/Reconstruction Non-Indications.....	39

Medial/Lateral Collateral Ligament (MCL/LCL) Repair/Reconstruction Indications

CMM.JT.IN.312

v2.0.2025

Medial/lateral collateral ligament (MCL/LCL) repair/reconstruction (with allograft or autograft) is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT arthrogram shows a tear/disruption of the medial or lateral collateral ligament (MCL/LCL)
- Physical exam demonstrates EITHER of the following findings:
 - positive Valgus Stress Test (medial)
 - positive Varus Stress Test (lateral)
- Symptoms include BOTH of the following:
 - function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
 - reports of knee instability which is noted as giving way, subjective weakness, or buckling.
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for LCL repair/reconstruction if there is an acute injury setting involving the the lateral collateral ligament (LCL) (including the posterolateral corner) with documentation of BOTH of the following additional conditions:
 - total disruption of the lateral collateral ligament (LCL) documented on MRI or CT arthrogram
 - joint instability documented on physical exam

Medial/Lateral Collateral Ligament (MCL/LCL) Repair/Reconstruction Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Medial collateral ligament (MCL) repair/reconstruction (including an isolated MCL repair) is considered **not medically necessary** in an acute injury setting.
- Medial/lateral collateral ligament (MCL/LCL) repair/reconstruction is considered **not medically necessary** for ANY other indication or condition.

Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT)

Guideline	Page
Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT) Indications.....	41
Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT) Non-Indications.....	42

Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT) Indications

CMM.JT.IN.312

v2.0.2025

Autologous chondrocyte implantation (ACI) or autologous chondrocyte transplantation (ACT) (using the MACI® implant) is considered **medically necessary** when ALL of the following criteria have been met:

- Body Mass Index (BMI) ≤ 35
- Age is 15-55 years
- Absence of inflammatory arthritis or other systemic disease affecting the joints
- Presence of ALL of the following of the following arthroscopic or imaging findings:
 - Kellgren-Lawrence Grade II or less on radiographs
 - normal articular cartilage at the lesion border (contained lesion))
 - A full-thickness distal femoral articular surface (i.e., medial condyle, lateral condyle, or trochlea) and/or patellar chondral defect of 1-10cm² in size that has been identified with ANY of the following:
 - CT arthrogram
 - MRI and the Modified Outerbridge Classification is Grade III or IV
 - Arthroscopy and the Outerbridge Classification is Grade III or IV
- Absence of an osteochondritis dissecans (OCD) lesion that requires bone grafting
- When the procedure is performed for femoral and patellar chondral lesions: absence of a Modified Outerbridge Classification Grade III or IV corresponding 'kissing lesion' defect on the distal femur (trochlea, condyles), patella, or tibia is also required.
- Physical exam demonstrates BOTH of the following findings:
 - a stable knee with intact or reconstructed ligaments (ACL or PCL) and menisci
 - **Note:** A concurrent ligament stabilization or meniscal procedure at the time of ACI would be acceptable.
 - normal tibial-femoral and/or patella-femoral alignment
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

Autologous Chondrocyte Implantation (ACI) or Autologous Chondrocyte Transplantation (ACT) Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Autologous chondrocyte implantation is considered **not medically necessary** for ANY other indication, condition, or when ANY of the following are present:
 - any knee joint surgery within six (6) months before screening (excluding surgery to procure a biopsy or a concomitant procedure to prepare the knee for a MACI® implant)
 - total meniscectomy, meniscal allograft, or bucket-handle tear or displaced tear requiring greater than 50% removal of the meniscus in the target knee
 - septic arthritis within one (1) year before screening
 - known history of hypersensitivity to gentamicin, other aminoglycosides, or products of porcine or bovine origin
 - uncorrected congenital blood coagulation disorders
 - cruciate ligament instability
- Hybrid autologous chondrocyte implantation performed with osteochondral autograft transfer system (Hybrid ACI/OATS) technique for the treatment of an osteochondral defect is considered **not medically necessary**.

Osteochondral Allograft/Autograft Transplantation Systems (OATS)/Mosaicplasty

Guideline	Page
Osteochondral Allograft/Autograft Transplantation Systems (OATS)/Mosaicplasty Indications.....	44
Osteochondral Allograft/Autograft Transplantation Systems (OATS)/Mosaicplasty Non-Indications.....	46

Osteochondral Allograft/Autograft Transplantation Systems (OATS)/ Mosaicplasty Indications

CMM.JT.IN.312

v2.0.2025

Osteochondral allograft/autograft transplantation (OATS)/mosaicplasty is considered **medically necessary** when ALL of the following criteria have been met:

- Body Mass Index (BMI) ≤ 35
- Age is ≤ 49 years
- Absence of inflammatory arthritis or other systemic disease affecting the joints
- Presence of ALL of the following imaging and/or arthroscopic findings:
 - Kellgren-Lawrence Grade II or less on radiographs
 - normal articular cartilage at the lesion border (contained lesion)
 - A full-thickness distal femoral articular surface (i.e., medial condyle, lateral condyle, or trochlea) and/or patellar chondral defect of 1-10cm² in size that has been identified with ANY of the following:
 - CT arthrogram
 - MRI and the Modified Outerbridge Classification is Grade III or IV
 - Arthroscopy and the Outerbridge Classification is Grade III or IV
 - Additional imaging findings required based on procedure type
 - osteochondral autograft transplants and mosaicplasty:
 - small (i.e., $\leq 2.5\text{cm}^2$ total) chondral defects with sharp, definite borders surrounded by normal-appearing hyaline cartilage
 - osteochondral allograft transplants:
 - larger (i.e., $\leq 10.0\text{cm}^2$ total) chondral defects with sharp, definite borders surrounded by normal-appearing hyaline cartilage
- When the procedure is performed for femoral and patellar chondral lesions: absence of a Modified Outerbridge Classification Grade III or IV corresponding 'kissing lesion' defect on the distal femur (trochlea, condyles), patella, or tibia is required.
- Physical exam demonstrates BOTH of the following findings:
 - a stable knee with intact or reconstructed ligaments (ACL or PCL) and menisci
 - **Note:** A concurrent ligament stabilization or meniscal procedure at the time of OATS would be acceptable.
 - normal tibial-femoral and/or patella-femoral alignment

- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

Osteochondral Allograft/Autograft Transplantation Systems (OATS)/ Mosaicplasty Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Osteochondral allograft/autograft transplantation (OATS)/mosaicplasty of the distal femoral articular or patellar surface is considered **not medically necessary** for ANY other indication or condition.
- Hybrid autologous chondrocyte implantation performed with osteochondral autograft transfer system (Hybrid ACI/OATS) technique for the treatment of an osteochondral defect is considered **not medically necessary**.

Abrasion Arthroplasty/ Subchondral Drilling/ Microfracturing

Guideline	Page
Abrasion Arthroplasty/Subchondral Drilling/Microfracturing Indications.....	48
Abrasion Arthroplasty/Subchondral Drilling/Microfracturing Non-Indications.....	49

Abrasion Arthroplasty/Subchondral Drilling/Microfracturing Indications

CMM.JT.IN.312

v2.0.2025

Abrasion arthroplasty, subchondral drilling, or microfracturing is considered **medically necessary** when ALL of the following criteria have been met:

- A full-thickness distal femoral articular surface (i.e., medial condyle, lateral condyle, or trochlea) and/or patellar chondral defect $\leq 2.5\text{cm}^2$ in size on the weight-bearing surface that has been identified with ANY of the following:
 - CT arthrogram
 - MRI and the Modified Outerbridge Classification is Grade III or IV
 - Arthroscopy and the Outerbridge Classification is Grade III or IV
- Physical exam demonstrates BOTH of the following findings:
 - a stable knee with intact or reconstructed ligaments (ACL or PCL) and menisci
 - **Note:** A concurrent ligament stabilization or meniscal procedure at the time of abrasion arthroplasty would be acceptable.
 - normal tibial-femoral and/or patella-femoral alignment
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

Abrasion Arthroplasty/Subchondral Drilling/Microfracturing Non-Indications

CMM.JT.NI.312

v2.0.2025

- Abrasion arthroplasty, subchondral drilling, or microfracturing is considered **not medically necessary** for ANY other indication or condition.

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Procedures for Patellofemoral Conditions

Guideline	Page
Procedures for Patellofemoral Conditions Indications.....	51
Procedures for Patellofemoral Conditions Non-Indications.....	54

Procedures for Patellofemoral Conditions Indications

CMM.JT.IN.312

v2.0.2025

Procedures for anterior knee pain (with or without recurrent patellar instability) include both bony and/or soft tissue surgical procedures

- Bony surgical procedures can include, but are not limited to, the following: tibial tubercle osteotomy/tubercleplasty (e.g., Fulkerson, Maquet) and trochleoplasty.
- Soft tissue surgical procedures can include, but are not limited to, the following: medial patellofemoral ligament (MPFL) reconstruction/repair; extensor realignment and/or muscle advancement or release (e.g., Campbell, Goldthwaite type procedure); and, lateral retinacular release.

Medial Patellofemoral Ligament (MPFL) Reconstruction/Repair

Medial patellofemoral ligament (MPFL) reconstruction/repair for anterior knee pain (with or without recurrent patellar instability) is considered **medically necessary** when ALL of the following criteria have been met:

- MPFL tear is identified/confirmed by ANY of the following:
 - identified on MRI, CT, or Ultrasound (US)
 - identified by arthroscopy
 - Physical exam demonstrates EITHER of the following findings:
 - MPFL palpation test findings (with the knee in full extension and the patella medially subluxated) noting tenderness to palpation of the origin of the MPFL
 - Patella Glide Test findings >75% lateral subluxation of the patella width at 30° of knee flexion
- Physical exam demonstrates ANY of the following findings:
 - positive J-Sign
 - positive Moving Patellar Apprehension Test
 - lateral patellar translation > one-half of the patellar width
 - tenderness of the medial or lateral facets
 - Patellar Grind Test (Clarke's Sign)
- Symptoms include ANY of the following:
 - anterior knee pain (e.g., loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)

- recurrent patellar instability which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
 - **Criteria exception:** An acute patellofemoral dislocation with a loose chondral or osteochondral fragment is not required to have reoccurring patellar instability.
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for an acute patellofemoral dislocation with a loose chondral or osteochondral fragment.

Trochleoplasty

Trochleoplasty for anterior knee pain (with or without recurrent patellar instability) is considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - absence of severe patellofemoral arthritis
 - trochlear dysplasia with ANY of the following findings:
 - supratrochlear spur
 - lateral trochlear inclination (LTI) $>11^{\circ}$
 - Crossing Sign
 - Double Contour Sign
- Physical exam demonstrates ANY of the following findings:
 - positive J-Sign
 - positive Moving Patellar Apprehension Test
 - lateral patellar translation $>$ one-half of the patellar width
 - tenderness of the medial or lateral facets
 - Patellar Grind Test (Clarke's Sign)
- Symptoms include ANY of the following:
 - anterior knee pain (e.g., loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
 - recurrent patellar instability which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
 - **Criteria exception:** An acute patellofemoral dislocation with a loose chondral or osteochondral fragment is not required to have reoccurring patellar instability.
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for an acute patellofemoral dislocation with a loose chondral or osteochondral fragment.

Procedures Other Than Medial Patellofemoral Ligament (MPFL) Reconstruction or Trochleoplasty

Procedures other than medial patellofemoral ligament (MPFL) reconstruction/repair or trochleoplasty performed for anterior knee pain (with or without recurrent patellar instability) is considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows ANY of the following findings:
 - radiographic evidence of patellar tilt $>20^{\circ}$
 - Patella Alta (e.g., Insall-Salvati, Blackburne-Peel, Caton-Deschamps ratios)
 - sulcus angle greater than 145°
 - increased tibial tubercle-posterior cruciate distance of $>24\text{mm}$
 - increased TT-TG (tibial tuberosity-trochlear groove) distance of 20mm
 - concordant osteochondral defect of the patellofemoral joint (MRI, CT, or previous arthroscopic procedure)
 - acute patellar dislocation with associated intra-articular fracture
- Physical exam demonstrates ANY of the following findings:
 - positive J-Sign
 - positive Moving Patellar Apprehension Test
 - lateral patellar translation $>$ one-half of the patellar width
 - tenderness of the medial or lateral facets
 - Patellar Grind Test (Clarke's Sign)
- Symptoms include ANY of the following:
 - anterior knee pain (e.g., loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
 - recurrent patellar instability which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
 - **Criteria exception:** An acute patellofemoral dislocation with a loose chondral or osteochondral fragment is not required to have reoccurring patellar instability.
- Failure of provider-directed non-surgical management for at least three (3) months duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for an acute patellofemoral dislocation with a loose chondral or osteochondral fragment.

Procedures for Patellofemoral Conditions Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Procedures for patellofemoral conditions are considered **not medically necessary** for ANY other indication or condition.

High Tibial Osteotomy

Guideline	Page
High Tibial Osteotomy Indications.....	56
High Tibial Osteotomy Non-Indications.....	57

High Tibial Osteotomy Indications

CMM.JT.IN.312

v2.0.2025

High tibial osteotomy is considered **medically necessary** when ALL of the following criteria have been met:

- Age is ≤60 years
- Imaging shows unicompartmental osteoarthritis involving less than one-third of the femoral condylar surface
- Physical exam demonstrates ALL of the following findings:
 - <15° of fixed varus deformity
 - affected knee is capable of at least 90° of flexion
 - joint stability in full extension
 - intact anterior cruciate ligament (ACL)
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least three (3) months duration

High Tibial Osteotomy Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- High tibial osteotomy is considered **not medically necessary** for ANY other indication, condition, or when ANY of the following are present:
 - inflammatory arthropathy (i.e., rheumatoid arthritis)
 - chondrocalcinosis
 - anterior cruciate ligament (ACL) tear
 - osteochondral defect >5mm in depth

Lysis of Adhesions

Guideline	Page
Lysis of Adhesions Indications.....	59
Lysis of Adhesions Non-Indications.....	60

Lysis of Adhesions Indications

CMM.JT.IN.312

v2.0.2025

Lysis of adhesions is considered **medically necessary** when ALL of the following criteria have been met:

- Performed for arthrofibrosis
- Physical exam demonstrates <90° of knee flexion by two (2) months after knee surgery or trauma
- Symptoms include function-limiting knee pain and/or loss of knee function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment
- Failure of provider-directed non-surgical management for at least two (2) months duration, including BOTH of the following:
 - anti-inflammatory medication and/or cortisone injection (unless contraindicated)
 - physical therapy (i.e., active exercise and manual therapy designed to increase joint mobility and range of motion)

Lysis of Adhesions Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Lysis of adhesions is considered **not medically necessary** for ANY other indication or condition.

Manipulation Under Anesthesia (MUA)

Guideline	Page
Manipulation Under Anesthesia (MUA) Indications.....	62
Manipulation Under Anesthesia (MUA) Non-Indications.....	63

Manipulation Under Anesthesia (MUA) Indications

CMM.JT.IN.312

v2.0.2025

- See Manipulation under Anesthesia (MUA) Non-Indications

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Manipulation Under Anesthesia (MUA) Non-Indications

CMM.JT.NI.312

v2.0.2025

Not Medically Necessary

- Manipulation under anesthesia (MUA) of a knee joint is included in all arthroscopic knee procedures and is therefore considered incidental to the base procedure requiring medical necessity review.

Procedures Not Addressed Elsewhere

Guideline	Page
Procedures Not Addressed Elsewhere Non-Indications.....	65

Procedures Not Addressed Elsewhere Non-Indications

CMM.JT.NI.312

v2.0.2025

Experimental, Investigational, or Unproven (EIU)

Based on lack of scientific evidence of efficacy and safety, the following are considered **experimental, investigational, or unproven**:

- knee subchondroplasty
- focal resurfacing of a single knee joint defect (e.g., Arthrosurface® femoral condyle implant, HemiCAP®, UniCAP®)

Codes (CMM-312)

Guideline	Page
Codes (CMM-312).....	67

Codes (CMM-312)

CMM.JT.PC.312
v2.0.2025

The inclusion of any code in this table does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

Code	Code Description/Definition
27331	Arthrotomy, knee; including joint exploration, biopsy, or removal of loose or foreign bodies
27332	Arthrotomy, with excision of semilunar cartilage (meniscectomy) knee; medial OR lateral
27333	Arthrotomy, with excision of semilunar cartilage (meniscectomy) knee; medial AND lateral
27334	Arthrotomy, with synovectomy, knee; anterior OR posterior
27335	Arthrotomy, with synovectomy, knee; anterior AND posterior including popliteal area
27340	Excision, prepatellar bursa
27347	Excision of lesion of meniscus or capsule (e.g., cyst, ganglion), knee
27355	Excision or curettage of bone cyst or benign tumor of femur
27356	Excision or curettage of bone cyst or benign tumor of femur; with allograft
27357	Excision or curettage of bone cyst or benign tumor of femur; with autograft (includes obtaining graft)
27358	Excision or curettage of bone cyst or benign tumor of femur; with internal fixation (List in addition to code for primary procedure)
27360	Partial excision (craterization, saucerization, or diaphysectomy) bone, femur, proximal tibia and/or fibula (e.g., osteomyelitis or bone abscess)
27403	Arthrotomy with meniscus repair, knee
27405	Repair, primary, torn ligament and/or capsule, knee; collateral

CMM-312: Knee Surgery- Arthroscopic and Open Procedures

Code	Code Description/Definition
27407	Repair, primary, torn ligament and/or capsule, knee; cruciate
27409	Repair, primary, torn ligament and/or capsule, knee; collateral and cruciate ligaments
27412	Autologous chondrocyte implantation, knee
27415	Osteochondral allograft, knee, open
27416	Osteochondral autograft(s), knee, open (e.g., mosaicplasty) (includes harvesting of autograft[s])
27418	Anterior tibial tubercleplasty (e.g., Maquet type procedure)
27420	Reconstruction of dislocating patella; (e.g., Hauser type procedure)
27422	Reconstruction of dislocating patella; with extensor realignment and/or muscle advancement or release (e.g., Campbell, Goldwaite type procedure)
27424	Reconstruction of dislocating patella; with patellectomy
27425	Lateral retinacular release, open
27427	Ligamentous reconstruction (augmentation), knee; extra-articular
27428	Ligamentous reconstruction (augmentation), knee; intra-articular (open)
27429	Ligamentous reconstruction (augmentation), knee; intra-articular (open) and extra-articular
27442	Arthroplasty, femoral condyles or tibial plateau(s), knee
27570	Manipulation of knee joint under general anesthesia (includes application of traction or other fixation devices)
29850	Arthroscopically aided treatment of intercondylar spine(s) and/or tuberosity fracture(s) of the knee, with or without manipulation; without internal or external fixation (includes arthroscopy)

Code	Code Description/Definition
29851	Arthroscopically aided treatment of intercondylar spine(s) and/or tuberosity fracture(s) of the knee, with or without manipulation; with internal or external fixation (includes arthroscopy)
29855	Arthroscopically aided treatment of tibial fracture, proximal (plateau); unicondylar, includes internal fixation, when performed (includes arthroscopy)
29856	Arthroscopically aided treatment of tibial fracture, proximal (plateau); bicondylar, includes internal fixation, when performed (includes arthroscopy)
29866	Arthroscopy, knee, surgical; osteochondral autograft(s) (e.g., mosaicplasty) (includes harvesting of the autograft[s])
29867	Arthroscopy, knee, surgical; osteochondral allograft (e.g., mosaicplasty)
29868	Arthroscopy, knee, surgical; meniscal transplantation (includes arthrotomy for meniscal insertion), medial or lateral
29870	Arthroscopy, knee, diagnostic; with or without synovial biopsy (separate procedure)
29871	Arthroscopy, knee, surgical; for infection, lavage and drainage
29873	Arthroscopy, knee, surgical; with lateral release
29874	Arthroscopy, knee, surgical; for removal of loose body or foreign body (e.g., osteochondritis dissecans fragmentation, chondral fragmentation)
29875	Arthroscopy, knee, surgical; synovectomy, limited (e.g., plica or shelf resection) (separate procedure)
29876	Arthroscopy, knee, surgical; synovectomy, major, two or more compartments (e.g., medial or lateral)
29877	Arthroscopy, knee, surgical; debridement/shaving of articular cartilage (chondroplasty)
29879	Arthroscopy, knee, surgical; abrasion arthroplasty (includes chondroplasty where necessary) or multiple drilling or microfracture

Code	Code Description/Definition
29880	Arthroscopy, knee, surgical; with meniscectomy (medial AND lateral, including any meniscal shaving) including debridement/shaving of articular cartilage (chondroplasty), same or separate compartment(s), when performed
29881	Arthroscopy, knee, surgical; with meniscectomy (medial OR lateral, including any meniscal shaving) including debridement/shaving of articular cartilage (chondroplasty), same or separate compartment(s), when performed
29882	Arthroscopy, knee, surgical; with meniscus repair (medial OR lateral)
29883	Arthroscopy, knee, surgical; with meniscus repair (medial AND lateral)
29884	Arthroscopy, knee, surgical; with lysis of adhesions, with or without manipulation (separate procedure)
29885	Arthroscopy, knee, surgical; drilling for osteochondritis dissecans with bone grafting, with or without internal fixation (including debridement of base of lesion)
29886	Arthroscopy, knee, surgical; drilling for intact osteochondritis dissecans lesion
29887	Arthroscopy, knee, surgical; drilling for intact osteochondritis dissecans lesion with internal fixation
29888	Arthroscopically aided anterior cruciate ligament repair/augmentation or reconstruction
29889	Arthroscopically aided posterior cruciate ligament repair/augmentation or reconstruction
J7330	Autologous cultured chondrocytes, implant

References (CMM-312)

Guideline	Page
References (CMM-312).....	72

References (CMM-312)

CMM.JT.RF.312

v2.0.2025

1. Aaron R, Skolnick A, Reinert S, Ciombor D. Arthroscopic debridement for osteoarthritis of the knee. *J Bone Joint Surg Am*. 2006;88(5):936-943.
2. Abrams G, Alentorn-Geli E, Harris J, Cole B. Treatment of a Lateral Tibial Plateau Osteochondritis Dissecans Lesion With Subchondral Injection of Calcium Phosphate. *Arthrosc Tech*. 2013;2(3):e271-e274. doi:10.1016/j.eats.2013.03.001.
3. Adler V, Pa L, Ko J, et al. Autologous chondrocyte transplantation for the treatment of articular defects of the knee. *Scr Med*. 2003;76(3):241-250.
4. Alleyne K, Galloway M. Management of osteochondral injuries of the knee. *Clin Sports Med*. 2001;20(2):343-364.
5. Altman R, Hochberg M, Moskowitz, R, et al.; Subcommittee on Osteoarthritis Guidelines. Recommendations for the medical management of osteoarthritis of the hip and knee. American College of Rheumatology Subcommittee on Osteoarthritis Guidelines. *Arthritis Rheum*. 2000;43(9):1905-1915.
6. Angadi D, Edwards D, Melton J. Calcium phosphate injection of symptomatic bone marrow lesions of the knee: what is the current clinical evidence?. *Knee Surg Relat Res*. 2020;32(1). doi:10.1186/s43019-019-0013-3.
7. Anley CM, Morris GV, Saithna A, et al. Defining the role of the tibial tubercle-trochlear groove and tibial tubercle-posterior cruciate ligament distances in the work-up of patients with patellofemoral disorders. *Am J Sports Med*. 2015;43(6):1348-1353. doi:10.1177/0363546515576128.
8. Askenberger M, Mostrom EB, Ekstrom W, et al. Operative repair of medial patellofemoral ligament injury versus knee brace in children with an acute first-time traumatic patellar dislocation. *Am J Sports Med*. 2018;46(10):2328-2340. doi:10.1177/0363546518770616.
9. Balcarek P, Oberthür S, Hopfensitz S, et al. Which patellae are likely to redislocate? *Knee Surg Sports Traumatol Arthrosc*. 2013;22(10):2308-2314. doi:10.1007/s00167-013-2650-5.
10. Balcarek P, Walde T, Frosch S, Schüttrumpf J, Wachowski M, Stürmer K. MRI but not arthroscopy accurately diagnoses femoral MPFL injury in first-time patellar dislocations. *Knee Surg Sports Traumatol Arthrosc*. 2012;20(8):1575-1580. doi:10.1007/s00167-011-1775-7.
11. Bartha L, Vajda A, Duska Z, et al. Autologous osteochondral mosaicplasty grafting. *J Orthop Sports Phys Ther*. 2006;36(10):739-750.
12. Beard D, Davies L, Cook J et al. The clinical and cost-effectiveness of total versus partial knee replacement in patients with medial compartment osteoarthritis (TOPKAT): 5-year outcomes of a randomised controlled trial. *Lancet*. 2019;394(10200):746-756. doi:10.1016/s0140-6736(19)31281-4.
13. Beard DJ, Davies L, Cook JA, et al. Rehabilitation versus surgical reconstruction for non-acute anterior cruciate ligament injury (ACL SNNAP): a pragmatic randomised controlled trial. *Lancet*. 2022;400(10352):605-615. doi:10.1016/s0140-6736(22)01424-6.
14. Bentley G, Biant L, Carrington R, et al. A prospective, randomised comparison of autologous chondrocyte implantation versus mosaicplasty for osteochondral defects in the knee. *J Bone Joint Surg Br*. 2003;85(2):223-230.
15. Berger Y, Ftaita S, Thienpont E. Does Medial Patellofemoral Osteoarthritis Influence Outcome Scores and Risk of Revision After Fixed-bearing Unicompartmental Knee Arthroplasty?. *Clin Orthop Relat Res*. 2019;477(9):2041-2047. doi:10.1097/corr.0000000000000738.
16. Bernstein J, Quach T. A perspective on the study of Moseley et al: Questioning the value of arthroscopic knee surgery for osteoarthritis. *Cleve Clin J Med*. 2003;70(5):401, 405-406, 408-410.
17. Biant LC, Bentley G, Vijayan S, et al. Long-term results of autologous chondrocyte implantation in the knee for chronic chondral and osteochondral defects. *Am J Sports Med*. 2014;42(9):2178-2183.
18. Biau D, Tournoux C, Katsahian S et al. Bone-patellar tendon-bone autografts versus hamstring autografts for reconstruction of anterior cruciate ligament: meta-analysis. *BMJ*. 2006;332(7548):995-1001.
19. Bradley J, Heilman D, Katz B, et al. Tidal irrigation as treatment for knee osteoarthritis: A sham-controlled, randomized, double-blinded evaluation. *Arthritis Rheum*. 2002;46(1):100-108.

20. Briggs T, Mahroof S, David L, et al. Histological evaluation of chondral defects after autologous chondrocyte implantation of the knee. *J Bone Joint Surg Br.* 2003;85(7):1077-1083.
21. Brignardello-Petersen R, Guyatt GH, Buchbinder R, et al. Knee arthroplasty versus conservative management in patients with degenerative knee disease: a systematic review. *BMJ Open.* 2017;7:e016114. doi:10.1136/bmjopen-2017-016114.
22. Brouwer, Reinoud W, Huizinga, Maarten R, Duivenvoorden, Tijs, van Raaij, Tom M, Verhagen, Arianne P, Bierma-Zeinstra, Sita MA, Verhaar, Jan AN. Osteotomy for treating knee osteoarthritis. *Cochrane Database Sys Rev.* 2014;12:CD004019. doi:10.1002/14651858.CD004019.pub4.
23. Burger J, Kleeblad L, Laas N, Pearle A. The Influence of Preoperative Radiographic Patellofemoral Degenerative Changes and Malalignment on Patellofemoral-Specific Outcome Scores Following Fixed-Bearing Medial Unicompartmental Knee Arthroplasty. *J Bone Joint Surg.* 2019;101(18):1662-1669. doi:10.2106/jbjs.18.01385.
24. Calvert G, Wright R. The use of arthroscopy in the athlete with knee osteoarthritis. *Clin Sports Med.* 2005;24(1):133-152.
25. Camathias C, Speth BM, Rutz E, Schlemmer T, Papp K, Vavken P, Studer K., Solitary Trochleoplasty for Treatment of Recurrent Patellar Dislocation, *JBJS Essent Surg Tech.* 2018;8(2):e11. doi:10.2106/JBJS.ST.17.00039. eCollection 2018 Jun 27.
26. Campbell AB, Knopp MV, Kolovich GP, et al: Preoperative MRI underestimates articular cartilage defect size compared with findings at arthroscopic knee surgery. *Am J Sports Med.* 2013;41:590-595.
27. Cavendish P, Everhart J, Peters N, Sommerfeldt M, Flanigan D. Osteochondral Allograft Transplantation for Knee Cartilage and Osteochondral Defects. *JBJS Rev.* 2019;7(6):e7. doi: 10.2106/jbjs.rvw.18.00123.
28. Chalmers P, Sherman S, Raphael B, Su E. Rheumatoid Synovectomy: Does the Surgical Approach Matter?. *Clin Orthop Relat Res.* 2011;469(7):2062-2071. doi:10.1007/s11999-010-1744-3.
29. Chambers K, Schulzer M. Arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med.* 2002;347:1718.
30. Chatain F, Adeleine P, Chambat P, Neyret P; Society Francaise d'Arthroscopie. A comparative study of medial versus lateral arthroscopic partial meniscectomy on stable knees: 10-year minimum follow-up. *Arthroscopy.* 2003;19(8):842-849.
31. Cohen S, Jones R. An evaluation of the efficacy of arthroscopic synovectomy of the knee in rheumatoid arthritis: 12-24 month results. *J Rheumatol.* 1987;14(3):452-455.
32. Cohen S, Sharkey B. Subchondroplasty for Treating Bone Marrow Lesions. *J Knee Surg.* 2015;29(07):555-563. doi:10.1055/s-0035-1568988.
33. Crawford DC, Safran MR. Osteochondritis Dissecans of the knee. *J Am Acad Orthop Surg.* 2006;14:90-100.
34. Daud A, Safir O, Gross A, Kuzyk P. Outcomes of Bulk Fresh Osteochondral Allografts for Cartilage Restoration in the Knee. *J Bone Joint Surg.* 2021;103(22):2115-2125. doi:10.2106/jbjs.20.00350.
35. DeFazio MW, Curry EJ, Gustin MJ, et al. Return to Sport After ACL Reconstruction With a BTB Versus Hamstring Tendon Autograft: A Systematic Review and Meta-analysis. *Orthop J Sports Med.* 2020;8(12):2325967120964919. doi:10.1177/2325967120964919.
36. Dejour DH, Editorial Commentary: Trochleoplasty: Is It Really That Fearsome and Dangerous a Technique?. *Arthroscopy.* 2020;36(8):2246-2248. doi:10.1016/j.arthro.2020.05.050.
37. Deirmengian CA, Dines JS, Vernace JV, et al: Use of a small-bore needle arthroscope to diagnose intra-articular knee pathology: comparison with magnetic resonance imaging. *Am J Orthop.* 2018;47(2).
38. Dervin G, Stiell I, Rody K, Grabowski J. Effect of arthroscopic debridement for osteoarthritis of the knee on health-related quality of life. *J Bone Joint Surg Am.* 2003;85-A(1):10-19.
39. Dhillon JS, Decilveo AP, Kraeutler MJ, Belk JW, McCulloch PC, Scillia AJ. Third-Generation Autologous Chondrocyte Implantation (Cells Cultured Within Collagen Membrane) Is Superior to Microfracture for Focal Chondral Defects of the Knee Joint: Systematic Review and Meta-analysis. *Arthroscopy.* 2022;38(8):2579-2586. doi:10.1016/j.arthro.2022.02.011.
40. Diduch DR, Kandil A, Burrus MT. Lateral patellar instability in the skeletally mature patient: evaluation and surgical management. *J Am Acad Orthop Surg.* 2018;26:429-439. doi:10.5435/JAAOS-D-16-00052.
41. Dixit S, DiFiori JP, Burton M, et al. Management of Patellofemoral Pain Syndrome. *Am Fam Physician.* 2007;75(2):194-202.
42. Dozin B, Malpeli M, Cancedda R, et al. Comparative evaluation of autologous chondrocyte implantation and mosaicplasty: A multicentered randomized clinical trial. *Clin J Sport Med.* 2005;15(4):220-226.

43. Drapeau-Zgoralski V, Swift B, Caines A, Kerrigan A, Carsen S, Pickell M. Lateral Patellar Instability. *JBJS*. 2023;105(5):397-409. doi:10.2106/jbjs.22.00756.
44. Duif C, Koutah MA, Ackermann O, et al. Combination of autologous chondrocyte implantation (ACI) and osteochondral autograft transfer system (OATS) for surgical repair of larger cartilage defects of the knee joint. A review illustrated by a case report. *Technol Health Care*. 2015;23(5):531-537. doi:10.3233/thc-151003.
45. Dwyer T, Whelan D, Shah P, Ajrawat P, Hoit G, Chahal J. Operative Versus Nonoperative Treatment of Femoroacetabular Impingement Syndrome: A Meta-analysis of Short-Term Outcomes. *Arthroscopy*. 2020;36(1):263-273. doi:10.1016/j.arthro.2019.07.025.
46. Ebert JR, Fallon M, Smith A, et al. Prospective Clinical and Radiologic Evaluation of Patellofemoral Matrix-Induced Autologous Chondrocyte Implantation. *Am J Sports Med*. 2015;43(6):1362-1372.
47. Ebert JR, Schneider A, Fallon M, Wood DJ, Janes GC. A Comparison of 2-Year Outcomes in Patients Undergoing Tibiofemoral or Patellofemoral Matrix-Induced Autologous Chondrocyte Implantation. *Am J Sports Med*. 2017;45(14):3243-3253. doi:10.1177/0363546517724761.
48. Englund M, Guermazi A, Roemer FW, et al. Meniscal tear in knees without surgery and the development of radiographic osteoarthritis among middle-aged and elderly persons: The multicenter osteoarthritis study. *Arthritis Rheum*. 2009;60(3):831-839.
49. Englund M, Roos E, Lohmander L. Impact of type of meniscal tear on radiographic and symptomatic knee osteoarthritis: a sixteen-year follow-up of meniscectomy with matched controls. *Arthritis Rheum*. 2003;48(8):2178-2187.
50. Erggelet C, Vavken P. Microfracture for the treatment of cartilage defects in the knee joint—A golden standard?. *J Clin Orthop Trauma*. 2016;7(3):145-152. doi:10.1016/j.jcot.2016.06.015.
51. Erickson BJ, Strickland SM, Gomoll AH. Indications, Techniques, Outcomes for Matrix-Induced Autologous Chondrocyte Implantation (MACI). *Oper Tech Sports Med*. 2018;26(3):175-182. doi:10.1053/j.otsm.2018.06.002.
52. Eseonu KC, Neale J, Lyons A, Kluzek S. Are Outcomes of Acute Meniscus Root Tear Repair Better Than Debridement or Nonoperative Management? A Systematic Review. *Am J Sports Med*. 2022;50(11):3130-3139. doi:10.1177/03635465211031250.
53. Familiari F, Cinque ME, Chahla J, et al. Clinical outcomes and failure rates of osteochondral allograft transplantation in the knee. *Am J Sports Med*. 2018;46(14):3541-3549. doi:10.1177/0363546517732531.
54. Farina E, Lowenstein N, Chang Y, Arant K, Katz J, Matzkin E. Meniscal and Mechanical Symptoms Are Associated with Cartilage Damage, Not Meniscal Pathology. *J Bone Joint Surg*. 2021;103(5):381-388. doi:10.2106/jbjs.20.01193.
55. Farr J. Autologous Chondrocyte Implantation Improves Patellofemoral Cartilage Treatment Outcomes. *CORR*® 2007;463:187-194.
56. Feeley BT, Lau BC. Biomechanics and Clinical Outcomes of Partial Meniscectomy. *J Am Acad Orthop Surg*. 2018;26(24):853-863. doi:10.5435/jaaos-d-17-00256.
57. Felson D, Buckwalter J. Debridement and lavage for osteoarthritis of the knee. *N Engl J Med*. 2002;347:132-133.
58. Felson D. Osteoarthritis of the knee. *N Engl J Med*. 2006;354:841-848.
59. Filardo G, Kon E, Andriolo L, Di Martino A, Zaffagnini S, Marcacci M. Treatment of “Patellofemoral” Cartilage Lesions with Matrix-Assisted Autologous Chondrocyte Transplantation: A Comparison of Patellar and Trochlear Lesions. *Am J Sports Med*. 2014;42(3):626-634. doi:10.1177/0363546513510884.
60. Fond J, Rodin D, Ahmad S, Nirschl R. Arthroscopic debridement for the treatment of osteoarthritis of the knee: 2- and 5-year results. *Arthroscopy*. 2002;18(8):829-834.
61. Forster M, Straw R. A prospective randomised trial comparing intra-articular Hyalgan injection and arthroscopic washout for knee osteoarthritis. *Knee*. 2003;10(3):291-293.
62. Fowler P, Lubliner J. The predictive value of five clinical signs in the evaluation of meniscal pathology. *Arthroscopy*. 1989;5(3):184-186. doi:10.1016/0749-8063(89)90168-0.
63. Gee M, Peterson C, Zhou M, Bottoni C. Anterior Cruciate Ligament Repair: Historical Perspective, Indications, Techniques, and Outcomes. *J Am Acad Orthop Surg*. 2020;28(23):963-971. doi:10.5435/jaaos-d-20-00077.
64. Getgood AMJ, Bryant DM, Litchfield R, et al. Lateral Extra-articular Tenodesis Reduces Failure of Hamstring Tendon Autograft Anterior Cruciate Ligament Reconstruction: 2-Year Outcomes from the STABILITY Study Randomized Clinical Trial. *Am J Sports Med*. 2020;48(2):285-297. doi:10.1177/0363546519896333.

65. Getgood A, Hewison C, Bryant D et al. No Difference in Functional Outcomes When Lateral Extra-Articular Tenodesis Is Added to Anterior Cruciate Ligament Reconstruction in Young Active Patients: The Stability Study. *Arthroscopy*. 2020;36(6):1690-1701. doi:10.1016/j.arthro.2020.02.015.
66. Gigante A, Enea D, Greco F, et al. Distal realignment and patellar autologous chondrocyte implantation: mid-term results in a selected population. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(1):2-10.
67. Gill TJ, Safran M, Mandelbaum B, et al. A prospective, blinded, multicenter clinical trial to compare the efficacy, accuracy, and safety of in-office diagnostic arthroscopy with magnetic resonance imaging and surgical diagnostic arthroscopy. *Arthroscopy*. 2018;34:2429-2435.
68. Gillogly SD, Arnold RM. Autologous Chondrocyte Implantation and Anteromedialization for Isolated Patellar Articular Cartilage Lesions. *Am J Sports Med*. 2014;42(4):912-920.
69. Gobbi A, Kon E, Berruto M, et al. Patellofemoral Full-Thickness Chondral Defects Treated with Second-Generation Autologous Chondrocyte Implantation. *Am J Sports Med*. 2009;37(6):1083-1092.
70. Gomoll AH, Gillogly SD, Cole BJ, et al. Autologous Chondrocyte Implantation in the Patella: A Multicenter Experience. *Am J Sports Med*. 2014;42(5):1074-1081.
71. Gomoll AH, Yoshioka H, Watanabe A, et al. Preoperative management of cartilage defects by MRI underestimates lesion size. *Cartilage*. 2011;2:389-393. doi:10.1177/1947603510397534.
72. Graf K, Sekiya J, Wojtys E. Long-term results after combined medial meniscal allograft transplantation and anterior cruciate ligament reconstruction: Minimum 8.5-year follow-up study. *Arthroscopy*. 2004;20(2):129-140.
73. Guenther D, Irrazazaval S, Bell KM, et al. The role of extra-articular tenodesis in combined ACL and anterolateral capsular injury. *J Bone Joint Surg Am*. 2017;99(19):1654-1660.
74. Gurusamy P, Pedowitz JM, Carroll AN, et al. Medial Patellofemoral Ligament Reconstruction for Adolescents With Acute First-Time Patellar Dislocation With an Associated Loose Body. *Am J Sports Med*. 2021;49(8):2159-2164. doi:10.1177/03635465211013543.
75. Haasper C, Zelle B, Knobloch K, et al. No mid-term difference in mosaicplasty in previously treated versus previously untreated individuals with osteochondral lesions of the talus. *Arch Orthop Trauma Surg*. 2008;128(5):499-504.
76. Halbrecht JL, Jackson DW: Office arthroscopy: a diagnostic alternative. *Arthroscopy*. 1992;8:320-326.
77. Hangody L, Vászárhelyi G, Hangody L, et al. Autologous osteochondral grafting-technique and long-term results. *Injury*. 2008;39(Suppl 1):S32-S39.
78. Harner C, Waltrip R, Bennett C, et al. Surgical management of knee dislocations. *J Bone Joint Surg Am*. 2004;86-A(2):262-73.
79. Harris JD, Siston RA, Pan X, Flanigan DC. Autologous chondrocyte implantation: a systematic review. *J Bone Joint Surg Am*. 2010;92(12):2220-2233. doi:10.2106/jbjs.j.00049.
80. Henderson I, Tuy B, Connell D, et al. Prospective clinical study of autologous chondrocyte implantation and correlation with MRI at three and 12 months. *J Bone Joint Surg Br*. 2003;85(7):1060-1066.
81. Heyse T, Chong L, Davis J, Haas S, Figgie M, Potter H. MRI Diagnosis of Patellar Clunk Syndrome Following Total Knee Arthroplasty. *HSS Journal*®. 2012;8(2):92-95. doi:10.1007/s11420-011-9258-4.
82. Hunt SA, Jazrawi LM, Sherman OH. Arthroscopic management of osteoarthritis of the knee. *J Am Acad Orthop Surg*. 2002;10(5):356-363. doi:10.5435/00124635-200209000-00007.
83. Hurley ET, Colasanti CA, Anil U, et al. Management of Patellar Instability: A Network Meta-analysis of Randomized Control Trials. *Am J Sports Med*. 2022;50(9):2561-2567. doi:10.1177/03635465211020000.
84. Jackson R, Dieterichs C. The results of arthroscopic lavage and debridement of osteoarthritic knees based on the severity of degeneration: A 4- to 6-year symptomatic follow-up. *Arthroscopy*. 2003;19(1):13-20.
85. Jakob R, Franz T, Gautier E, Mainil-Varlet P. Autologous osteochondral grafting in the knee: Indication, results, and reflections. *Clin Orthop*. 2002;(401):170-184.
86. Johnson LL. Arthroscopic abrasion arthroplasty: a review. *Clin Orthop Relat Res*. 2001;(391 Suppl):S306-S317.
87. Jurgensmeier K, Jurgensmeier D, Kunz D, Fuerst P, Warth L, Daines S. Intra-articular Injections of the Hip and Knee With Triamcinolone vs Ketorolac: A Randomized Controlled Trial. *J Arthroplasty*. 2021;36(2):416-422. doi:10.1016/j.arth.2020.08.036.
88. Kalson N, Borthwick L, Mann D, et al. International consensus on the definition and classification of fibrosis of the knee joint. *Bone Joint J*. 2016;98-B(11):1479-1488. doi:10.1302/0301-620x.98b10.37957.
89. Karataglis D, Green M, Learmonth D. Autologous osteochondral transplantation for the treatment of chondral defects of the knee. *Knee*. 2006;13(1):32-35. doi:10.1016/j.knee.2005.05.006.

90. Karataglis D, Learmonth D. Management of big osteochondral defects of the knee using osteochondral allografts with the MEGA-OATS technique. *Knee*. 2005;12(5):389-393.
91. Katz JN, Wright J, Spindler KP, et al. Predictors and outcomes of crossover to surgery from physical therapy for meniscal tear and osteoarthritis: a randomized trial comparing physical therapy and surgery. *J Bone Joint Surgery Am*. 2016;98(22):1890-1896.
92. Kelly M. Role of arthroscopic debridement in the arthritic knee. *J Arthroplasty*. 2006;21(Suppl 1):9-10.
93. Krych AJ, Reardon PJ, Johnson NR, et al. Non-operative management of medial meniscus posterior horn root tears is associated with worsening arthritis and poor clinical outcome at 5-year follow-up. *Knee Surg Sports Traumatol Arthrosc*. 2017;25(2):383-389. doi:10.1007/s00167-016-4359-8.
94. Kirkley A, Birmingham T, Litchfield R, et al. A Randomized Trial of Arthroscopic Surgery for Osteoarthritis of the Knee. *N Engl J Med*. 2008;59:1097-1107,1169-1170.
95. Kise NJ, Risberg MA, Stensrud S, et al. Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle aged patients: randomized control trial with two year follow-up. *BMJ*. 2016;354:i3740.
96. Kocher MS, Logan CA, Kramer DE. Discoid Lateral Meniscus in Children: Diagnosis, Management, and Outcomes. *J Am Acad Orthop Surg*. 2017;25:736-743.
97. Kohn M, Sassoon A, Fernando N. Classifications in Brief: Kellgren-Lawrence Classification of Osteoarthritis. *Clin Orthopaed and Rel Res®*. 2016;474(8):1886-1893. doi:10.1007/s11999-016-4732-4.
98. Kon E, Filardo G, Di Martino A, Marcacci M. ACI and MACI. *J Knee Surg*. 2012;25(1):17-22. doi:10.1055/s-0031-1299651.
99. Kreuz P, Steinwachs M, Erggelet C, et al. Mosaicplasty with autogenous talar autograft for osteochondral lesions of the talus after failed primary arthroscopic management: A prospective study with a 4-year follow-up. *Am J Sports Med*. 2006;34(1):55-63.
100. Kurosaka M, Yagi M, Yoshiya S, Muratsu H, Mizuno K. Efficacy of the axially loaded pivot shift test for the diagnosis of a meniscal tear. *Int Orthop*. 1999;23(5):271-274. doi:10.1007/s002640050369.
101. Kvist J, Pettersson M. Knee-Related Quality of Life Compared Between 20 and 35 Years After an Anterior Cruciate Ligament Injury Treated Surgically With Primary Repair or Reconstruction, or Nonsurgically. *Am J Sports Med*. 2024;52(2):311-319. doi:10.1177/03635465231218237.
102. Lahav A, Burks R, Greis P, et al. Clinical outcomes following osteochondral autologous transplantation (OATS). *J Knee Surg*. 2006;19(3):169-173.
103. Laupattarakasem W, Laopaiboon M, Laupattarakasem P, Sumananont C. Arthroscopic debridement for knee osteoarthritis. *Cochrane Database Syst Rev*. 2008;(1):CD005118.
104. Lee D, Park Y, Song S, Hwang S, Park J, Kang D. Which Technique Is Better for Treating Patellar Dislocation? A Systematic Review and Meta-analysis. *Arthroscopy*. 2018;34(11):3082-3093.e1. doi:10.1016/j.arthro.2018.06.052.
105. Linko E, Harilainen A, Malmivaara A, Seitsalo S. Surgical versus conservative interventions for anterior cruciate ligament ruptures in adults. *Cochrane Database Syst Rev*. 2005;(2):CD001356.
106. Ma H, Hung S, Wang S, et al. Osteochondral autografts transfer for post-traumatic osteochondral defect of the knee- 2 to 5 years follow-up. *Injury*. 2004;35(12):1286-1292.
107. MACI® (autologous cultured chondrocytes on porcine collagen membrane) Product Insert revised 06/2017.
108. MACI® prescribing information (December 2016). U.S. Food and Drug Administration.
109. Macmull S, Jaiswal PK, Bentley G, et al. The role of autologous chondrocyte implantation in the treatment of symptomatic chondromalacia patellae. *Int Orthop:SICOT*. 2012;36:1371-1377.
110. Marcacci M, Kon E, Zaffagnini S, et al. Multiple osteochondral arthroscopic grafting (mosaicplasty) for cartilage defects of the knee: Prospective study results at 2-year follow-up. *Arthroscopy*. 2005;21(4):462-470.
111. Marx R. Arthroscopic surgery for osteoarthritis of the knee? *N Engl J Med*. 2008;359(11):1169-1170.
112. McMillan S, Schwartz M, Jennings B, et al: In-office diagnostic needle arthroscopy: understanding the potential value for the US healthcare system. *Am J Orthop*. 2017;46:252-256.
113. Migliorini F, Lüning C, Eschweiler J, et al. Isolated Arthroscopic Lateral Retinacular Release for Lateral Patellar Compression Syndrome. *Life*. 2021;11(4):295. doi:10.3390/life11040295.
114. Minas T, Bryant T. The Role of Autologous Chondrocyte Implantation in the Patellofemoral Joint. *CORR®*. 2005;436:30-39.

115. Mistovich RJ, Urwin JW, Fabricant PD, et al. Patellar tendon-lateral trochlear ridge distance. A novel measurement of patellofemoral instability. *Am J Sports Med.* 2018;46(14):3400-3406. doi:10.1177/0363546518809982.
116. Mohindra M, Jain J. *Fundamentals of Orthopedics*. Jaypee Brothers Medical Publishers (P) Ltd;358.
117. Moseley J, O'Malley K, Petersen N, et al. A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *N Engl J Med.* 2002;347:81-88.
118. Murray M, Fleming B, Badger G et al. Bridge-Enhanced Anterior Cruciate Ligament Repair Is Not Inferior to Autograft Anterior Cruciate Ligament Reconstruction at 2 Years: Results of a Prospective Randomized Clinical Trial. *Am J Sports Med.* 2020;48(6):1305-1315. doi:10.1177/0363546520913532.
119. Nawaz SZ, Bentley G, Briggs TW, et al. Autologous chondrocyte implantation in the knee: mid-term to long-term results. *J Bone Joint Surg Am.* 2014;96(10):824-830. doi:10.2106/JBJS.L.01695.
120. Niemeyer P, Steinwachs M, Erggelet C, et al. Autologous chondrocyte implantation for the treatment of retropatellar cartilage defects: clinical results referred to defect localisation. *Arch Orthop Trauma Surg.* 2008;128(11):1223-1231. doi:10.1007/s00402-007-0413-9.
121. Nomura E, Horiuchi Y, Inoue M. Correlation of MR imaging findings and open exploration of medial patellofemoral ligament injuries in acute patellar dislocations. *Knee.* 2002;9(2):139-143. doi:10.1016/s0968-0160(02)00002-9.
122. Noorduyn JCA, van de Graaf VA, Willigenburg NW, et al. Effect of Physical Therapy vs Arthroscopic Partial Meniscectomy in People with Degenerative Meniscal Tears: Five-Year Follow-up of the ESCAPE Randomized Clinical Trial. *JAMA Network Open.* 2022;5(7):e2220394-e2220394. doi:10.1001/jamanetworkopen.2022.20394.
123. Noyes F, Barber-Westin S, Rankin M. Meniscal transplantation in symptomatic individuals less than fifty years old. *J Bone Joint Surg Am.* 2005;87(Suppl 1)(Pt.2):149-165.
124. O'Connor D, Johnston RV, Brignardello-Petersen R, et al. Arthroscopic surgery for degenerative knee disease (osteoarthritis including degenerative meniscal tears). *Cochrane Database Syst Rev.* 2022;2022(3). doi:10.1002/14651858.cd014328.
125. Onggo JR, Rasaratnam HK, Nambiar M, et al. Anterior Cruciate Ligament Reconstruction Alone Versus With Lateral Extra-articular Tenodesis With Minimum 2-Year Follow-up: A Meta-analysis and Systematic Review of Randomized Controlled Trials. *Am J Sports Med.* 2022;50(4):1137-1145. doi:10.1177/03635465211004946.
126. Pareek A, Reardon PJ, Macalena JA, et al. Osteochondral autograft transfer versus microfracture in the knee: a meta-analysis of prospective comparative studies at midterm. *Arthroscopy.* 2016;32(10):2118-2130. doi:10.1016/j.arthro.2016.05.038.
127. Pagliuzzi G, Napoli F, Previtali D, Filardo G, Zaffagnini S, Candrian C. A Meta-analysis of Surgical Versus Nonsurgical Treatment of Primary Patella Dislocation. *Arthroscopy.* 2019;35(8):2469-2481. doi:10.1016/j.arthro.2019.03.047.
128. Pascual-Garrido C, Slabaugh MA, L'Heureux DR, et al. Recommendations and treatment outcomes for patellofemoral articular cartilage defects with autologous chondrocyte implantation: prospective evaluation at average 4-year follow-up. *Am J Sports Med.* 2009;37(Suppl 1):33S-41S.
129. Patel KA, Hartigan DE, Makovicka JL, Dulle DL 3rd, Chhabra A. Diagnostic evaluation of the knee in the office setting using small-bore needle arthroscopy. *Arthrosc Tech.* 2018;7(1):e17-e21. doi:10.1016/j.eats.2017.08.044.
130. Pauyo T, Park JP, Bozzo I, Bernstein M. Patellofemoral Instability Part I: Evaluation and Nonsurgical Treatment. *JAAOS.* 2022;30(22):e1431-e1442. doi:10.5435/jaaos-d-22-00254.
131. Pearse E, Craig D. Partial meniscectomy in the presence of severe osteoarthritis does not hasten the symptomatic progression of osteoarthritis. *Arthroscopy.* 2003;19(9):963-968.
132. Peterson L, Minas T, Brittberg M, Lindahl A. Treatment of osteochondritis dissecans of the knee with autologous chondrocyte transplantation. *J Bone Joint Surg Am.* 2003;85(Suppl 2):17-24.
133. Peterson L, Vasiliadis HS, Brittberg M, Lindahl A. Autologous chondrocyte implantation: a long-term follow-up. *Am J Sports Med.* 2010;38(6):1117-1124.
134. Peterson R, Shelton W, Bomboy A. Allograft versus autograft patellar tendon anterior cruciate ligament reconstruction: A 5-year follow-up. *Arthroscopy.* 2001;17(1):9-13.
135. Puzzitiello R, Waterman B, Agarwalla A et al. Primary Medial Patellofemoral Ligament Repair Versus Reconstruction: Rates and Risk Factors for Instability Recurrence in a Young, Active Patient Population. *Arthroscopy.* 2019;35(10):2909-2915. doi:10.1016/j.arthro.2019.05.007.

136. Rahardja R, Zhu M, Love H, Clatworthy MG, Monk AP, Young SW. Effect of Graft Choice on Revision and Contralateral Anterior Cruciate Ligament Reconstruction: Results from the New Zealand ACL Registry. *Am J Sports Med*. 2020;48(1):63-69. doi:10.1177/0363546519885148.
137. Richter D, Schenck R, Wascher D, Treme G. Knee Articular Cartilage Repair and Restoration Techniques. *Sports Health: A Multidisciplinary Approach*. 2015;8(2):153-160. doi:10.1177/1941738115611350.
138. Roos E, Ostenberg A, Roos H, et al. Long-term outcome of meniscectomy: symptoms, function, and performance tests in individuals with or without radiographic osteoarthritis compared to matched controls. *Osteoarthritis Cartilage*. 2001;9(4):316-324.
139. Roos E, Roos H, Ryd L, Lohmander L. Substantial disability 3 months after arthroscopic partial meniscectomy: A prospective study of individual-relevant outcomes. *Arthroscopy*. 2000;16(6):619-626.
140. Rosenstiel N, Praz C, Ouanezar H, et al. Combined Anterior Cruciate and Anterolateral Ligament Reconstruction in the Professional Athlete: Clinical Outcomes From the Scientific Anterior Cruciate Ligament Network International Study Group in a Series of 70 Patients With a Minimum Follow-Up of 2 Years. *Arthroscopy*. 2019;35(3):885-892. doi:10.1016/j.arthro.2018.09.020.
141. Rossi R, Dettoni F, Bruzzone M, et al. Clinical examination of the knee: know your tools for diagnosis of knee injuries. *BMC Sports Sci Med Rehabil*. 2011;3(25). doi:10.1186/1758-2555-3-25.
142. Ruano-Ravina A, Jato Diaz M. Autologous chondrocyte implantation: a systematic review. *Osteoarthritis Cartilage*. 2006;14(1):47-51.
143. Ryu R, Dunbar V, Morse G. Meniscal allograft replacement: a 1-year to 6-year experience. *Arthroscopy*. 2002;18(9):989-994.
144. Saris D, Price A, Widuchowski W, et al. on behalf of the SUMMIT Study Group. Matrix-Applied Characterized Autologous Cultured Chondrocytes Versus Microfracture: Two-Year Follow-up of a Prospective Randomized Trial. *Am J of Sports Med*. 2014;42(6):1384-1394. doi:10.1177/0363546514528093.
145. Sanders T, Morrison W, Singleton B, Miller M, Cornum K. Medial Patellofemoral Ligament Injury Following Acute Transient Dislocation of the Patella: MR Findings with Surgical Correlation in 14 Patients. *J Computer Assisted Tomography*. 2001;25(6):957-962. doi:10.1097/00004728-200111000-00021.
146. Sanders TL, Pareek A, Johnson NR, Stuart MJ, Dahm DL, Krych AJ. Patellofemoral arthritis after lateral patellar dislocation: a matched population-based analysis. *Am J Sports Med*. 2017;45(5):1012-1017.
147. Seitlinger G, Scheurecker G, Hogler R, et al. Tibial tubercle-posterior cruciate ligament distance: A new measurement to define the position of the tibial tubercle in patients with patellar dislocation. *Am J Sports Med*. 2012;40(5):1119-1125. doi:10.1177/0363546512438762.
148. Sekiya J, Giffin J, Irrgang J, et al. Clinical outcomes after combined meniscal allograft transplantation and anterior cruciate ligament reconstruction. *Am J Sports Med*. 2003;31(6):896-906.
149. Sharpe J, Ahmed S, Fleetcroft J, Martin R. The treatment of osteochondral lesions using a combination of autologous chondrocyte implantation and autograft: Three-year follow-up. *J Bone Joint Surg Br*. 2005;87(5):730-735.
150. Sherman SL, Garrity J, Bauer K, Cook J et al. Fresh osteochondral allograft transplantation for the knee: current concepts. *J Am Acad Orthop Surg*. 2014;22(2):121-133.
151. Siemieniuk RA, Harris IA, Agoritsas T, et al. Arthroscopic surgery for generative knee arthritis and meniscal tears: a clinical practice guideline. *BMJ*. 2017;357:j1982.
152. Sihvonen R, Paavola M, Malmivaara A, et al. FIDELITY (Finnish Degenerative Meniscal Lesion Study) Investigators. Arthroscopic partial meniscectomy versus placebo surgery of a degenerative meniscus tear: a 2-year follow-up of the randomized controlled trial. *Ann Rheum Dis*. 2018;77(2):188-195.
153. Solomon D, Avorn J, Warsi A, et al. Which individuals with knee problems are likely to benefit from nonarthroplasty surgery? Development of a clinical prediction rule. *Arch Intern Med*. 2004;164(5):509-513.
154. Sonnery-Cottet B, Siatha A, Cavalier M, Kajetanek C, Temponi EF, Daggest M, Helito CP, Thaunat M. Anterolateral ligament reconstruction is associated with significantly reduced ACL graft rupture rates at a minimum of follow-up of 2 years: a prospective comparative study of 502 patients from the SANTI study Group. *Am J Sports Med*. 2017;45(7):1547-1557.
155. Spindler KP, Hutson LJ, Zajichek A, et al. MOON Knee Group. Anterior cruciate ligament reconstruction in high school and college athletes: does autograft choice influence anterior cruciate ligament reconstruction rates? *Am J Sports Med*. 2020;48(2):298-309.
156. Stuart M, Lubowitz J. What, if any, are the indications for arthroscopic debridement of the osteoarthritic knee? *Arthroscopy*. 2006;22(3):238-239.

157. Szachnowski P, Wei N, Arnold WJ, et al: Complications of office-based arthroscopy of the knee. *J Rheumatol*. 1995;22:1722-1725.
158. Thorlund JB, Juhl CB, Roos EM, et al. Arthroscopic surgery for degenerative knee: systematic review and meta-analysis of benefits and harms. *BMJ*. 2015;350:h2747. doi: 10.1136/bmj.h2747.
159. Tibbo M, Limberg A, Salib C et al. Acquired Idiopathic Stiffness After Total Knee Arthroplasty. *J Bone Joint Surg*. 2019;101(14):1320-1330. doi:10.2106/jbjs.18.01217.
160. Trinh TQ, Harris JD, Siston RA, Flanigan DC. Improved outcomes with combined autologous chondrocyte implantation and patellofemoral osteotomy versus isolated autologous chondrocyte implantation. *Arthroscopy*. 2013;29(3):566-574.
161. van de Graaf VA, Noorduyn JCA, Willigenburg NW, et al. Effect of early surgery vs physical therapy on knee function among patients with nonobstructive meniscal tears. *JAMA*. 2018;320(13):1328-1337. doi:10.1001/jama.2018.13308.
162. Voigt JD, Mosier M, Huber B. Diagnostic needle arthroscopy and the economics of improved diagnostic accuracy: a cost analysis. *Appl Health Econ Health Policy*. 2014;12:523-535.
163. von Keudell A, Han R, Bryant T, Minas T. Autologous Chondrocyte Implantation to Isolated Patella Cartilage Defects. *Cartilage*. 2017;8(2):146-154.
164. Wang D, Jones KJ, Eliasberg CD, et al. Condyle-Specific Matching Does Not Improve Midterm Clinical Outcomes of Osteochondral Allograft Transplantation in the Knee. *J Bone Joint Surg Am*. 2017;99:1614-1620.
165. Yeung M, Leblanc MC, Ayeni OR, et al. Indications for Medial Patellofemoral Ligament Reconstruction: A Systematic Review. *J Knee Surg*. 2015;29(7):543-554.
166. Zamborsky R, Danisovic L. Surgical Techniques for Knee Cartilage Repair: An Updated Large-Scale Systematic Review and Network Meta-analysis of Randomized Controlled Trials. *Arthroscopy*. 2020;36(3):845-858. doi:10.1016/j.arthro.2019.11.096.
167. Zhang G, Zheng L, Ding H, Li E, Sun B, Shi H. Evaluation of medial patellofemoral ligament tears after acute lateral patellar dislocation: comparison of high-frequency ultrasound and MR. *Euro Radiol*. 2015;25(1):274-281.
168. Zhang W, Moskowitz R, Nuki G et al. OARSI recommendations for the management of hip and knee osteoarthritis, Part II: OARSI evidence-based, expert consensus guidelines. *Osteoarthritis Cartilage*. 2008;16(2):137-162.