

Oxlumo® (lumasiran) Injection

When requesting Oxlumo® (lumasiran) injection, the individual requiring treatment must be diagnosed with the following FDA-approved indication and meet the specific coverage guidelines and applicable safety criteria for the covered indication.

FDA-Approved Indication

Oxlumo is indicated for the treatment of primary hyperoxaluria type 1 (PH1) to lower urinary and plasma oxalate levels in pediatric and adult patients.

Coverage Guidelines

Primary Hyperoxaluria Type 1

The individual must meet **all** of the following criteria for initial authorization:

- Has had a genetic test confirming the diagnosis of Primary Hyperoxaluria Type 1 via identification of an alanine:glyoxylate aminotransferase gene (AGXT) biallelic pathogenic variants;
- Meets one of the following:
 - o Has a urinary oxalate excretion greater than or equal to 0.5 mmol/24 hours/1.73 meters² with the absence of secondary sources of oxalate; OR
 - o Has a urinary oxalate:creatinine ratio above the age-specific upper limit of normal; OR
 - o Has a plasma oxalate level greater than or equal to 20 µmol/L;
- Has not previously received a liver transplant for Primary Hyperoxaluria Type 1;
- Oxlumo is prescribed by or in consultation with a nephrologist or urologist.

The individual must meet **all** of the following criteria for re-authorization:

- Is continuing to receive benefit from Oxlumo (e.g., reduced urinary oxalate excretion, decreased urinary oxalate:creatinine ratio, reduced plasma oxalate levels from baseline [i.e., prior to Oxlumo therapy], or improved or stabilized clinical signs/symptoms of Primary Hyperoxaluria Type 1 [e.g., nephrocalcinosis, formation of renal stones, renal impairment]);
- Has not previously received a liver transplant for Primary Hyperoxaluria Type 1.

Approval duration (initial): 6 months

Approval duration (renewal): 12 months

Dosing Recommendation

Oxlumo Weight-Based Dosing Regimen

Body Weight	Loading Dose	Maintenance Dose*
Less than 10 kg	6 mg/kg once monthly for 3 doses	3 mg/kg once monthly
10 kg to less than 20 kg	6 mg/kg once monthly for 3 doses	6 mg/kg once every 3 months (quarterly)
20 kg and above	3 mg/kg once monthly for 3 doses	3 mg/kg once every 3 months (quarterly)

*Beginning 1 month after the last loading dose

References

1. Oxlumo™ subcutaneous injection [prescribing information]. Cambridge, MA: Alnylam; April 2025.
2. Milliner DS, Harris PC, Cogal AG, et al. Primary Hyperoxaluria Type 1. Gene Reviews® Available at: <https://www.ncbi.nlm.nih.gov/books/NBK1283/>. Updated August 15, 2024. Accessed on October 28, 2025.
3. Primary Hyperoxaluria: MedlinePlus Genetics. U.S. National Library of Medicine; National Institutes of Health; Department of Health and Human Services. Available at: <https://medlineplus.gov/genetics/condition/primary-hyperoxaluria/#resources>. Accessed on October 28, 2025.
4. Cochat P, Rumsby G. Primary hyperoxaluria. *N Engl J Med*. 2013;369(7):649-658.
5. Garrelfs SF, Frishberg Y, Hulton SA, et al. Lumasiran, an RNAi therapeutic for primary hyperoxaluria Type 1. *N Engl J Med*. 2021;384(13):1216-1226.
6. Sas DJ, Magen D, Hayes W, et al. Phase 3 trial of lumasiran for primary hyperoxaluria type 1: a new RNAi therapeutic in infants and young children. *Genet Med*. 2022;24(3):654-662.
7. Michael M, Groothoff JW, Shasha-Lavsky H, et al. Lumasiran for advanced primary hyperoxaluria type 1: phase 3 ILLUMINATE-C. *Am J Kidney Dis*. 2022 July 14. [Epub ahead of print].
8. Michael M, Harvey E, Milliner DS, et al. Diagnosis and management of primary hyperoxalurias: best practices. *Pediatr Nephrol*. 2024;39(11):3143-3155.