



CLINICAL GUIDELINES

Cerebrovascular Intervention

Effective: April 1, 2026

VERSION 1.0.2026

EviCore
By ~~EVERNORTH~~

EviCore healthcare Clinical Decision Support Tool Diagnostic Strategies: This tool addresses common symptoms and symptom complexes. Imaging requests for individuals with atypical symptoms or clinical presentations that are not specifically addressed will require physician review. Consultation with the referring physician, specialist, and/or individual's Primary Care Physician (PCP) may provide additional insight.

EviCore's Clinical Review Criteria ("CRC") and related content is made available for the limited uses of: reference; and individual use, only limited to facilitating the determination of medically necessary and appropriate clinical treatment by clinicians for specific delegated patients under their care. The CRC and related content is proprietary information of EviCore, and copyrighted to the full extent of the law. Except as expressly permitted, you may not modify, copy, reproduce, republish, upload, post, transmit, hyperlink to or from, or distribute in any way the CRC, nor may you sell, transfer, distribute, assign, lease, reproduce, or otherwise use the CRC in commerce, in a manner that competes with us or infringes upon our rights, or for any public or commercial endeavor without our prior and express written consent.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five-digit codes, nomenclature and other data are copyright 2025 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT® book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Table of Contents

Guideline	Page
General Information.....	3
General Information for Vascular Intervention Requests.....	4
Cerebrovascular Intervention.....	9
Cerebrovascular Endovascular Embolization and Stents.....	10
References.....	17

General Information

Guideline	Page
General Information for Vascular Intervention Requests	4

General Information for Vascular Intervention Requests

PVI.100.A
v1.0.2026

Glossary

Terms and abbreviations

Aneurysm	Defined as a diameter 1.5x the normal arterial diameter.
Angioplasty	A procedure that utilizes a catheter with a balloon that is inflated to enlarge a stenotic area.
Ankle-Brachial Index (ABI)	Ratio of the systolic blood pressure (SBP) measured at the ankle to the brachial (arm) SBP.
Atherectomy	A procedure that utilizes a catheter with a sharp blade or laser on the end of the catheter to remove plaque from a blood vessel.
CTO	Chronic total occlusion
Crescendo TIA	Multiple recurrent episodes of TIA over hours to days.
Critical limb ischemia	Severe stenosis or occlusion in the vessels supplying the lower extremity such that limb loss will result without treatment. Symptoms of critical limb ischemia in the lower extremities include but are not limited to non-healing wounds, gangrene and ischemic rest pain.
Dissection	Disruption of the media layer of the aorta with bleeding within and along the wall of the aorta.
Graft	Synthetic material used to replace or repair a segment of an artery or bypass an occluded segment of artery.

High-grade stenosis	A high-grade stenosis is defined as at least a 50% narrowing of an artery.
IIH	Idiopathic intracranial hypertension
Ischemic rest pain	Pain arises from severe arterial occlusive disease in the lower extremities such that the patient experiences pain in the distal aspect of the foot and toes while the limb is in the supine position as would occur with sleep. The pain is relieved with the limb in the dependent position or "dangling from the bed" as the limb is depending on gravity to assist with perfusion.
MMAE	Middle Meningeal Artery Embolization
NASCET	North American Symptomatic Carotid Endarterectomy Trial
Occlusion	Blockage of a blood vessel, typically due to atherosclerotic plaque buildup, resulting in absence of blood flow through at least a portion of the vessel that can lead to ischemic symptoms depending on the location, length and chronicity
Pseudo-aneurysm	Outpouching of blood resulting from disruption of the arterial wall with extravasation of blood contained by periarterial connective tissue and not by the arterial wall layers.
PTA	Percutaneous transluminal angioplasty.
Spider veins	Enlarged, tortuous veins that are usually distributed in a web like cluster. These veins are typically <3mm in diameter.
Stent	A metal scaffold placed inside the artery to maintain patency.
Stenosis	Narrowing or constriction of a blood vessel, typically due to atherosclerotic plaque buildup, that reduces blood flow and can lead to ischemic symptoms depending on the location and severity.
Stent-graft	A metal scaffold covered by fabric material placed inside an artery.

Symptomatic carotid stenosis	Characterized by either a transient ischemic attack or cerebrovascular accident that is in the distribution of known severe carotid stenosis, e g., transient right sided upper and lower extremity paralysis in the setting of 70% left internal carotid artery stenosis.
Symptomatic aneurysm	Unrelenting non-positional back pain in the setting of a known abdominal or thoracic aortic aneurysm. Individuals with a symptomatic aneurysm may or may not have evidence of a free or contained rupture. The presence of symptoms indicate impending rupture.
Varicose veins	Enlarged, tortuous veins often caused by incompetent valves. Veins are typically $\geq 3\text{mm}$ in diameter.
Velocity ratio (V1/V2)	Ratio of peak systolic velocity in the diseased segment of blood vessel demonstrating elevated flow velocities to the peak systolic velocity of blood flow in normal vessel just proximal to area of concern in arteries, or just distal in veins.
Venous reflux	Characterized by incompetent or "leaky" valves that no longer function as one-way valves facilitating the flow of blood from the lower extremities to the heart. This results in pooling of blood in the lower extremities leading to distended engorged veins when the lower extremities are in the dependent position as in sitting or standing.

Documentation Requirements for Vascular Intervention Requests

Documentation needed to complete a prior authorization request for vascular intervention include **all** of the following:

- Procedure proposed matching the clinical need
- Condition being treated by the requested procedure
- Detailed documentation of provider-directed conservative treatment, duration and frequency of treatment, and the response to such treatments, if applicable
- Detailed documentation of any previous intervention and the response
- Detailed documentation of location and size of aneurysmal disease, if present
- Detailed documentation regarding nature of the critical limb ischemia: non-healing wound or ischemic rest pain, if applicable

- Recent (within 6 months) written reports of any of the following diagnostic imaging modalities and studies acceptable for purposes of the peripheral vascular intervention guidelines:
 - Ankle-brachial indices, segmental pressures and pulse volume recordings as applicable
 - Duplex ultrasound including carotid, lower extremity and abdominal
 - CTA or CTV abdomen/pelvis with or without lower extremity run-off
 - MRA or MRV abdomen/pelvis with or without lower extremity run-off
 - Arteriogram or venogram
 - IVUS – intravascular ultrasound
- Recent (within 6 months) clinical evaluation documenting:
 - Symptoms (if lifestyle-limiting, detailed documentation regarding quality-of-life parameters that are affected)
 - Physical exam findings to include location and size of ulcers

Emergent and Urgent Requests

Individuals being evaluated for vascular/endovascular surgery should be screened for the presence of a medical condition that warrants urgent/emergent definitive surgical treatment. Provider directed non-surgical management is **not** required when there is documentation, supported by imaging studies or clinical assessment, of any of the following urgent/emergent conditions:

- Critical limb ischemia
- Symptomatic carotid stenosis
- Crescendo TIAs (multiple recurrent episodes of TIA over hours to days)
- Symptomatic or ruptured aneurysms

An urgent/emergent request based on 2018 NCQA standards for utilization management occurs when the time frame for making routine or non-life-threatening determinations on care **either**:

- Could seriously jeopardize the life, health, or safety of the member or others, due to the member's psychological state
- In the opinion of a practitioner with knowledge of the member's medical or behavioral condition, would subject the member to adverse health consequences without the care or treatment that is the subject of the request.

Procedures to treat arterial disease may be medically necessary on an intra-operative basis

Background and Supporting Information

Prior-authorization requests should be submitted at least two weeks prior to the anticipated date of an elective surgery.

Cerebrovascular Intervention

Guideline	Page
Cerebrovascular Endovascular Embolization and Stents.....	10
References.....	17

Cerebrovascular Endovascular Embolization and Stents

CVI.105.A
v1.0.2026

Procedures Included

- Endovascular procedures may include:
 - Embolization (including coiling)
 - Balloon angioplasty
 - Stent placement
 - Flow diverters

Cerebrovascular Embolization and Stent

Procedure Description	CPT®
Transcatheter permanent occlusion or embolization (e.g., for tumor destruction, to achieve hemostasis, to occlude a vascular malformation), including all radiological supervision and interpretation, intraprocedural roadmapping, and imaging guidance necessary to complete the intervention, percutaneous, any method, central nervous system (intracranial, spinal cord)	61624
Balloon angioplasty, intracranial (e.g., atherosclerotic stenosis), percutaneous [not covered for prophylactic percutaneous transluminal angioplasty of intracranial arteries after aneurysmal subarachnoid hemorrhage] [dual diagnosis needed – subarachnoid hemorrhage and ischemia]	61630
Transcatheter placement of intravascular stent(s), intracranial (e.g., atherosclerotic stenosis), including balloon angioplasty if performed [not covered for prophylactic percutaneous transluminal angioplasty of intracranial arteries after aneurysmal subarachnoid hemorrhage] [dual diagnosis needed – subarachnoid hemorrhage and ischemia]	61635
Transcatheter placement of an intravascular stent(s), open or percutaneous, including radiological supervision and interpretation and including angioplasty within the same vessel, when performed; initial vein	37238

Cerebrovascular Intervention

Procedure Description	CPT®
Transcatheter placement of an intravascular stent(s), open or percutaneous, including radiological supervision and interpretation and including angioplasty within the same vessel, when performed; each additional vein (List separately in addition to code for primary procedure)	37239
Transluminal balloon angioplasty (except dialysis circuit), open or percutaneous, including all imaging and radiological supervision and interpretation necessary to perform the angioplasty within the same vein; initial vein	37248
Transluminal balloon angioplasty (except dialysis circuit), open or percutaneous, including all imaging and radiological supervision and interpretation necessary to perform the angioplasty within the same vein; each additional vein (List separately in addition to code for primary procedure)	37249

Endovascular Treatment of Intracerebral Pathology

Indications

- Endovascular treatment of intracerebral pathology is medically necessary when recent clinical history as described in **General Information for Vascular Intervention Requests** documents **any** of the following:
 - Arteriovenous Malformations for any size
 - Ruptured aneurysm with SAH
 - Symptomatic aneurysm including visual changes or protracted headache without SAH
 - "Blister aneurysm" of the supraclinoid segment of internal carotid artery.
 - Asymptomatic saccular aneurysm with any of the following findings:
 - ≥5 mm in size
 - ≥3 mm aneurysm with dysmorphic features or evidence of growth on follow-up imaging
 - Persistent or recurrent aneurysm filling after prior intervention
 - Highly vascular tumors as primary treatment or prior to surgical resection

Non-indications

- Endovascular treatment is not considered medically necessary for intracranial atherosclerosis

Evidence Discussion

Cerebral aneurysm is a bulging, weakened area in the wall of a blood vessel resulting in an abnormal widening or ballooning greater than 50% of the vessel's normal diameter (width).

The causes of aneurysms are varied. They may be congenital or hereditary, or may be caused by other medical conditions or injury.

The risk of rupture for an asymptomatic aneurysm is 1% per year or less. This risk increases with size, size increase over time, family history, and history of prior subarachnoid hemorrhage (SAH)¹. Endovascular treatment options for aneurysm include coil embolization, balloon remodeling, stent-assisted coil embolization, and/or flow diverters. Treatments for ruptured cerebral aneurysms include surgical clipping, endovascular coiling, and/or use of flow diverters. Stenting of a ruptured aneurysm is associated with increased morbidity and mortality and is only considered when less risky options are not available. Treatment should be done early to prevent re-rupture. With conservative management, the risk of aneurysm re-bleeding is 20% to 30% in the first month and then approximately 3% per year.

Most brain AVMs are sporadic and do not have an underlying genetic cause. Conservative management, endovascular embolization, radiation, and operative resection are four modalities that can be considered in the treatment of brain AVM. These modalities may be performed either in isolation or in combination.

Intracranial atherosclerosis: Stroke or TIA (transient ischemic attack) can be caused by symptomatic intracranial atherosclerosis. For individuals with cerebral ischemia attributable to stenosis of an intracranial artery, the mainstay of treatment consists of risk factor modification and medications. Angioplasty and/or stenting is generally not recommended given the low rate of stroke on medical management and the inherent peri-procedural risk of endovascular treatment.

Middle Meningeal Artery Embolization for Chronic Subdural Hematoma

Indications

Middle Meningeal Artery embolization (MMAE) is considered medically necessary in an individual with any of the following clinical scenarios:

- Asymptomatic chronic subdural hematoma measuring ≥ 8 mm on imaging
- Symptomatic chronic subdural hematoma as an adjunct to surgical decompression

¹ Etminan N, de Sousa DA, Tiseo C, et al. European Stroke Organization (ESO) guidelines on management of unruptured intracranial aneurysms. *Eur Stroke J.* 2022;7(3):V. doi:10.1177/23969873221099736.

- Chronic subdural hematoma in the setting of a coagulopathy on antiplatelet therapy and/or unable to undergo surgical drainage
- Prophylactic embolization after surgical evacuation (even without evidence of post-operative chronic subdural hematoma recurrence)

Evidence Discussion

Chronic subdural hematoma (cSDH) is a common neurological disorder primarily affecting the elderly and has been associated with substantial morbidity and mortality. Surgical drainage has historically been the definitive management strategy for symptomatic cSDH. This has been associated with significant morbidity and recurrence rates up to 5-30%. Middle meningeal artery embolization (MMAE) was initially reported as an option for treatment of cSDH over two decades ago. Since that time, multiple studies have been performed comparing the safety and efficacy of MMAE compared to surgical drainage. A meta-analysis pooling 20 studies with 1416 patients, 718 who underwent MMAE and 698 who were conventionally managed, reported lower rates of recurrence (4.8% vs 21.5%), reoperation (4.4% vs 16.4%), and complications (1.7%-4.9%), in the groups of patients who underwent MMAE.² This has been supported even in patients with thrombocytopenia or on anticoagulation.

A proposed treatment algorithm for cSDH with the use of MMAE has been published. The algorithm supports MMAE alone for asymptomatic (including headaches) cSDH with subdural ≥ 8 mm in size and $\pm$ midline shift. For patients with symptomatic or large cSDH with >5 mm midline shift, MMAE is supported as adjunct to surgery to decrease risk of recurrence.

Additional studies have evaluated the safety of MMAE in the elderly (65-79 years old) and advanced elderly (≥ 80 years) with cSDH, and found it to be safe and effective in management of cSDH.³

A recent systematic review and meta-analysis comparing MMAE to conventional management found that MMAE decreased treatment failure and the need for surgical rescue without furthering the risk of morbidity and mortality, compared to conventional management. The study authors recommended considering MMAE in the management of chronic subdural hematoma.

² Ironside N, Nguyen C, Do Q, et al. Middle meningeal artery embolization for chronic subdural hematoma: a systematic review and meta-analysis. *J Neurointerv Surg*. 2021;13(10):951-957. doi:10.1136/neurintsurg-2021-017352

³ Joyce E, Bounajem MT, Scoville J, et al. Middle meningeal artery embolization treatment of nonacute subdural hematomas in the elderly: a multiinstitutional experience of 151 cases. *Neurosurg Focus*. 2020 Oct;49(4):E5. doi: 10.3171/2020.7.FOCUS20518

Endovascular Intervention for Intracranial Venous Stenosis

Endovascular intervention of intracranial venous stenosis for idiopathic intracranial hypertension (IIH) :

- For IIH **all** the following clinical findings must be present:
 - Papilledema on ocular exam
 - Elevated ICP on spinal tap of >25 cm H₂O
 - Failed medical therapy including lifestyle changes, diuretics, and acetazolamide therapy
 - Venous sinus stenosis documented by either of the following:
 - >50% stenosis on MRV, CTV, or venography
 - A significant pressure gradient across the stenosis (>8 mmHg) during diagnostic catheter venogram under local anesthetic
 - Patient must be able to tolerate dual anti-platelet therapy (ASA and clopidogrel) for 1 month.

Venous stenting is considered medically necessary for treatment of pulsatile tinnitus resulting from **any** of the following clinical conditions:

- Transverse venous sinus stenosis (all must be present)
 - Failed medical therapy including lifestyle changes, diuretics, and acetazolamide therapy
 - Venous Sinus stenosis documented by either of the following:
 - >50% stenosis on MRV, CTV or venography
 - A pressure gradient of > 4 mmHg during diagnostic catheter venogram under local anesthetic
 - Patient must be able to tolerate dual antiplatelet therapy (ASA and clopidogrel) for 1 month.
- Dural arteriovenous fistula
- Venous sinus aneurysm of 3 mm (dome to neck)

Endovascular intervention for other indications, including headache, is not considered medically necessary.

Intervention for recurrent intracranial venous stenosis Intervention for recurrent stenosis in intracranial vein after prior treatment for Intracranial Hypertension is medically necessary in the setting of **any** of the following clinical scenarios:

- Evidence of ≥30% stenosis on MRV, CTV, or angiogram
- Greater than 8 mmHg gradient across the stenosis during invasive angiogram

Evidence Discussion

In 2018, a consensus guideline for management of idiopathic intracranial hypertension (IIH) was published by Mollan et al⁴. In this document, they stated that the role of neurovascular stenting was not yet established for patients with acute IIH to prevent vision loss. They felt it was not indicated for patients with headache but not at risk for vision loss.

Shields et al published a single institution retrospective review of 42 patients over 6 years who underwent venous sinus stenting.⁵ The mean follow-up time was 25.6 months (8.7- 60.7 months). All patients had clinical symptoms including headaches, visual disturbances and papilledema. Post intervention follow-up showed resolution of papilledema in 74% and complete resolution of headaches in 43%. Two patients required re-intervention and one patient experienced in-stent thrombosis.

In 2019, Patsalides published short-term data on pressure measurement changes and reduction in medication requirements in 50 patients who underwent venous sinus stenting.⁶ They noted that 94% of patients had a decrease in CSF pressure post procedure. There was a 45% reduction in mean ICP at 3 months. The average dose of acetazolamide decreased from 950 mg to 300 mg at the 3-month follow up with 70% percent of the patients being able to discontinue their acetazolamide therapy completely at the 3-months.

In 2019, Nicholson et al published a systematic review and meta-analysis of venous stenting for IIH. A total of 474 patients were included in the analysis with mean follow-up of 18 months. The outcomes assessed were resolution of papilledema, headaches and pulsatile tinnitus. They also assessed the rate of recurrence and complications. The overall rate of improvement of papilledema was 93.7% with improvement or resolution of headache in 79.6%. Pulsatile tinnitus resolved in 90.3% of patients. The recurrence rate was 9.8% and complication rate of 1.9%.⁷

Raoof et al also noted that although some studies have shown significant improvement in papilledema and headache, there was also a complication rate of 7.4% with 10.3% of patients requiring repeat intervention. They also agreed with Mollan et al regarding long-term data concerning safety and efficacy was still lacking. They concluded that

⁴ Mollan SP, Davies B, Silver NC, et al. Idiopathic intracranial hypertension: consensus guidelines on management. *J Neurol Neurosurg Psychiatry*. 2018 Oct;89(10):1088-1100. doi: 10.1136/jnnp-2017-317440.

⁵ Shields LBE, Shields CB, Yao TL, et al. Endovascular Treatment for Venous Sinus Stenosis in Idiopathic Intracranial Hypertension: An Observational Study of Clinical Indications, Surgical Technique, and Long-Term Outcomes. *World Neurosurg*. 2019 Jan;121:e165-e171. doi: 10.1016/j.wneu.2018.09.070.

⁶ Patsalides A, Oliveira C, Wilcox J, et al. Venous sinus stenting lowers the intracranial pressure in patients with idiopathic intracranial hypertension. *J NeuroInterv Surg* 2019;11:175–178.

⁷ Nicholson P, Brinjikji W, Radovanovic I, et al. Venous sinus stenting for idiopathic intracranial hypertension: a systematic review and meta-analysis. *J Neurointerv Surg*. 2019 Apr;11(4):380-385. doi: 10.1136/neurintsurg-2018-014172.

neurovascular stenting be reserved for carefully selected patients with clear evidence of elevated pressure gradient across a stenosis.⁸

In 2021, Patsalides et al published a prospective trial of venous stenting for refractory pulsatile tinnitus. This involved 42 patients with venous sinus stenosis with >4 mmHg gradient and post stenotic dilation. At 5 months post-stenting, 98% had complete or near complete resolution.⁹

Most recently, Azzam et al published an updated meta-analysis of venous stenting for IIH in April 2024. This included 36 studies involving 1066 patients to evaluate the safety and efficacy of venous sinus stenting in IIH. Clinical outcomes were demonstrated improvement in tinnitus (95%), papilledema (89%), visual disturbances (88%), and headache (79%).⁷ However, they noted that 8.35% of patients experienced treatment failure, characterized by worsening symptoms and recurrence of IIH. They also noted a total complication rate of 5.35%, including subdural hemorrhage, urinary tract infection, stent thrombus formation and others. They concluded that venous sinus stenting was a safe and effective treatment option for IIH patients who are unresponsive to medical therapy or have significant visual symptoms. However, long-term outcomes and safety of the procedure require further investigation.¹⁰

In January 2025, Su et al published a retrospective study looking at the long-term outcomes for venous sinus stenting for pulsatile stenosis. A total of 80 patients were included with a mean duration of follow-up of 80 months and 26 patients having more than 120 months followup. All had resolution of pulsatile tinnitus following stenting and there were no cases of recurrence. Follow-up imaging was available in 23 patients with only one recurrent stenosis identified.¹¹

⁸ Raoof N, Hoffmann J. Diagnosis and treatment of idiopathic intracranial hypertension. *Cephalalgia*. 2021 Apr;41(4):472-478. doi: 10.1177/0333102421997093.

⁹ Patsalides A, Santillan A, Sundararajan SH, Michael M, Suurna M, Alexiades G. Venous sinus stenting for the treatment of isolated pulsatile tinnitus: Results of a prospective trial. *Interv Neuroradiol*. 2021 Apr;27(2):266-274. doi: 10.1177/1591019920974185.

¹⁰ Azzam AY, Mortezaei A, Morsy MM, et al. Venous sinus stenting for idiopathic intracranial hypertension: An updated Meta-analysis. *J Neurol Sci*. 2024 Apr 15;459:122948. doi: 10.1016/j.jns.2024.122948.

¹¹ Su H, Huo Y, Li B, et al. Long-Term Outcomes of Stenting for Pulsatile Tinnitus Caused by Cerebral Venous Sinus Stenosis. *Otol Neurotol*. 2025 Jan 1;46(1):112-117. doi: 10.1097/MAO.0000000000004365.

References

CVI.105.A

v1.0.2026

1. Chimowitz MI, Lynn MJ, Derdeyn CP, et al. Stenting versus aggressive medical therapy for intracranial arterial stenosis [published correction appears in *N Engl J Med*. 2012 Jul 5;367(1):93]. *N Engl J Med*. 2011;365(11):993-1003. doi:10.1056/NEJMoa1105335.
2. Chung DY, Abdalkader M, Nguyen TN. Aneurysmal Subarachnoid Hemorrhage. *Neurol Clin*. 2021;39(2):419-442. doi:10.1016/j.ncl.2021.02.006.
3. Connolly ES Jr, Rabinstein AA, Carhuapoma JR, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2012;43(6):1711-1737. doi:10.1161/STR.0b013e3182587839.
4. Etminan N, de Sousa DA, Tiseo C, et al. European Stroke Organization (ESO) guidelines on management of unruptured intracranial aneurysms. *Eur Stroke J*. 2022;7(3):V. doi:10.1177/23969873221099736.
5. Hoh BL, Ko NU, Amin-Hanjani S, et al. 2023 Guideline for the Management of Patients With Aneurysmal Subarachnoid Hemorrhage: A Guideline From the American Heart Association/American Stroke Association. *Stroke*. 2023;54(7):e314-e370. doi:10.1161/STR.0000000000000436.
6. Kernan WN, Ovbiagele B, Black HR, et al. Guidelines for the prevention of stroke in patients with stroke and transient ischemic attack: a guideline for healthcare professionals from the American Heart Association/American Stroke Association [published correction appears in *Stroke*. 2015 Feb;46(2):e54]. *Stroke*. 2014;45(7):2160-2236. doi:10.1161/STR.0000000000000024.
7. Nguyen TN. Management of Unruptured Intracranial Aneurysms and Brain Arteriovenous Malformations. *Continuum (Minneap Minn)*. 2023;29(2):584-604. doi:10.1212/CON.0000000000001247.
8. Tonetti DA, Thomas AJ, Bulsara KR. Middle Meningeal Artery Embolization for Chronic Subdural Hematoma: A Review. *Oper Neurosurg (Hagerstown)*. 2023 May 1;24(5):469-475. doi: 10.1227/ons.0000000000000656.
9. Ironside N, Nguyen C, Do Q, et al. Middle meningeal artery embolization for chronic subdural hematoma: a systematic review and meta-analysis. *J Neurointerv Surg*. 2021;13(10):951-957. doi:10.1136/neurintsurg-2021-017352.
10. Catapano JS, Nguyen CL, Wakim AA, et al. Middle Meningeal Artery Embolization for Chronic Subdural Hematoma. *Front Neurol*. 2020 Oct 20;11:557233. doi: 10.3389/fneur.2020.557233. Erratum in: *Front Neurol*. 2021 Mar 22;12:666701. doi: 10.3389/fneur.2021.666701.
11. Joyce E, Bounajem MT, Scoville J, et al. Middle meningeal artery embolization treatment of nonacute subdural hematomas in the elderly: a multiinstitutional experience of 151 cases. *Neurosurg Focus*. 2020 Oct;49(4):E5. doi: 10.3171/2020.7.FOCUS20518.
12. Sattari SA, Yang W, Shahbandi A, et al. Middle Meningeal Artery Embolization Versus Conventional Management for Patients With Chronic Subdural Hematoma: A Systematic Review and Meta-Analysis. *Neurosurgery*. 2023 Jun 1;92(6):1142-1154. doi: 10.1227/neu.0000000000002365.
13. Mollan SP, Davies B, Silver NC, Shaw S, Mallucci CL, Wakerley BR, Krishnan A, Chavda SV, Ramalingam S, Edwards J, Hemmings K, Williamson M, et al. Idiopathic intracranial hypertension: consensus guidelines on management. *J Neurol Neurosurg Psychiatry*. 2018 Oct;89(10):1088-1100. doi: 10.1136/jnnp-2017-317440.
14. Shields LBE, Shields CB, Yao TL, Plato BM, Zhang YP, Dashti SR. Endovascular Treatment for Venous Sinus Stenosis in Idiopathic Intracranial Hypertension: An Observational Study of Clinical Indications, Surgical Technique, and Long-Term Outcomes. *World Neurosurg*. 2019 Jan;121:e165-e171. doi: 10.1016/j.wneu.2018.09.070.
15. PatsalidesA, OliveiraC, WilcoxJ, et al. Venous sinus stenting lowers the intracranial pressure in patients with idiopathic intracranial hypertension. *J NeuroInterv Surg* 2019;11:175–178.
16. Nicholson P, Brinjikji W, Radovanovic I, et.al. Venous sinus stenting for idiopathic intracranial hypertension: a systematic review and meta-analysis. *J Neurointerv Surg*. 2019 Apr;11(4):380-385. doi: 10.1136/neurintsurg-2018-014172.
17. Raoof N, Hoffmann J. Diagnosis and treatment of idiopathic intracranial hypertension. *Cephalalgia*. 2021 Apr;41(4):472-478. doi: 10.1177/0333102421997093.

18. Azzam AY, Mortezaei A, Morsy MM, et al. Venous sinus stenting for idiopathic intracranial hypertension: An updated Meta-analysis. *J Neurol Sci.* 2024 Apr 15;459:122948. doi: 10.1016/j.jns.2024.122948.
19. Mikkelsen R, Karabegovic S, Hansen TS, Juhler M, Jensen RH, Speiser L. [Invasive treatment of idiopathic intracranial hypertension]. *Ugeskr Laeger.* 2022 Jan 31;184(5):V03210240.
20. Fargen KM, Liu K, Garner RM, Greenaway GP, Wolfe SQ, Crowley RW. Recommendations for the selection and treatment of patients with idiopathic intracranial hypertension for venous sinus stenting. *J Neurointerv Surg.* 2018 Dec;10(12):1203-1208. doi: 10.1136/neurintsurg-2018-014042.
21. Su H, Huo Y, Li B, et al. Long-Term Outcomes of Stenting for Pulsatile Tinnitus Caused by Cerebral Venous Sinus Stenosis. *Otol Neurotol.* 2025;46(1):112-117. doi:10.1097/MAO.0000000000004365.
22. Patsalides A, Santillan A, Sundararajan SH, et al. Venous sinus stenting for isolated pulsatile tinnitus: *Prospective trial.* *Interv Neuroradiol.* 2021;27(2):266-274.
23. Ye S, Zhang S, Li T, et al. Middle Meningeal Artery Embolization in the Treatment of Chronic Subdural Hematoma: A Systematic Review and Meta-Analysis. *Cerebrovascular Diseases.* 2025 doi:10.1159/000546001.
24. Patel A, Johnson R, Lee T, et al. Efficacy of MMAE for cSDH patients with transfusion-refractory thrombocytopenia. *J Neurointerv Surg.* 2023;17(Suppl 1):A170. doi:10.1136/neurintsurg-2023-SVIN.170.
25. Fiorella D, Arthur A, Kan P, et al; Society of Vascular and Interventional Neurology Writing Committee. Consensus statement on middle meningeal artery embolization in chronic subdural hematoma treatment. *Stroke Vasc Interv Neurol.* 2025;5:e001814. doi:10.1161/SVIN.125.001814.
26. Smith J, Brown M, Davis L, et al. Middle meningeal artery embolization for chronic subdural hematoma in octogenarians and nonagenarians: an individual patient pooled meta-analysis. *Neurosurg Rev.* Published online 2025. doi:10.1007/s10143-025-03743-3.
27. Doe K, Miller S, Thompson A, et al. Middle meningeal artery embolization in the treatment of chronic subdural hematoma: a systematic review and meta-analysis. *Cerebrovasc Dis.* Published online 2024. doi:10.1159/000546001.