

CIGNA MEDICAL COVERAGE POLICIES - MUSCULOSKELETAL CMM-315: Shoulder Surgery - Arthroscopic and Open Procedures

Effective Date: August 04, 2026



Instructions for use

The following coverage policy applies to health benefit plans administered by Cigna. Coverage policies are intended to provide guidance in interpreting certain standard Cigna benefit plans and are used by medical directors and other health care professionals in making medical necessity and other coverage determinations. Please note the terms of a customer's particular benefit plan document may differ significantly from the standard benefit plans upon which these coverage policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a coverage policy.

In the event of a conflict, a customer's benefit plan document always supersedes the information in the coverage policy. In the absence of federal or state coverage mandates, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of:

1. The terms of the applicable benefit plan document in effect on the date of service
2. Any applicable laws and regulations
3. Any relevant collateral source materials including coverage policies
4. The specific facts of the particular situation

Coverage policies relate exclusively to the administration of health benefit plans. Coverage policies are not recommendations for treatment and should never be used as treatment guidelines.

This evidence-based medical coverage policy has been developed by EviCore, Inc. Some information in this coverage policy may not apply to all benefit plans administered by Cigna.

These guidelines include procedures EviCore does not review for Cigna. Please refer to the [Cigna CPT code list](#) for the current list of high-tech imaging procedures that EviCore reviews for Cigna.

CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association (AMA). CPT® five-digit codes, nomenclature and other data are copyright 2026 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in the CPT book. AMA does not directly or indirectly practice medicine or dispense medical services. AMA assumes no liability for the data contained herein or not contained herein.

Table of Contents

Guideline	Page
Definitions.....	3
General Guidelines.....	8
Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body.....	10
Diagnostic Arthroscopy.....	12
Loose Body/Foreign Body Removal.....	15
Synovectomy.....	18
Debridement.....	21
Rotator Cuff Repair.....	25
Distal Clavicle Excision.....	29
Subacromial Decompression/Acromioplasty.....	32
Labral Repair.....	35
Biceps Tenodesis.....	38
Arthroscopic or Open Surgical Procedures for Shoulder Instability and/or Laxity.....	41
Arthroscopic Capsular Release.....	44
Lysis of Adhesions.....	47
Manipulation Under Anesthesia (MUA).....	50
Arthroscopic or Open Coracoplasty/Subcoracoid Decompression.....	53
Codes (CMM-315).....	55
References (CMM-315).....	59

Definitions

Guideline	Page
Definitions.....	4

Definitions

CMM.JT.DF.315

v1.0.2026

**Adhesive Capsulitis/
Arthrofibrosis**

a condition of the shoulder characterized by stiffness, loss of motion (contracture), and pain due to scarring in and/or around the shoulder joint. Conditions that have been suggested to predispose an individual to adhesive capsulitis include trauma, surgery to the shoulder, inflammatory diseases, diabetes, hyperthyroidism, and dyslipidemia. Often called frozen shoulder, adhesive capsulitis is clinically divided into classes:

- Primary adhesive capsulitis is characterized by a significant limitation of both active and passive motions of the shoulder; individuals are typically unable to recall a possible cause of the condition (idiopathic adhesive capsulitis).
- Secondary adhesive capsulitis is characterized by a trauma or a possible cause prior to the onset of the symptoms, such as fracture of the humerus, rotator cuff repair, shoulder girdle injury/surgery, or prolonged immobilization.

**Distal Clavicle
Excision**

the removal of the end of the clavicle at the acromioclavicular (AC) joint. The superior AC ligament remains intact so that the joint remains stable.

**Impingement
Syndrome**

a condition that commonly results from friction, abrasion, and inflammation of the rotator cuff and the long head of the biceps tendon with the subacromial arch (anterior lip of the acromion, coracoacromial ligament, and acromioclavicular joint) from acute trauma, repetitive use, or degenerative changes.

Labral Tear

a tear of the glenoid labrum. Tears are typically classified by the position of the tear in relation to the glenoid.

- Bankart tear: a tear in the labrum located in the front, lower (anterior, inferior) part of the glenoid. This type of tear occurs most commonly during a shoulder dislocation and makes the shoulder more prone to recurrent dislocations.
- SLAP tear (Superior Labral, Anterior, and Posterior tear): a tear in the labrum that covers the top part of the glenoid from the front to the back. A SLAP tear occurs at the point where the long head of the biceps tendon attaches. This type of tear occurs most commonly during falls on an outstretched arm.

Non-Surgical Management (with regard to the treatment of shoulder pain)

any provider-directed non-surgical treatment, which has been demonstrated in the scientific literature as efficacious and/or is considered reasonable care in the treatment of shoulder pain. The types of treatment involved can include, but are not limited to, the following: relative rest/activity modification; supervised physiotherapy modalities and therapeutic exercises; prescription and non-prescription medications; assistive devices; and/or injections.

Rotator Cuff Arthropathy

a condition that results from ALL of the following:

- rotator cuff dysfunction (e.g., secondary to unrepairable/irreparable massive rotator cuff tear)
- advanced glenohumeral arthritis
- radiographically diminished acromio-humeral distance

Rotator Cuff Tear (RCT)

a disruption of the tendon(s) of the rotator cuff muscles which attach the humerus to the scapula. These tendons are important in shoulder movements and maintaining glenohumeral joint stability. The supraspinatus tendon is most commonly involved, but the infraspinatus, teres minor, and subscapularis tendons can also be torn.

- Defining whether a rotator cuff tear is acute has relevance to treatment. In evaluating individuals, the surgeon should attempt to properly identify individuals with acute tears as opposed to individuals with pre-existing chronic tears that become symptomatic after an injury event. A discrete traumatic event is more suggestive of acute tear. Physical exam findings including supraspinatus and infraspinatus muscle atrophy, as well as internal and external rotation lag signs, may be indicative of larger and more chronic rotator cuff tears.
- Evaluation of rotator cuff muscle quality with CT or MRI is an important consideration. Chronic and larger tears are associated with muscle atrophy and fatty replacement, both of which correlate with inferior functional outcome after rotator cuff repair. It is thought that early repair of acute rotator cuff tears might mitigate the development of chronic tendon and muscle pathology and improve functional outcomes.
- Classification of rotator cuff tears (based upon surgical findings):
 - Partial-thickness tears, also called incomplete tears (Ellman):
 - Grade 1: <3mm deep (<25% thickness)
 - Grade 2: 3-6mm in depth, but not exceeding 50% of the tendon thickness
 - Grade 3: >6mm deep (>50% thickness)
 - Full-thickness tears, also called complete tears (Cofield):
 - Small: <1cm
 - Medium: 1-3cm
 - Large: 3-5cm
 - Massive: >5cm

Shoulder Dislocation	the complete loss of the humeral articulation with the glenoid fossa, usually as a result of acute trauma.
Shoulder Instability and/or Laxity	<p>a partial loss of the glenohumeral articulation of which there are two categories:</p> <ul style="list-style-type: none">• Post-traumatic shoulder instability includes stretched or torn ligaments of the shoulder as a result of a previous injury.• Atraumatic instability and/or laxity includes generalized looseness of the joints, "double-jointed", or "multi-directional instability", usually representing a type of congenital ligamentous laxity.
Shoulder Subluxation	a partial loss of humeral articulation with the glenoid fossa (incomplete or partial dislocation), usually as a result of repetitive trauma to the degree that symptoms are produced.
Subacromial Decompression	the removal of bone or other abnormality to enlarge the space between the rotator cuff musculature and the acromion.
Acromioplasty	the removal of bone from the acromion and partial resection of the coracoacromial ligament.

General Guidelines

Guideline	Page
General Guidelines.....	9

Shoulder Surgery - Arthroscopic and Open Procedures

General Guidelines

CMM.JT.GG.315

v1.0.2026

Application of Guideline

- The determination of medical necessity for the performance of shoulder surgery is always made on a case-by-case basis.

Health Equity Considerations

Health equity is the highest level of health for all individuals; health inequity is the avoidable difference in health status or distribution of health resources due to the social conditions in which individuals are born, grow, live, work, and age. Social determinants of health are the conditions in the environment that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples include the following: safe housing, transportation, and neighborhoods; racism, discrimination, and violence; education, job opportunities, and income; access to nutritious foods and physical activity opportunities; access to clean air and water; and language and literacy skills.

Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body

Shoulder Surgery - Arthroscopic and Open Procedures

Guideline	Page
Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body.....	11

Arthroscopic or Open Procedures for Fracture, Tumor, Infection, or Foreign Body

CMM.JT.IN.315

v1.0.2026

- Arthroscopic or open shoulder surgery may be considered **medically necessary** when surgery is being performed for fracture, tumor, infection, or foreign body that has led to, or will likely lead to, progressive destruction.

Shoulder Surgery - Arthroscopic and Open Procedures

Diagnostic Arthroscopy

Guideline	Page
Diagnostic Arthroscopy Indications.....	13
Diagnostic Arthroscopy Non-Indications.....	14

Diagnostic Arthroscopy Indications

CMM.JT.IN.315

v1.0.2026

Diagnostic arthroscopy is considered **medically necessary** as a stand-alone procedure when ALL of the following criteria have been met:

- MRI or CT is inconclusive for internal derangement/pathology
- Physical exam demonstrates ALL of the following findings (as compared to the non-involved side):
 - functionally-limited range of motion (active or passive)
 - measurable loss of strength
 - positive Neer impingement test or Hawkins-Kennedy impingement test
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment) for at least six (6) months' duration
- Failure of provider-directed non-surgical management for at least three (3) months' duration
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Diagnostic Arthroscopy Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Diagnostic arthroscopy is considered **not medically necessary** for ANY other indication or condition.
- Based on lack of scientific evidence of efficacy and safety, "in-office" diagnostic arthroscopy (e.g., Mi-Eye™, VisionScope®) is considered **not medically necessary**.

Loose Body/Foreign Body Removal

Shoulder Surgery - Arthroscopic and Open Procedures

Guideline	Page
Loose Body/Foreign Body Removal Indications.....	16
Loose Body/Foreign Body Removal Non-Indications.....	17

Loose Body/Foreign Body Removal Indications

CMM.JT.IN.315

v1.0.2026

Loose body or foreign body removal is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT shows findings of a loose body or foreign body within the shoulder joint
- Symptoms include BOTH of the following:
 - function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
 - mechanical symptoms including painful locking, clicking, catching, or popping
- Failure of provider-directed non-surgical management for at least three (3) months' duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required if the loose body or foreign body has caused an acute restriction of shoulder joint range of motion (i.e., locking).
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Loose Body/Foreign Body Removal Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Loose body or foreign body removal is considered **not medically necessary** for ANY other indication or condition.

Shoulder Surgery - Arthroscopic and Open Procedures

Synovectomy

Guideline	Page
Synovectomy Indications.....	19
Synovectomy Non-Indications.....	20

Synovectomy Indications

CMM.JT.IN.315

v1.0.2026

Synovectomy (partial or complete) is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT arthrogram shows findings of synovitis
- Presence of ANY of the following conditions:
 - inflammatory arthritis (i.e., rheumatoid arthritis, gout, pseudogout, psoriatic arthritis)
 - pigmented villonodular synovitis (PVNS)
 - synovial chondromatosis
 - Lyme synovitis
 - hemophilia
 - hemochromatosis
 - non-specific synovitis (e.g., proliferative synovitis, post-operative synovitis as a sequela from a shoulder replacement, etc.)
 - recurrent hemarthrosis
- Physical exam demonstrates functionally-limited range of motion (active or passive) (as compared to the non-involved side)
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
- Failure of provider-directed non-surgical management for at least three (3) months' duration
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Synovectomy Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Synovectomy is considered **not medically necessary** for ANY other indication or condition.

Debridement

Guideline	Page
Debridement Indications.....	22
Debridement Non-Indications.....	24

Debridement Indications

CMM.JT.IN.315

v1.0.2026

Debridement (limited or extensive) is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT shows findings of underlying pathology which correlates with the individual's reported symptoms and physical exam findings
- Physical exam demonstrates BOTH of the following findings (as compared to the non-involved side):
 - EITHER of the following findings:
 - functionally-limited range of motion
 - measurable loss of strength
 - ANY of the following positive orthopedic tests/signs:
 - drop arm test
 - painful arc test
 - Jobe test or empty can test
 - external rotation lag sign (dropping sign)
 - lift-off test
 - belly press test (Napoleon)
 - cross body adduction test
 - resisted AC joint extension test
 - Neer impingement test
 - Hawkins-Kennedy impingement test
 - O'Brien's test
 - biceps load test
 - clunk test
 - anterior slide test
 - compression rotation test
 - Speed's test
 - upper cut test
 - Yergason's test
 - proximal biceps (groove) tenderness
 - modified dynamic labral shear
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)

- Failure of provider-directed non-surgical management for at least three (3) months' duration
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Debridement Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Debridement is considered **not medically necessary** for ANY other indication or condition.

Rotator Cuff Repair

Guideline	Page
Rotator Cuff Repair Indications.....	26
Rotator Cuff Repair Non-Indications.....	28

Shoulder Surgery - Arthroscopic and Open Procedures

Rotator Cuff Repair Indications

CMM.JT.IN.315

v1.0.2026

Rotator cuff repair is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT shows EITHER of the following findings:
 - Grade 2 or 3 partial-thickness rotator cuff tear (Ellman classification)
 - full-thickness rotator cuff tear (Cofield classification)
- Physical exam demonstrates BOTH of the following findings (as compared to the non-involved side):
 - EITHER of the following findings:
 - functionally-limited range of motion
 - measurable loss of strength of the rotator cuff musculature
 - ANY of the following positive orthopedic tests/signs:
 - drop arm test
 - painful arc test
 - Jobe test or empty can test
 - lateral Jobe test
 - external rotation lag sign (dropping sign)
 - external rotation resistance test
 - full can test
 - hug up test
 - internal rotation lag sign
 - internal resistance rotation test (IRRT)
 - internal rotation resistance test at maximal 90° of abduction and maximal external rotation test (IRRTM)
 - lift-off test
 - bear hug test
 - belly press test (Napoleon)
 - belly-off test
 - Neer impingement test
 - Hawkins-Kennedy impingement test
 - Hornblower test (Patte)
 - Yocum test
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)

- Failure of provider-directed non-surgical management for at least three (3) months' duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for an individual who suffers a discrete traumatic event that results in an acute full-thickness rotator cuff tear without evidence of a pre-existing chronic rotator cuff tear.
 - **Note:** The presence of fatty infiltration and/or muscle atrophy on MRI or CT is considered evidence of a pre-existing chronic rotator cuff tear.
 - Therefore, when fatty infiltration and/or muscle atrophy is also present on MRI or CT, three (3) months of provider-directed non-surgical management is required, regardless of whether a discrete traumatic event occurred.
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Rotator Cuff Repair Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Rotator cuff repair is considered **not medically necessary** for ANY other indication or condition.

Experimental, Investigational, or Unproven (EIU)

- Based on lack of scientific evidence of efficacy and safety, the following is considered **experimental, investigational, or unproven**:
 - superior capsular reconstruction using any type of graft (e.g., autograft, allograft, xenograft, or synthetic)

Distal Clavicle Excision

Guideline	Page
Distal Clavicle Excision Indications.....	30
Distal Clavicle Excision Non-Indications.....	31

Distal Clavicle Excision Indications

CMM.JT.IN.315

v1.0.2026

Distal clavicle excision is considered **medically necessary** when ALL of the following criteria have been met:

- Imaging shows BOTH of the following findings:
 - plain radiographs show findings of underlying pathology in the subacromial space and/or at the AC joint (e.g., AC joint arthritis)
 - MRI or CT shows findings of underlying pathology (e.g., AC joint arthritis, impingement, etc.) which correlates with the individual's reported symptoms and physical exam findings
 - **Criteria exception:** Advanced imaging is not required for an isolated distal clavicle excision when the procedure is not associated with subacromial decompression/acromioplasty surgery.
- Physical exam demonstrates BOTH of the following findings (as compared to the non-involved side):
 - localized tenderness to palpation of the acromioclavicular (AC) joint
 - ANY of the following positive orthopedic tests/signs:
 - cross body adduction test
 - resisted AC joint extension test
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
- Failure of provider-directed non-surgical management for at least three (3) months' duration
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Distal Clavicle Excision Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Distal clavicle excision is considered **not medically necessary** for ANY other indication or condition.

Subacromial Decompression/ Acromioplasty

Shoulder Surgery - Arthroscopic and Open Procedures

Guideline	Page
Subacromial Decompression/Acromioplasty Indications.....	33
Subacromial Decompression/Acromioplasty Non-Indications.....	34

Subacromial Decompression/ Acromioplasty Indications

CMM.JT.IN.315

v1.0.2026

Subacromial decompression/acromioplasty is considered **medically necessary** when ALL of the following criteria have been met:

- When requested as an arthroscopic procedure (add-on procedure), it is required to be performed with other medically necessary primary arthroscopic shoulder surgical procedures (with the exception of diagnostic shoulder arthroscopy)
- Imaging shows BOTH of the following findings:
 - plain radiographs show findings of underlying pathology in the subacromial space and/or at the AC joint (e.g., AC joint arthritis, Type 3 hooked acromion, down-sloping acromion, etc.)
 - MRI or CT shows findings of underlying pathology (e.g., AC joint arthritis, impingement, etc.) which correlates with the individual's reported symptoms and physical exam findings
- Physical exam demonstrates ANY of the following positive orthopedic tests/signs (as compared to the non-involved side):
 - Neer impingement test
 - Hawkins-Kennedy impingement test
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
- Failure of provider-directed non-surgical management for at least three (3) months' duration
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Subacromial Decompression/ Acromioplasty Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Subacromial decompression/acromioplasty is considered **not medically necessary** for ANY other indication or condition.

Shoulder Surgery - Arthroscopic and Open Procedures

Labral Repair

Guideline	Page
Labral Repair Indications.....	36
Labral Repair Non-Indications.....	37

Labral Repair Indications

CMM.JT.IN.315

v1.0.2026

Labral repair is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT shows labral tear (e.g., SLAP) and correlates with the individual's reported symptoms and physical exam findings
- Physical exam demonstrates BOTH of the following findings (as compared to the non-involved side):
 - minimally limited or full shoulder range of motion
 - ANY of the following positive orthopedic tests/signs:
 - O'Brien's test
 - biceps load test
 - clunk test
 - anterior slide test
 - compression rotation test
 - Speed's test
 - modified dynamic labral shear
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
- Failure of provider-directed non-surgical management for at least three (3) months' duration
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Labral Repair Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Labral repair is considered **not medically necessary** for ANY other indication or condition.

Biceps Tenodesis

Guideline	Page
Biceps Tenodesis Indications.....	39
Biceps Tenodesis Non-Indications.....	40

Biceps Tenodesis Indications

CMM.JT.IN.315

v1.0.2026

Biceps tenodesis is considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT shows labral tear/biceps tendon pathology (e.g., SLAP or full-thickness subscapularis tear) and correlates with the individual's reported symptoms and physical exam findings
- Physical exam demonstrates BOTH of the following findings (as compared to the non-involved side):
 - minimally limited or full shoulder range of motion
 - ANY of the following positive orthopedic tests/signs:
 - O'Brien's test
 - biceps load test
 - clunk test
 - anterior slide test
 - compression rotation test
 - Speed's test
 - upper cut test
 - Yergason's test
 - proximal biceps (groove) tenderness
 - modified dynamic labral shear
 - Popeye sign
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
- Failure of provider-directed non-surgical management for at least three (3) months' duration
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required for an individual who suffers a discrete traumatic event that results in an acute proximal biceps tendon rupture.
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Biceps Tenodesis Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Biceps tenodesis is considered **not medically necessary** for ANY other indication or condition.

Arthroscopic or Open Surgical Procedures for Shoulder Instability and/or Laxity

Shoulder Surgery - Arthroscopic and Open Procedures

Guideline	Page
Arthroscopic or Open Surgical Procedures for Shoulder Instability and/or Laxity Indications.....	42
Arthroscopic or Open Surgical Procedures for Shoulder Instability and/or Laxity Non-Indications.....	43

Arthroscopic or Open Surgical Procedures for Shoulder Instability and/or Laxity Indications

CMM.JT.IN.315
v1.0.2026

Arthroscopic or open surgical procedures for shoulder instability and/or laxity are considered **medically necessary** when ALL of the following criteria have been met:

- MRI or CT shows labral/capsular/bone pathology (e.g., Bankart, capsular tear, Hill-Sachs deformity) and correlates with the individual's reported symptoms and physical exam findings
- Physical exam demonstrates ANY of the following positive orthopedic tests/signs (as compared to the non-involved side):
 - anterior apprehension test
 - posterior apprehension test
 - sulcus sign
 - load and shift test
- Documented history of post-traumatic or atraumatic instability and/or laxity that has resulted in function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment)
- Failure of provider-directed non-surgical management for at least three (3) months' duration that includes shoulder stabilization/strengthening exercises
 - **Criteria exception:** Three (3) months of provider-directed non-surgical management is not required in an acute traumatic injury setting when EITHER of the following conditions are present:
 - irreducible shoulder dislocation
 - anterior shoulder instability in competitive contact or collision athletes
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Arthroscopic or Open Surgical Procedures for Shoulder Instability and/ or Laxity Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Arthroscopic or open surgical procedures for shoulder instability and/or laxity are considered **not medically necessary** for ANY other indication or condition.

Experimental, Investigational, or Unproven (EIU)

- Based on lack of scientific evidence of efficacy and safety, the following is considered **experimental, investigational, or unproven**:
 - superior capsular reconstruction using any type of graft (e.g., autograft, allograft, xenograft, or synthetic)

Arthroscopic Capsular Release

Guideline	Page
Arthroscopic Capsular Release Indications.....	45
Arthroscopic Capsular Release Non-Indications.....	46

Shoulder Surgery - Arthroscopic and Open Procedures

Arthroscopic Capsular Release Indications

CMM.JT.IN.315

v1.0.2026

Arthroscopic capsular release is considered **medically necessary** when ALL of the following criteria have been met:

- Performed for chronic refractory adhesive capsulitis or arthrofibrosis
- Physical exam demonstrates functionally-limited and painful global loss of active and passive range of motion of at least 50% (as compared to the non-involved side)
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment) for at least six (6) months' duration
- Failure of provider-directed non-surgical management for at least three (3) months' duration, including a combination of anti-inflammatory medication, cortisone injection, and a trial of physical therapy (i.e., active exercise and manual therapy designed to increase joint mobility and range of motion)
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Arthroscopic Capsular Release Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Arthroscopic capsular release is considered **not medically necessary** for ANY other indication or condition.

Shoulder Surgery - Arthroscopic and Open Procedures

Lysis of Adhesions

Guideline	Page
Lysis of Adhesions Indications.....	48
Lysis of Adhesions Non-Indications.....	49

Lysis of Adhesions Indications

CMM.JT.IN.315

v1.0.2026

Lysis of adhesions is considered **medically necessary** when ALL of the following criteria have been met:

- Performed for chronic refractory adhesive capsulitis or arthrofibrosis
- Physical exam demonstrates functionally-limited and painful global loss of active and passive range of motion of at least 50% (as compared to the non-involved side)
- Symptoms include function-limiting pain (e.g., loss of shoulder function which interferes with the ability to carry out age-appropriate activities of daily living and/or demands of employment) for at least six (6) months' duration
- Failure of provider-directed non-surgical management for at least three (3) months' duration, including a combination of anti-inflammatory medication, cortisone injection, and a trial of physical therapy (i.e., active exercise and manual therapy designed to increase joint mobility and range of motion)
- Other potential pathological conditions have been excluded, including, but not limited to the following: fracture, thoracic outlet syndrome, brachial plexus disorders, referred neck pain, cervical radiculopathy, and advanced glenohumeral osteoarthritis.

Lysis of Adhesions Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Lysis of adhesions is considered **not medically necessary** for ANY other indication or condition.

Manipulation Under Anesthesia (MUA)

Guideline	Page
Manipulation Under Anesthesia (MUA) Indications.....	51
Manipulation Under Anesthesia (MUA) Non-Indications.....	52

Shoulder Surgery - Arthroscopic and Open Procedures

Manipulation Under Anesthesia (MUA) Indications

CMM.JT.IN.315

v1.0.2026

- See Manipulation Under Anesthesia (MUA) Non-Indications.

Manipulation Under Anesthesia (MUA) Non-Indications

CMM.JT.NI.315

v1.0.2026

Not Medically Necessary

- Manipulation under anesthesia (MUA) of a shoulder joint is included in all arthroscopic shoulder procedures and is therefore considered incidental to the base procedure requiring medical necessity review.

Arthroscopic or Open Coracoplasty/Subcoracoid Decompression

Shoulder Surgery - Arthroscopic and Open Procedures

Guideline	Page
-----------	------

Arthroscopic or Open Coracoplasty/Subcoracoid Decompression Non-Indications.....	54
--	----

Arthroscopic or Open Coracoplasty/ Subcoracoid Decompression Non- Indications

CMM.JT.NI.315

v1.0.2026

Experimental, Investigational, or Unproven (EIU)

- Based on lack of scientific evidence of efficacy and safety, arthroscopic or open coracoplasty/subcoracoid decompression for the treatment of subcoracoid impingement is considered **experimental, investigational, or unproven**.

Codes (CMM-315)

Guideline	Page
Codes (CMM-315).....	56

Codes (CMM-315)

CMM.JT.PC.315

v1.0.2026

The inclusion of any code in this table does not imply that the code is under management or requires prior authorization. Refer to the applicable health plan for management details. Prior authorization of a code listed in this table is not a guarantee of payment. The Certificate of Coverage or Evidence of Coverage policy outlines the terms and conditions of the member's health insurance policy.

Code	Code Description/Definition
23000	Removal of subdeltoid calcareous deposits, open
23020	Capsular contracture release (e.g. Sever type procedure)
23106	Arthrotomy; sternoclavicular joint, with synovectomy, with or without biopsy
23120	Claviclectomy; partial
23130	Acromioplasty or acromionectomy, partial, with or without coracoacromial ligament release
23145	Excision or curettage of bone cyst or benign tumor of clavicle or scapula; with autograft (includes obtaining graft)
23155	Excision or curettage of bone cyst or benign tumor of proximal humerus; with autograft (includes obtaining graft)
23172	Sequestrectomy (e.g. for osteomyelitis or bone abscess), scapula
23174	Sequestrectomy (e.g. for osteomyelitis or bone abscess), humeral head to surgical neck
23410	Repair of ruptured musculotendinous cuff (e.g., rotator cuff) open; acute
23412	Repair of ruptured musculotendinous cuff (e.g., rotator cuff) open; chronic
23415	Coracoacromial ligament release, with or without acromioplasty
23420	Reconstruction of complete shoulder (rotator) cuff avulsion, chronic (includes acromioplasty)

Shoulder Surgery - Arthroscopic and Open Procedures

Code	Code Description/Definition
23430	Tenodesis of long tendon of biceps
23440	Resection or transplantation of long tendon of biceps
23450	Capsulorrhaphy, anterior; Putti-Platt procedure or Magnuson type operation
23455	Capsulorrhaphy, anterior; with labral repair (e.g., Bankart procedure)
23460	Capsulorrhaphy, anterior, any type; with bone block
23462	Capsulorrhaphy, anterior, any type; with coracoid process transfer
23465	Capsulorrhaphy, glenohumeral joint, posterior, with or without bone block
23466	Capsulorrhaphy, glenohumeral joint, any type multi-directional instability
23700	Manipulation under anesthesia, shoulder joint, including application of fixation apparatus (dislocation excluded)
29805	Arthroscopy, shoulder, diagnostic, with or without synovial biopsy (separate procedure)
29806	Arthroscopy, shoulder, surgical; capsulorrhaphy
29807	Arthroscopy, shoulder, surgical; repair of SLAP lesion
29819	Arthroscopy, shoulder, surgical; with removal of loose body or foreign body
29820	Arthroscopy, shoulder, surgical; synovectomy, partial
29821	Arthroscopy, shoulder, surgical; synovectomy, complete
29822	Arthroscopy, shoulder, surgical; debridement, limited, 1 or 2 discrete structures (e.g., humeral bone, humeral articular cartilage, glenoid bone, glenoid articular cartilage, biceps tendon, biceps anchor complex, labrum, articular capsule, articular side of the rotator cuff, bursal side of the rotator cuff, subacromial bursa, foreign body[ies])

Code	Code Description/Definition
29823	Arthroscopy, shoulder, surgical; debridement, extensive, 3 or more discrete structures (e.g., humeral bone, humeral articular cartilage, glenoid bone, glenoid articular cartilage, biceps tendon, biceps anchor complex, labrum, articular capsule, articular side of the rotator cuff, bursal side of the rotator cuff, subacromial bursa, foreign body[ies])
29824	Arthroscopy, shoulder, surgical; distal claviclectomy including distal articular surface (Mumford procedure)
29825	Arthroscopy, shoulder, surgical; with lysis and resection of adhesions, with or without manipulation
29826	Arthroscopy, shoulder, surgical; decompression of subacromial space with partial acromioplasty, with coracoacromial ligament (i.e., arch) release when performed (List separately in addition to code for primary procedure)
29827	Arthroscopy, shoulder, surgical; with rotator cuff repair
29828	Arthroscopy, shoulder, surgical; biceps tenodesis

References (CMM-315)

Guideline	Page
References (CMM-315).....	60

References (CMM-315)

CMM.JT.RF.315

v1.0.2026

1. Abdel Khalik H, Lameire DL, Leroux T, Bhandari M, Khan M. Arthroscopic stabilization surgery for first-time anterior shoulder dislocations: a systematic review and meta-analysis. *J Shoulder Elbow Surg.* 2024;33(8):1858-1872. doi:10.1016/j.jse.2024.01.037.
2. Ahmed AF, Polisetty TS, Wang C, et al. Higher return to sport and lower revision rates when performing arthroscopic Bankart repair with remplissage for anterior shoulder instability with a Hill-Sachs lesion: a meta-analysis. *J Shoulder Elbow Surg.* 2024;33(8):1836-1846.
3. Altintas B, Anderson N, Pitta R, et al. Repair of Rotator Cuff Tears in the Elderly: Does It Make Sense? A Systematic Review. *Am J Sports Med.* 2019;48(3):744-753. doi:10.1177/0363546519834574.
4. American Academy of Orthopaedic Surgeons (AAOS); AAOS Clinical Practice Guidelines Unit. *Guideline and Evidence Report: Optimizing the Management of Rotator Cuff Problems.* Dec 2010. Rosemont, IL. © American Academy of Orthopaedic Surgeons (AAOS).
5. American Academy of Orthopaedic Surgeons (AAOS). Evidence-Based Clinical Practice Guideline: *Management of Rotator Cuff Injuries.* Aug 2025. Rosemont, IL. © American Academy of Orthopaedic Surgeons (AAOS).
6. Arciero RA, Wheeler JH, Ryan JB, et al. Arthroscopic Bankart Repair versus Nonoperative Treatment for Acute, Initial Anterior Shoulder Dislocations. *Am J Sports Med.* 1994;22(5):589-594.
7. Audenaert E, Nuffel JV, Schepens A, Verhelst M, Verdonk R. Reconstruction of massive rotator cuff lesions with a synthetic interposition graft: a prospective study of 41 patients. *Knee Surg Sports Traumatol Arthrosc.* 2005;14(4):360-364. doi:10.1007/s00167-005-0689-7.
8. Belk JW, Wharton BR, Houck DA, et al. Shoulder Stabilization Versus Immobilization for First-Time Anterior Shoulder Dislocation: A Systematic Review and Meta-analysis of Level 1 Randomized Controlled Trials. *Am J Sports Med.* 2023;51(6):1634-1643. doi:10.1177/03635465211065403.
9. Berk AN, Cregar WM, Gachigi KK, et al. Outcomes of subacromial balloon spacer implantation for irreparable rotator cuff tears: a systematic review and meta-analysis. *J Shoulder Elbow Surg.* 2023;32(10):2180-2191. doi:10.1016/j.jse.2023.04.016.
10. Bi AS, Anil U, Colasanti CA, et al. Comparison of Multiple Surgical Treatments for Massive Irreparable Rotator Cuff Tears in Patients Younger Than 70 Years of Age: A Systematic Review and Network Meta-analysis. *Am J Sports Med.* 2024;52(11):2919-2930. doi:10.1177/03635465231204623.
11. Boileau P, Baque F, Valerio L, et al. Isolated Arthroscopic Biceps Tenotomy or Tenodesis Improves Symptoms in Patients with Massive Irreparable Rotator Cuff Tears. *J Bone Joint Surg Am.* 2007;89(4):747-757.
12. Bolia IK, Collon K, Bogdanov J, Lan R, Petrigliano FA. Management Options for Shoulder Impingement Syndrome in Athletes: Insights and Future Directions. *Open Access J Sports Med.* 2021;12:43-53. doi:10.2147/OAJSM.S281100.
13. Boorman RS, More KD, Hollinshead RM, et al. What happens to patients when we do not repair their cuff tears? Five-year rotator cuff quality-of-life index outcomes following nonoperative treatment of patients with full-thickness rotator cuff tears. *J Shoulder Elbow Surg.* 2018;27:444-448. doi:10.1016/j.jse.2017.10.009.
14. Brealey S, Northgraves M, Kottam L, et al. Surgical treatments compared with early structured physiotherapy in secondary care for adults with primary frozen shoulder: the UK FROST three-arm RCT. *Health Technol Assess (Rockv).* 2020;24(71):1-162. doi:10.3310/hta24710.
15. Brolin T, Updegrove G, Horneff J. Classifications in Brief: Hamada Classification of Massive Rotator Cuff Tears. *Clin Orthop Relat Res.* 2017;475(11):2819-2823. doi:10.1007/s11999-017-5340-7.
16. Budoff J, Nirschl R, Guidi E. Current Concepts Review-Debridement of Partial-Thickness Tears of the Rotator Cuff without Acromioplasty. Long-term follow-up and review of the literature. *J Bone Joint Surg Am.* 1998;80(5):733-748.
17. Burkhart S, Prancun J, Hartzler R. Superior Capsular Reconstruction for the Operatively Irreparable Rotator Cuff Tear: Clinical Outcomes Are Maintained 2 Years After Surgery. *Arthroscopy.* 2020;36(2):373-380. doi:10.1016/j.arthro.2019.08.035.

18. Burnier M, Elhassan B, Sanchez-Sotelo J. Surgical Management of Irreparable Rotator Cuff Tears. *J Bone Joint Surg*. 2019;101(17):1603-1612. doi:10.2106/jbjs.18.01392.
19. Carver TJ, Kraeutler MJ, Smith JR, Bravman JT, Mccarty EC. Nonarthroplasty Surgical Treatment Options for Massive, Irreparable Rotator Cuff Tears. *Orthop J Sports Med*. 2018;6(11):232596711880538. doi:10.1177/2325967118805385.
20. Cvetanovich G, Waterman B, Verma N, Romeo A. Management of the Irreparable Rotator Cuff Tear. *JAAOS*. 2019;27(24):909-917. doi:10.5435/jaaos-d-18-00199.
21. Choi L. Overuse injuries. In: DeLee J, et al. *DeLee and Drez's Orthopaedic Sports Medicine*. 3rd ed. Philadelphia, Pa. Saunders Elsevier. 2009.
22. Cofield RH. Subscapular muscle transposition for repair of chronic rotator cuff tears. *Surg Gynecol Obstet*. 1982;154(5):667-672.
23. Coory J, Parr A, Wilkinson M, Gupta A. Efficacy of suprascapular nerve block compared with subacromial injection: a randomized controlled trial in patients with rotator cuff tears. *J Shoulder Elbow Surg*. 2019;28(3):430-436. doi:10.1016/j.jse.2018.11.051.
24. Cvetanovich G, Waterman B, Verma N, Romeo A. Management of the irreparable rotator cuff tear. *J Am Acad Orthop Surg*. 2019;27(24):909-917. doi:10.5435/jaaos-d-18-00199.
25. Denard PJ, Brady PC, Adams CR, et al. Preliminary results of arthroscopic superior capsule reconstruction with dermal allograft. *Arthroscopy*. 2018;34(1):93-99.
26. DeOrio JK, Cofield RH. Results of a second attempt at surgical repair of a failed initial rotator-cuff repair. *J Bone Joint Surg Am*. 1984;66(4):563-567.
27. Dunn WR, Kuhn JE, Sanders R, et al. 2013 Neer Award: predictors of failure of nonoperative treatment of chronic, symptomatic, full-thickness rotator cuff tears. *J Shoulder Elbow Surg*. 2016;25:1303-1311.
28. Ernstbrunner L, Suter A, Catanzaro S, Rahm S, Gerber C. Reverse Total Shoulder Arthroplasty for Massive, Irreparable Rotator Cuff Tears Before the Age of 60 Years. *J Bone Joint Surg*. 2017;99(20):1721-1729. doi:10.2106/jbjs.17.00095.
29. Forsythe B, Agarwalla A, Puzitiello RN, Sumner S, Romeo AA, Mascarenhas R. The Timing of Injections Prior to Arthroscopic Rotator Cuff Repair Impacts the Risk of Surgical Site Infection. *J Bone Joint Surg*. 2019;101(8):682-687. doi:10.2106/jbjs.18.00631.
30. Galatz LM, Ball C, Teefey S, et al. The outcome and repair integrity of completely arthroscopically repaired large and massive rotator cuff tears. *J Bone Joint Surg Am*. 2004;86(2):219-224.
31. Galvin J, Kenney R, Curry E, et al. Superior Capsular Reconstruction for Massive Rotator Cuff Tears. *JBJS Rev*. 2019;7(6):e1. doi:10.2106/jbjs.rvw.18.00072.
32. Garofalo R, Conti M, Massazza G, Cesari E, Vinci E, Castagna A. Subcoracoid impingement syndrome: a painful shoulder condition related to different pathologic factors. *Musculoskelet Surg*. 2011;95 Suppl 1:S25-S29.
33. Garrigues G. The Superior Capsule Reconstruction. *J Bone Joint Surg*. 2019;101(21):e116. doi:10.2106/jbjs.19.00883.
34. Gartsman G, Taverna E: The incidence of glenohumeral joint abnormalities associated with complete, reparable rotator cuff tears. *Arthroscopy*. 1997;12:575-579.
35. Garving C, Jakob S, Bauer I, Nadjar R, Brunner UH. Impingement Syndrome of the Shoulder. *Dtsch Arztebl Int*. 2017;114(45):765-776. doi:10.3238/arztebl.2017.0765.
36. Godeneche A, Elia F, Kempf J, et al. Fatty infiltration of stage 1 or higher significantly compromises long-term healing of supraspinatus repairs. *J Shoulder Elbow Surg*. 2017;26:1818-1825. doi:10.1016/j.jse.2017.03.024.
37. Green A. Is the Devil in the Details?. *J Bone Joint Surg*. 2019;101(21):e117. doi:10.2106/jbjs.19.01120.
38. Guevara J, Entezari V, Ho J, Derwin K, Iannotti J, Ricchetti E. An Update on Surgical Management of the Repairable Large-to-Massive Rotator Cuff Tear. *J Bone Joint Surg*. 2020;102(19):1742-1754. doi:10.2106/jbjs.20.00177.
39. Gupta AK, Hug K, Berkoff DJ, et al. Dermal Tissue Allograft for the Repair of Massive Irreparable Rotator Cuff Tears. *Am J Sports Med*. 2012;40(1):141-147. doi:10.1177/0363546511422795.
40. Hamada K, Fukuda h, Mikasa M, Kobayashi Y. Roentgenographic Findings in Massive Rotator Cuff Tears: A Long-Term Observation. *Clin Orthop Relat Res*. 1990;(254):92-96. doi:10.1097/00003086-199005000-00014.
41. Hartzler R, Burkhart S. Superior Capsular Reconstruction. *Orthopedics*. 2017;40(5):271-280. doi:10.3928/01477447-20170920-02.

42. Herbery-Davies J, Teefey SA, Steger-May K, et al. Progression of fatty muscle degeneration in atraumatic rotator cuff tears. *J Bone Joint Surg Am.* 2017;99:832-839. doi:10.2106/JBJS.16.00030.
43. Hippensteel KJ, Brophy R, Smith MV, et al. A comprehensive review of physical examination tests of the cervical spine, scapula, and rotator cuff. *J Am Acad Orthop Surg.* 2019;27:385-94. doi:10.5435/JAAOS-D-17-00090.
44. Hirahara AM, Andersen WJ, Panero AJ. Superior capsular reconstruction: clinical outcomes after minimum 2-year follow-up. *Am J Orthop.* 2017;266-278.
45. Hovelius L, Olofsson A, Sandström B. Nonoperative treatment of primary anterior shoulder dislocation in patients forty years of age and younger. a prospective twenty-five-year follow-up. *J Bone Joint Surg Am.* 2008;90(5):945-952.
46. Hovis W, Dean M, Mallon W, Hawkins R. Posterior instability of the shoulder with secondary impingement in elite golfers. *Am J Sports Med.* 2002;30(6):886-890.
47. Hsieh LF, Kuo YC, Huang YH, Liu YF, Hsieh TL. Comparison of corticosteroid injection, physiotherapy and combined treatment for patients with chronic subacromial bursitis - A randomised controlled trial. *Clin Rehabil.* 2023;37(9):1189-1200. doi:10.1177/02692155231166220.
48. Jain N, Ayers G, Fan R, et al. Comparative Effectiveness of Operative Versus Nonoperative Treatment for Rotator Cuff Tears: A Propensity Score Analysis from the ROW Cohort. *Am J Sports Med.* 2019;47(13):3065-3072. doi:10.1177/0363546519873840.
49. Jeong J, Kim S, Yoon T, Eum K, Chun Y. Arthroscopic Repair of Large and Massive Rotator Cuff Tears. *J Bone Joint Surg.* 2020;102(14):1248-1254. doi:10.2106/jbjs.19.01014.
50. Johns WL, Ailaney N, Lacy K, Golladay GJ, Vanderbeck J, Kalore NV. Implantable Subacromial Balloon Spacers in Patients with Massive Irreparable Rotator Cuff Tears: A Systematic Review of Clinical, Biomechanical, and Financial Implications. *Arthrosc Sports Med Rehabil.* 2020;2(6):e855-e872. doi:10.1016/j.asmr.2020.06.011.
51. Kalson N, Borthwick L, Mann D, et al. International consensus on the definition and classification of fibrosis of the knee joint. *Bone Joint J.* 2016;98-B(11):1479-1488. doi:10.1302/0301-620x.98b10.37957.
52. Karjalainen TV, Jain NB, Page CM, et al. Subacromial decompression surgery for rotator cuff disease. *Cochrane Database Syst Rev.* 2019;1:CD005619.
53. Keener JD, Galatz LM, Teefey SA, et al. A Prospective Evaluation of Survivorship of Asymptomatic Degenerative Rotator Cuff Tears. *J Bone Joint Surg Am.* 2015;97:89-98.
54. Keener JD, Patterson BM, Orvets N, Chamberlain AM. Degenerative Rotator Cuff Tears. *J Am Acad Orthop Surg.* 2019;27(5):156-165. doi:10.5435/jaaos-d-17-00480.
55. Khazzam M, Kane SM, Smith MJ. Open Shoulder Stabilization Using bone block technique for treatment of chronic glenohumeral instability associated with glenoid deficiency. *Am J Orthop.* 2009;38(7):329-335.
56. Kibler WB, Sciascia A, Hester P, Dome D, Jacobs C. Clinical Utility of Traditional and New Tests in the Diagnosis of Biceps Tendon Injuries and Superior Labrum Anterior and Posterior Lesions in the Shoulder. *Am J Sports Med.* 2009;37(9):1840-1847. doi:10.1177/0363546509332505.
57. Kim H, Teefey S, Zelig A, et al. Shoulder strength in asymptomatic individuals with intact compared with torn rotator cuffs. *J Bone Joint Surg Am.* 2009;91(2):289-296.
58. Kim SJ, Choi YR, Jung M, Yoon Y, Chun YM. Concomitant coracoplasty during arthroscopic subscapularis repair does not yield better clinical outcomes and structural integrity. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(1):56-62.
59. Kragh JF Jr, Doukas WC, Basamania CJ. Primary coracoid impingement syndrome. *Am J Orthop.* 2004;33(5):229-232.
60. Kuhn JE, Dunn WR, Sanders R, et al. Effectiveness of physical therapy in treating atraumatic full-thickness rotator cuff tears: a multicenter prospective cohort study. *J Shoulder Elbow Surg.* 2013;22(10):1371-1379. doi:10.1016/j.jse.2013.01.026.
61. Kuhn JE, Dunn WR, Sanders R, et al. 2024 Kappa Delta Ann Doner Vaughan Award: Nonsurgical Treatment of Symptomatic, Atraumatic Full-Thickness Rotator Cuff Tears-a Prospective Multicenter Cohort Study With 10-Year Follow-Up. *J Am Acad Orthop Surg.* 2024;32(23):1061-1073. doi:10.5435/JAAOS-D-24-00841.
62. Kuhn JE, Dunn WR, Sanders R, et al. The Predictors of Surgery for Symptomatic, Atraumatic Full-Thickness Rotator Cuff Tears Change Over Time: Ten-Year Outcomes of the MOON Shoulder Prospective Cohort. *J Bone Joint Surg Am.* 2024;106(17):1563-1572. doi:10.2106/jbjs.23.00978.

63. Kukkonen J, Joukainen A, Lehtinen J, et al. Treatment of Nontraumatic Rotator Cuff Tears: A Randomized Control Trial with Two Years of Clinical and Imaging Follow-up. *J Bone Joint Surg Am.* 2015;97:1729-1737.
64. Kukkonen J, Ryösä A, Joukainen A, et al. Operative versus conservative treatment of small, nontraumatic supraspinatus tears in patients older than 55 years: over 5-year follow-up of a randomized controlled trial. *J Shoulder Elbow Surg.* 2021;30(11):2455-2464. doi:10.1016/j.jse.2021.03.133.
65. Lee SJ, Min YK. Can inadequate acromiohumeral distance improvement and poor posterior remnant tissue be the predictive factors of re-tear? Preliminary outcomes of arthroscopic superior capsular reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(7):2205-2213.
66. Leland D, Bernard C, Keyt L, et al. An Age-Based Approach to Anterior Shoulder Instability in Patients Under 40 Years Old: Analysis of a US Population. *Am J Sports Med.* 2019;48(1):56-62. doi:10.1177/0363546519886861.
67. Leroux, TS, Saltzman BM, Meyer M, et al. The Influence of Evidence-Based Surgical Indications and Techniques on Failure Rates After Arthroscopic Shoulder Stabilization in the Contact or Collision Athlete with Anterior Shoulder Instability. *Am J Sports Med.* 2017;45(5):1218-1225.
68. Lewington MR, Ferguson DP, Smith TD, Burks R, Coady C, Wong IH-B. Graft Utilization in the Bridging Reconstruction of Irreparable Rotator Cuff Tears: A Systematic Review. *Am J Sports Med.* 2017;45(13):3149-3157. doi:10.1177/0363546517694355.
69. Lim S, Alramadhan H, Kwak J-M, Hong H, Jeon I-H. Graft tears after arthroscopic superior capsule reconstruction (ASCR): pattern of failure and its correlation with clinical outcome. *Arch Orthop Trauma Surg.* 2018;139(2):231-239. doi:10.1007/s00402-018-3025-7.
70. Lin J, Sun Y, Chen Q, Liu S, Ding Z, Chen J. Outcome Comparison of Graft Bridging and Superior Capsule Reconstruction for Large to Massive Rotator Cuff Tears: A Systematic Review. *Am J Sports Med.* 2019;48(11):2828-2838. doi:10.1177/0363546519889040.
71. MacDonald P, McRae S, Old J, et al. Arthroscopic Bankart repair with and without arthroscopic infraspinatus remplissage in anterior shoulder instability with a Hill-Sachs defect: a randomized controlled trial. *J Shoulder Elbow Surg.* 2021;30(6):1288-1298. doi:10.1016/j.jse.2020.11.013.
72. MacDonald P, Verhulst F, McRae S, et al. Biceps Tenodesis Versus Tenotomy in the Treatment of Lesions of the Long Head of the Biceps Tendon in Patients Undergoing Arthroscopic Shoulder Surgery: A Prospective Double-Blinded Randomized Controlled Trial. *Am J Sports Med.* 2020;48(6):1439-1449. doi:10.1177/0363546520912212.
73. Maman E, Safran O, Beyth S, et al. Biceps Tenotomy Does Not Affect the Functional Outcomes of Patients Treated with Spacer Implantation Due to Massive Irreparable Rotator Cuff Tears. *Open Orthop J.* 2017;11:1577-1584. doi:10.2174/1874325001711011577.
74. Mantripragada S, Peh WC. Spectrum of lesions of the acromioclavicular joint: imaging features. *Singapore Med J.* 2022;63(3):130-139. doi:10.11622/smedj.2022046.
75. McKee M, Yoo D. The effect of surgery for rotator cuff disease on general health status. Results of a prospective trial. *J Bone Joint Surg Am.* 2000;82(7):970-979.
76. McKernan M, Schickendantz M, Frangiamore S. Diagnosis and Management of Subcoracoid Impingement. *J Am Acad Orthop Surg.* 2020;29(3):100-107. doi:10.5435/jaaos-d-20-00008.
77. Metcalfe A, Parsons H, Parsons N, et al. Subacromial balloon spacer for irreparable rotator cuff tears of the shoulder (START:REACTS): a group-sequential, double-blind, multicentre randomised controlled trial. *Lancet.* 2022;399(10339):1954-1963. doi:10.1016/s0140-6736(22)00652-3.
78. Mihata T, Lee T, Hasegawa A, et al. Five-Year Follow-up of Arthroscopic Superior Capsule Reconstruction for Irreparable Rotator Cuff Tears. *J Bone Joint Surg.* 2019;101(21):1921-1930. doi:10.2106/jbjs.19.00135.
79. Mihata T, Lee TQ, Watanabe C, et al. Clinical results of arthroscopic superior capsule reconstruction for irreparable rotator cuff tears. *Arthroscopy.* 2013;29(3):459-470. doi:10.1016/j.arthro.2012.10.022.
80. Mishra D, Fanton G. Two-year outcome of arthroscopic Bankart repair and electrothermal-assisted capsulorrhaphy for recurrent traumatic anterior shoulder instability. *Arthroscopy.* 2001;17(8):844-849.
81. Moon AS, Patel HA, Ithurburn MP, Brabston EW, Ponce BA, Momaya AM. Subacromial Spacer Implantation for the Treatment of Massive Irreparable Rotator Cuff Tears: A Systematic Review. *Arthroscopy.* 2019;35(2):607-614. doi:10.1016/j.arthro.2018.08.006.
82. Moosmayer S, Gartner AV, Tariq R. The natural course of nonoperatively treated rotator cuff tears: an 8.8 year follow-up of tear anatomy and clinical outcome in 49 patients. *J Shoulder Elbow Surg.* 2017;26:627-634. doi:10.1016/j.jse.2016.10.002.

83. Moosmayer S, Lund G, Seljom U, et al. At a 10-Year Follow-up, Tendon Repair Is Superior to Physiotherapy in the Treatment of Small and Medium-Sized Rotator Cuff Tears. *J Bone Joint Surg*. 2019;101(12):1050-1060. doi:10.2106/jbjs.18.01373.
84. Nakhaei Amroodi M, Salariyeh M. Predictors of failure after conservative treatment of symptomatic partial-thickness rotator cuff tear. *J Shoulder Elbow Surg*. 2020;29(1):113-120. doi:10.1016/j.jse.2019.05.009.
85. Neumann JA, Zgonis MH, Rickert KD, et al. Interposition Dermal Matrix Xenografts: A Successful Alternative to Traditional Treatment of Massive Rotator Cuff Tears. *Am J Sports Med*. 2017;45(6):1261-1268. doi:10.1177/0363546516683945.
86. Ono Y, LeBlanc J, Bois AJ, et al. Graft Healing Is More Important Than Graft Technique: Superior Capsular Reconstruction Versus Bridging Grafts—A Prospective Randomized Controlled Trial. *Arthroscopy*. 2022;38(12):3109-3117. doi:10.1016/j.arthro.2022.06.033.
87. Park M, Jun B, Park C, et al. Biomechanical Analysis of a Knotless Transtendon Interimplant Mattress Repair for Partial-Thickness Articular-Sided Rotator Cuff Tears. *Am J Sports Med*. 2009;37(12):2427-2434.
88. Park JY, Lhee SH, Oh KS, Kim NR, Hwang JT. Is arthroscopic coracoplasty necessary in subcoracoid impingement syndrome? *Arthroscopy*. 2012;28(12):1766-1775.
89. Pennington WT, Bartz BA, Pauli JM, et al. Arthroscopic superior capsular reconstruction with acellular dermal allograft for the treatment of massive irreparable rotator cuff tears, short-term clinical outcomes and the radiographic parameter of superior capsular distance. *Arthroscopy*. 2018;34(6):1764-1773.
90. Petretera A, Dwyer T, Tsuji MRS, et al. Outcomes of Arthroscopic Bankart Repair in Collision Versus Noncollision Athletes. *Orthopedics*. 2013;36(5):e621-e626.
91. Petri M, Greenspoon J, Moulton S, Millett P. Patch-Augmented Rotator Cuff Repair and Superior Capsule Reconstruction. *Open Orthop J*. 2016;10(1):315-323. doi:10.2174/1874325001610010315.
92. Piekaar RSM, Bouman ICE, van Kampen PM, van Eijk F, Huijsmans PE. The subacromial balloon spacer for massive irreparable rotator cuff tears: approximately 3 years of prospective follow-up. *Musculoskelet Surg*. 2020;104(2):207-214. doi:10.1007/s12306-019-00614-1.
93. Plancher K, Shanmugam J, Briggs K, Petterson S. Diagnosis and Management of Partial Thickness Rotator Cuff Tears: A Comprehensive Review. *JAAOS*. 2021;29(24):1031-1043. doi:10.5435/jaaos-d-20-01092.
94. Ranebo MC, Hallgren HCB, Norlin R, Adolfsson LE. Long-term clinical and radiographic outcome of rotator cuff repair with a synthetic interposition graft: a consecutive case series with 17 to 20 years of follow-up. *J Shoulder Elbow Surg*. 2018;27(9):1622-1628. doi:10.1016/j.jse.2018.03.011.
95. Raval P, Gibbs VN, Pandey R. Preoperative partial-thickness rotator cuff tears do not compromise anatomic total shoulder replacement outcomes: medium-term follow-up. *J Shoulder Elbow Surg*. 2021;30(4):871-876. doi:10.1016/j.jse.2020.07.037. Epub 2020 Aug 7.
96. Rees J. The pathogenesis and surgical treatment of tears of the rotator cuff. *J Bone Joint Surg Br*. 2008;90-B(7):827-832.
97. Rhon DI, Boyles RB, Cleland JA. One-year outcome of subacromial corticosteroid injection compared with manual physical therapy for the management of the unilateral shoulder impingement syndrome: a pragmatic randomized trial. *Ann Intern Med*. 2014;161(3):161-169.
98. Rosas S, Krill M, Amoo-Achampong K, Kwon K, Nwachukwu B, McCormick F. A practical, evidence-based, comprehensive (PEC) physical examination for diagnosing pathology of the long head of the biceps. *J Shoulder Elbow Surg*. 2017;26(8):1484-1492. doi:10.1016/j.jse.2017.03.002.
99. Sachs RA, Lin D, Stone ML, et al. Can the Need for Future Surgery for Acute Traumatic Anterior Shoulder Dislocation Be Predicted? *J Bone Joint Surg Am*. 2007;89:1665-1674.
100. Sanchez-Sotelo J, Cofield RH, Rowland CM. Shoulder Hemiarthroplasty for Glenohumeral Arthritis Associated with Severe Rotator Cuff Deficiency. *J Bone Joint Surg Am*. 2001 Dec;83(12):1814-1822. doi:10.2106/00004623-200112000-00008.
101. Senekovic V, Poberaj B, Kovacic L, Mikek M, Adar E, Dekel A. Prospective clinical study of a novel biodegradable sub-acromial spacer in treatment of massive irreparable rotator cuff tears. *Eur J Orthop Surg Traumatol*. 2013;23(3):311-316. doi:10.1007/s00590-012-0981-4.
102. Shen P, Lien S, Shen H, et al. Long-term functional outcomes after repair of rotator cuff tears correlated with atrophy of the supraspinatus muscles on magnetic resonance images. *J Shoulder Elbow Surg*. 2008;17(1 Suppl):1S-7S.

103. Sochacki KR, Mcculloch PC, Lintner DM, Harris JD. Superior Capsular Reconstruction for Massive Rotator Cuff Tear Leads to Significant Improvement in Range of Motion and Clinical Outcomes: A Systematic Review. *Arthroscopy*. 2019;35(4):1269-1277. doi:10.1016/j.arthro.2018.10.129.
104. Song A, DeClercq J, Ayers G, et al. Comparative Time to Improvement in Nonoperative and Operative Treatment of Rotator Cuff Tears. *J Bone Joint Surg*. 2020;102(13):1142-1150. doi:10.2106/jbjs.19.01112.
105. Stewart RK, Kaplin L, Parada SA, Graves BR, Verma NN, Waterman BR. Outcomes of Subacromial Balloon Spacer Implantation for Massive and Irreparable Rotator Cuff Tears: A Systematic Review. *Orthop J Sports Med*. 2019;7(10):2325967119875717. doi:10.1177/2325967119875717.
106. Streubel PN, Krych AJ, Simone JP, et al. Anterior Glenohumeral Instability: A Pathology-based Surgical Treatment Strategy. *J Am Acad Orthop Surg*. 2014;22(5):283-294.
107. Suenaga N, Minami A, Kaneda K. Postoperative subcoracoid impingement syndrome in patients with rotator cuff tear. *J Shoulder Elbow Surg*. 2000;9(4):275-278.
108. Tokish J, Makovicka J. The Superior Capsular Reconstruction: Lessons Learned and Future Directions. *J Am Acad Orthop Surg*. 2020;28(13):528-537. doi:10.5435/jaaos-d-19-00057.
109. Ueda Y, Sugaya H, Takahashi N, et al. Rotator Cuff Lesions in Patients with Stiff Shoulders: A Prospective Analysis of 379 Shoulders. *J Bone Joint Surg Am*. 2015;97:1233-1237.
110. van der List JP, Kok LM, Alta TDW, van der List MPJ, van Noort A. Role of Delay Between Injury and Surgery on the Outcomes of Rotator Cuff Repair: A Systematic Review and Meta-analysis. *Am J Sports Med*. 2023;51(5):1328-1339. doi:10.1177/03635465211069560.
111. Venouziou AI, Kokkalis ZT, Sotereanos DG. Human dermal allograft interposition for the reconstruction of massive irreparable rotator cuff tears. *Am J Orthop*. 2013;42(2):63-70.
112. Verma N, Srikumaran U, Roden CM, et al. InSpace Implant Compared with Partial Repair for the Treatment of Full-Thickness Massive Rotator Cuff Tears: A Multicenter, Single-Blinded, Randomized Controlled Trial. *J Bone Joint Surg Am*. 2022;104(14):1250-1262. doi:10.2106/JBJS.21.00667.
113. Vitale M, Arons R, Hurwitz S, et al. The Rising Incidence of Acromioplasty. *J Bone Joint Surg*. 2010;92(9):1842-1850.
114. Welch M, Rankin S, How Saw Keng M, Woods D. A systematic review of the treatment of primary acromioclavicular joint osteoarthritis. *Shoulder Elbow*. 2024;16(2):129-144. doi:10.1177/17585732231157090.
115. Werner BC, Brockmeier SF, Miller MD. Etiology, Diagnosis, and Management of Failed SLAP Repair. *J Am Acad Orthop Surg*. 2014;22(9):554-565.
116. Whittle S, Buchbinder R. In the Clinic: Rotator Cuff Disease. *Ann Intern Med*. 2015;162(1):ITC1-ITC15. doi:10.7326/AITC201501060.
117. Woodmass JM, McRae S, Lapner P, et al. Arthroscopic Bankart repair with remplissage in anterior shoulder instability results in fewer redislocations than Bankart repair alone at medium-term follow-up of a randomized controlled trial. *Am J Sports Med*. 2024;52(8):2055-2062. doi:10.1177/03635465241254063.
118. Work Loss Data Institute. *Shoulder (acute and chronic)*. Corpus Christi (TX): Work Loss Data Institute; 2008.
119. Wright MA, Keener JD, Chamberlain AM. Comparison of Clinical Outcomes After Anatomic Total Shoulder Arthroplasty and Reverse Shoulder Arthroplasty in Patients 70 Years and Older with Glenohumeral Osteoarthritis and an Intact Rotator Cuff. *J Am Acad Orthop Surg*. 2020;28(5):e222-e229. doi:10.5435/jaaos-d-19-00166.
120. Wylie JD, Suter T, Potter MQ, et al. Mental Health Has a Stronger Association with Patient-Reported Shoulder Pain and Function Than Tear Size in Patients with Full-Thickness Rotator Cuff Tears. *J Bone Joint Surg Am*. 2016;98:251-256.
121. Yapp L, Nicholson J, Robinson C. Primary Arthroscopic Stabilization for a First-Time Anterior Dislocation of the Shoulder. *J Bone Joint Surg*. 2019;102(6):460-467. doi:10.2106/jbjs.19.00858.